



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 03:13 AM GMT

PDB ID : 2IF8
Title : Crystal structure of Inositol Phosphate Multikinase Ipk2 in complex with ADP and Mn²⁺ from *S. cerevisiae*
Authors : Holmes, W.; Jogl, G.
Deposited on : 2006-09-20
Resolution : 2.40 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| R_{free} | 66092 | 2207 (2.40-2.40) |
| Clashscore | 79885 | 2789 (2.40-2.40) |
| Ramachandran outliers | 78287 | 2736 (2.40-2.40) |
| Sidechain outliers | 78261 | 2737 (2.40-2.40) |
| RSRZ outliers | 66119 | 2210 (2.40-2.40) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | A | 363 |  |
| 1 | B | 363 |  |

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4568 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inositol polyphosphate multikinase.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|---------|-------|
| 1 | A | 255 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 2083 | 1338 | 335 | 400 | 10 | | | |
| 1 | B | 256 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 2091 | 1342 | 336 | 403 | 10 | | | |

There are 16 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|------------------|------------|
| A | 356 | LEU | - | CLONING ARTIFACT | UNP P07250 |
| A | 357 | GLU | - | CLONING ARTIFACT | UNP P07250 |
| A | 358 | HIS | - | EXPRESSION TAG | UNP P07250 |
| A | 359 | HIS | - | EXPRESSION TAG | UNP P07250 |
| A | 360 | HIS | - | EXPRESSION TAG | UNP P07250 |
| A | 361 | HIS | - | EXPRESSION TAG | UNP P07250 |
| A | 362 | HIS | - | EXPRESSION TAG | UNP P07250 |
| A | 363 | HIS | - | EXPRESSION TAG | UNP P07250 |
| B | 356 | LEU | - | CLONING ARTIFACT | UNP P07250 |
| B | 357 | GLU | - | CLONING ARTIFACT | UNP P07250 |
| B | 358 | HIS | - | EXPRESSION TAG | UNP P07250 |
| B | 359 | HIS | - | EXPRESSION TAG | UNP P07250 |
| B | 360 | HIS | - | EXPRESSION TAG | UNP P07250 |
| B | 361 | HIS | - | EXPRESSION TAG | UNP P07250 |
| B | 362 | HIS | - | EXPRESSION TAG | UNP P07250 |
| B | 363 | HIS | - | EXPRESSION TAG | UNP P07250 |


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 2 | A | 1 | Total | Mn | 0 | 0 |
| | | | 1 | 1 | | |

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---------|---------|
| 3 | A | 1 | Total | Ca | 0 | 0 |
| | | | 1 | 1 | | |

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|----|---|----|---|---------|---------|
| 4 | A | 1 | Total | C | N | O | P | 0 | 0 |
| | | | 27 | 10 | 5 | 10 | 2 | | |

- Molecule 5 is water.


| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|-------|-----|---------|---------|
| 5 | A | 212 | Total | O | 0 | 0 |
| | | | 212 | 212 | | |
| 5 | B | 153 | Total | O | 0 | 0 |
| | | | 153 | 153 | | |

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Inositol polyphosphate multikinase

Chain A:



- Molecule 1: Inositol polyphosphate multikinase

Chain B:



4 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 63 | Depositor |
| Cell constants a, b, c, α , β , γ | 185.00Å 185.00Å 50.00Å 90.00° 90.00° 120.00° | Depositor |
| Resolution (Å) | 30.00 – 2.40 26.98 – 2.40 | Depositor EDS |
| % Data completeness (in resolution range) | 98.5 (30.00-2.40) 98.6 (26.98-2.40) | Depositor EDS |
| R_{merge} | 0.09 | Depositor |
| R_{sym} | (Not available) | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.80 (at 2.39Å) | Xtriage |
| Refinement program | REFMAC 5.2.0019 | Depositor |
| R, R_{free} | 0.193 , 0.246 0.194 , 0.241 | Depositor DCC |
| R_{free} test set | 1922 reflections (5.29%) | DCC |
| Wilson B-factor (Å ²) | 41.0 | Xtriage |
| Anisotropy | 0.299 | Xtriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.38 , 69.8 | EDS |
| Estimated twinning fraction | 0.047 for h,-h-k,-l | Xtriage |
| L-test for twinning | $\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$ | Xtriage |
| Outliers | 0 of 38288 reflections | Xtriage |
| F_o, F_c correlation | 0.95 | EDS |
| Total number of atoms | 4568 | wwPDB-VP |
| Average B, all atoms (Å ²) | 47.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------------|-------------|---------------|
| | | RMSZ | $\# Z > 5$ | RMSZ | $\# Z > 5$ |
| 1 | A | 0.68 | 1/2122 (0.0%) | 0.73 | 0/2854 |
| 1 | B | 0.58 | 0/2130 | 0.70 | 1/2865 (0.0%) |
| All | All | 0.63 | 1/4252 (0.0%) | 0.71 | 1/5719 (0.0%) |

All (1) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|-------|-------------|----------|
| 1 | A | 168 | CYS | CB-SG | -7.57 | 1.69 | 1.82 |

All (1) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|------|-------------|----------|
| 1 | B | 38 | LEU | CA-CB-CG | 5.38 | 127.68 | 115.30 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 2083 | 0 | 2069 | 25 | 0 |
| 1 | B | 2091 | 0 | 2073 | 38 | 0 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 2 | A | 1 | 0 | 0 | 0 | 0 |
| 3 | A | 1 | 0 | 0 | 0 | 0 |
| 4 | A | 27 | 0 | 12 | 0 | 0 |
| 5 | A | 212 | 0 | 0 | 4 | 1 |
| 5 | B | 153 | 0 | 0 | 9 | 0 |
| All | All | 4568 | 0 | 4154 | 62 | 1 |

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below.

| Atom-1 | Atom-2 | Distance(Å) | Clash(Å) |
|-----------------|-----------------|-------------|----------|
| 1:B:25:ASP:HB2 | 5:B:496:HOH:O | 1.28 | 1.25 |
| 1:B:197:PHE:HB3 | 5:B:487:HOH:O | 1.38 | 1.23 |
| 1:A:317:SER:N | 5:A:561:HOH:O | 1.87 | 1.05 |
| 1:A:147:LYS:HB3 | 5:A:520:HOH:O | 1.70 | 0.91 |
| 1:A:174:LYS:HE2 | 1:A:189:GLU:OE1 | 1.95 | 0.67 |

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1 | Atom-2 | Distance(Å) | Clash(Å) |
|---------------|----------------------|-------------|----------|
| 5:A:562:HOH:O | 5:A:590:HOH:O[4_665] | 1.90 | 0.30 |

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|---------|----------|-------------|----|
| 1 | A | 247/363 (68%) | 241 (98%) | 3 (1%) | 3 (1%) | 19 | 26 |
| 1 | B | 248/363 (68%) | 241 (97%) | 5 (2%) | 2 (1%) | 27 | 39 |
| All | All | 495/726 (68%) | 482 (97%) | 8 (2%) | 5 (1%) | 22 | 32 |

All (5) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 178 | VAL |
| 1 | B | 179 | LEU |
| 1 | B | 325 | ASP |
| 1 | A | 177 | SER |
| 1 | A | 325 | ASP |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1 | A | 234/327 (72%) | 222 (95%) | 12 (5%) | 33 | 50 |
| 1 | B | 235/327 (72%) | 217 (92%) | 18 (8%) | 18 | 28 |
| All | All | 469/654 (72%) | 439 (94%) | 30 (6%) | 25 | 37 |

5 of 30 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 27 | LEU |
| 1 | B | 73 | LEU |
| 1 | B | 331 | ILE |
| 1 | B | 45 | GLN |
| 1 | B | 113 | GLN |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 45 | GLN |
| 1 | B | 45 | GLN |
| 1 | B | 223 | HIS |

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|-------------|-------------|------|-------------|
| | | | | | Counts | RMSZ | $\# Z > 2$ | Counts | RMSZ | $\# Z > 2$ |
| 4 | ADP | A | 401 | 2 | 29,29,29 | 1.27 | 4 (13%) | 45,45,45 | 1.73 | 7 (15%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|------------|---------|
| 4 | ADP | A | 401 | 2 | - | 0/16/32/32 | 0/1/3/3 |

All (4) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|--------|-------|-------------|----------|
| 4 | A | 401 | ADP | C5-C4 | 3.52 | 1.48 | 1.40 |
| 4 | A | 401 | ADP | C4-N9 | -2.67 | 1.33 | 1.37 |
| 4 | A | 401 | ADP | PB-O3A | 2.53 | 1.64 | 1.60 |
| 4 | A | 401 | ADP | PA-O3A | 2.48 | 1.64 | 1.59 |

The worst 5 of 7 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|------------|-------|-------------|----------|
| 4 | A | 401 | ADP | N3-C2-N1 | -6.06 | 123.64 | 128.71 |
| 4 | A | 401 | ADP | N3-C4-N9 | 4.88 | 134.24 | 125.43 |
| 4 | A | 401 | ADP | O4'-C1'-N9 | 4.70 | 112.81 | 108.44 |
| 4 | A | 401 | ADP | C5-C4-N3 | -2.81 | 119.57 | 125.70 |
| 4 | A | 401 | ADP | C4-C5-N7 | -2.77 | 107.15 | 109.52 |

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | OWAB(Å ²) | Q<0.9 |
|-----|-------|---------------|--------|---------------|-----------------------|-------|
| 1 | A | 255/363 (70%) | -0.00 | 16 (6%) 19 18 | 29, 39, 62, 97 | 0 |
| 1 | B | 256/363 (70%) | 0.51 | 27 (10%) 7 6 | 34, 49, 77, 96 | 0 |
| All | All | 511/726 (70%) | 0.25 | 43 (8%) 11 10 | 29, 45, 73, 97 | 0 |

The worst 5 of 43 RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 1 | B | 111 | SER | 8.9 |
| 1 | B | 285 | ILE | 7.8 |
| 1 | B | 286 | ASP | 7.1 |
| 1 | A | 286 | ASP | 6.8 |
| 1 | B | 145 | LEU | 6.7 |

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSR | LLDF | B-factors(\AA^2) | Q<0.9 |
|-----|------|-------|-----|-------|------|-------|-----------------------------|-------|
| 2 | MN | A | 402 | 1/1 | 0.20 | 0.77 | 112,112,112,112 | 0 |
| 4 | ADP | A | 401 | 27/27 | 0.19 | 0.34 | 82,86,95,96 | 0 |
| 3 | CA | A | 403 | 1/1 | 0.06 | -2.00 | 50,50,50,50 | 0 |

6.5 Other polymers ⓘ

There are no such residues in this entry.