



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 08:36 PM GMT

PDB ID : 2IFF
Title : STRUCTURE OF AN ANTIBODY-LYSOZYME COMPLEX: EFFECT OF
A CONSERVATIVE MUTATION
Authors : Chacko, S.; Davies, D.R.
Deposited on : 1994-02-03
Resolution : 2.65 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2700 (2.70-2.62)
Ramachandran outliers	78287	2657 (2.70-2.62)
Sidechain outliers	78261	2657 (2.70-2.62)
RSRZ outliers	66119	2234 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	L	212	
2	H	215	
3	Y	129	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4325 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG1 HYHEL-5 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	212	Total	C	N	O	S	0	0	0
			1635	1014	273	338	10			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	18	LYS	ARG	CONFLICT	GB 1042224
L	26	SER	ASN	CONFLICT	GB 1042224
L	30	ASN	SER	CONFLICT	GB 1042224
L	33	TYR	HIS	CONFLICT	GB 1042224
L	59	VAL	ALA	CONFLICT	GB 1042224
L	79	THR	ALA	CONFLICT	GB 1042224
L	91	GLY	SER	CONFLICT	GB 1042224
L	92	ARG	SER	CONFLICT	GB 1042224
L	93	ASN	HIS	CONFLICT	GB 1042224
L	?	-	TYR	DELETION	GB 1042224
L	111	PRO	GLN	CONFLICT	GB 1042224

- Molecule 2 is a protein called IGG1 HYHEL-5 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	214	Total	C	N	O	S	0	0	0
			1607	1015	263	322	7			

- Molecule 3 is a protein called HEN EGG WHITE LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Y	129	Total	C	N	O	S	0	1	0
			1003	616	192	185	10			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	68	LYS	ARG	CONFLICT	UNP P00698


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	30	Total O 30 30	0	0
4	L	30	Total O 30 30	0	0
4	Y	20	Total O 20 20	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: IGG1 HYHEL-5 FAB (LIGHT CHAIN)



• Molecule 2: IGG1 HYHEL-5 FAB (HEAVY CHAIN)



• Molecule 3: HEN EGG WHITE LYSOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.80Å 74.80Å 79.00Å 90.00° 101.80° 90.00°	Depositor
Resolution (Å)	10.00 – 2.65 43.59 – 2.58	Depositor EDS
% Data completeness (in resolution range)	69.2 (10.00-2.65) 65.0 (43.59-2.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.66 (at 2.58Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	(Not available) , (Not available) 0.160 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 75.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 12927 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4325	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.73	0/1673	1.45	23/2270 (1.0%)
2	H	0.75	0/1652	1.44	17/2257 (0.8%)
3	Y	0.74	0/1028	1.65	32/1387 (2.3%)
All	All	0.74	0/4353	1.49	72/5914 (1.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1
2	H	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 72 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	Y	108	TRP	CD1-CG-CD2	9.15	113.62	106.30
2	H	190	PRO	O-C-N	-9.13	108.09	122.70
3	Y	62	TRP	CD1-CG-CD2	8.96	113.47	106.30
1	L	146	TRP	CD1-CG-CD2	8.95	113.46	106.30
3	Y	28	TRP	CD1-CG-CD2	8.87	113.40	106.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	4	LEU	Mainchain
1	L	169	SER	Mainchain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1635	0	1557	29	0
2	H	1607	0	1547	36	0
3	Y	1003	0	968	10	0
4	H	30	0	0	2	0
4	L	30	0	0	0	0
4	Y	20	0	0	0	0
All	All	4325	0	4072	73	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:H:26:GLY:HA3	4:H:793:HOH:O	1.49	1.10
2:H:187:PRO:O	2:H:190:PRO:HD2	1.79	0.81
1:L:143:ASN:HB3	1:L:195:THR:HB	1.62	0.81
2:H:122:PRO:HB3	2:H:148:TYR:HB3	1.69	0.73
1:L:142:ILE:HD12	1:L:196:HIS:HB2	1.78	0.66

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	210/212 (99%)	187 (89%)	19 (9%)	4 (2%)	12	26
2	H	212/215 (99%)	190 (90%)	18 (8%)	4 (2%)	12	26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	Y	128/129 (99%)	116 (91%)	9 (7%)	3 (2%)	10	21
All	All	550/556 (99%)	493 (90%)	46 (8%)	11 (2%)	11	25

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	55	SER
1	L	197	LYS
2	H	189	SER
2	H	136	ASN
3	Y	22	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	187/187 (100%)	163 (87%)	24 (13%)	6	14
2	H	182/182 (100%)	163 (90%)	19 (10%)	10	21
3	Y	106/105 (101%)	95 (90%)	11 (10%)	10	21
All	All	475/474 (100%)	421 (89%)	54 (11%)	8	17

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	11	LEU
2	H	97	LEU
3	Y	62	TRP
2	H	28	THR
2	H	46	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	155	ASN
2	H	98	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	Y	27	ASN
3	Y	121	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	212/212 (100%)	-0.62	1 (0%) 88 91	5, 23, 51, 73	0
2	H	214/215 (99%)	-0.66	5 (2%) 57 61	3, 20, 47, 83	0
3	Y	129/129 (100%)	-0.80	0 100 100	5, 18, 39, 50	0
All	All	555/556 (99%)	-0.68	6 (1%) 77 81	3, 21, 47, 83	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	135	THR	4.5
2	H	136	ASN	3.7
1	L	211	GLU	3.0
2	H	3	GLN	2.9
2	H	2	VAL	2.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.