



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 03:24 PM GMT

PDB ID : 4IQB
Title : High Resolution Crystal Structure of C.elegans Thymidylate Synthase
Authors : Wilk, P.; Dowiercial, A.; Banaszak, K.; Jarmula, A.; Rypniewski, W.; Rode, W.
Deposited on : 2013-01-11
Resolution : 1.13 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1225 (1.20-1.04)
Clashscore	79885	1354 (1.20-1.04)
Ramachandran outliers	78287	1287 (1.20-1.04)
Sidechain outliers	78261	1285 (1.20-1.04)
RSRZ outliers	66119	1225 (1.20-1.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	SO4	A	401	-	X
2	SO4	B	401	-	X

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 5927 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thymidylate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	0	31	0
			2474	1587	419	448	20			
1	B	288	Total	C	N	O	S	0	34	0
			2501	1608	423	449	21			

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	463	Total 463	O 463	0	0
3	B	479	Total 479	O 479	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Thymidylate synthase

Chain A: 



• Molecule 1: Thymidylate synthase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.33Å 98.37Å 69.16Å 90.00° 111.77° 90.00°	Depositor
Resolution (Å)	29.30 – 1.13 29.35 – 1.13	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.30-1.13) 99.1 (29.35-1.13)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.13Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.114 , 0.139 0.126 , 0.149	Depositor DCC
R_{free} test set	11419 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	9.9	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 228161 reflections	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5927	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.14	12/2620 (0.5%)	1.34	26/3536 (0.7%)
1	B	1.13	8/2660 (0.3%)	1.25	26/3587 (0.7%)
All	All	1.13	20/5280 (0.4%)	1.30	52/7123 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	178	ARG	CZ-NH2	-7.49	1.23	1.33
1	A	154	ASP	CG-OD2	-6.78	1.09	1.25
1	B	129	GLU	CD-OE2	-6.64	1.18	1.25
1	A	197[A]	CYS	CB-SG	-6.04	1.72	1.82
1	A	197[B]	CYS	CB-SG	-6.04	1.72	1.82
1	A	168	ARG	CZ-NH2	-6.01	1.25	1.33
1	B	276	TYR	CE2-CZ	-5.89	1.30	1.38
1	A	83	TRP	CZ3-CH2	-5.66	1.30	1.40
1	B	260	TYR	CB-CG	5.58	1.60	1.51
1	B	220	ASP	N-CA	-5.46	1.35	1.46
1	A	39[A]	GLU	CG-CD	5.37	1.59	1.51
1	A	39[B]	GLU	CG-CD	5.37	1.59	1.51
1	A	169	GLN	CG-CD	5.32	1.63	1.51
1	B	58[A]	SER	CB-OG	-5.32	1.35	1.42
1	B	58[B]	SER	CB-OG	-5.32	1.35	1.42
1	A	105	GLU	CG-CD	5.31	1.59	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	276	TYR	CE2-CZ	-5.30	1.31	1.38
1	B	261	SER	CB-OG	-5.17	1.35	1.42
1	A	296	ASP	CG-OD1	-5.15	1.13	1.25
1	B	44	GLU	CD-OE1	-5.03	1.20	1.25

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69[A]	ARG	NE-CZ-NH1	17.29	128.94	120.30
1	A	69[B]	ARG	NE-CZ-NH1	17.29	128.94	120.30
1	A	178	ARG	NE-CZ-NH2	-16.19	112.20	120.30
1	A	69[A]	ARG	NE-CZ-NH2	-15.16	112.72	120.30
1	A	69[B]	ARG	NE-CZ-NH2	-15.16	112.72	120.30
1	A	177	ARG	NE-CZ-NH2	-12.54	114.03	120.30
1	B	178[A]	ARG	NE-CZ-NH2	-10.25	115.17	120.30
1	B	178[B]	ARG	NE-CZ-NH2	-10.25	115.17	120.30
1	A	34	TYR	CB-CG-CD1	10.10	127.06	121.00
1	B	177	ARG	NE-CZ-NH1	-9.79	115.40	120.30
1	A	34	TYR	CB-CG-CD2	-9.04	115.58	121.00
1	B	206[A]	ASP	CB-CG-OD1	8.76	126.18	118.30
1	B	206[B]	ASP	CB-CG-OD1	8.76	126.18	118.30
1	A	177	ARG	NE-CZ-NH1	8.39	124.49	120.30
1	A	254	LEU	CB-CG-CD2	8.17	124.88	111.00
1	B	220	ASP	CB-CG-OD2	-7.74	111.34	118.30
1	B	282	ARG	NE-CZ-NH1	-7.72	116.44	120.30
1	A	69[A]	ARG	CD-NE-CZ	-7.57	113.01	123.60
1	A	69[B]	ARG	CD-NE-CZ	-7.57	113.01	123.60
1	A	128[A]	ARG	NE-CZ-NH1	6.78	123.69	120.30
1	A	128[B]	ARG	NE-CZ-NH1	6.78	123.69	120.30
1	B	129	GLU	OE1-CD-OE2	-6.72	115.23	123.30
1	A	178	ARG	NH1-CZ-NH2	6.48	126.52	119.40
1	A	128[A]	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	A	128[B]	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	A	34	TYR	CZ-CE2-CD2	-6.16	114.25	119.80
1	A	117	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	B	51[A]	ARG	NE-CZ-NH2	6.02	123.31	120.30
1	B	51[B]	ARG	NE-CZ-NH2	6.02	123.31	120.30
1	B	116	ASP	CB-CG-OD1	5.82	123.54	118.30
1	B	286	ASP	CB-CG-OD1	5.81	123.53	118.30
1	A	125	PHE	CB-CG-CD2	5.77	124.84	120.80
1	A	128[A]	ARG	CG-CD-NE	-5.71	99.82	111.80
1	A	128[B]	ARG	CG-CD-NE	-5.71	99.82	111.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	168[A]	ARG	NE-CZ-NH1	-5.56	117.52	120.30
1	B	168[B]	ARG	NE-CZ-NH1	-5.56	117.52	120.30
1	B	273	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	B	32	TYR	CZ-CE2-CD2	-5.52	114.83	119.80
1	B	246	LYS	CD-CE-NZ	5.51	124.37	111.70
1	B	121	ASP	CB-CG-OD2	5.48	123.23	118.30
1	A	25	VAL	CA-CB-CG1	5.35	118.93	110.90
1	B	207[A]	ASN	CB-CA-C	5.34	121.08	110.40
1	B	207[B]	ASN	CB-CA-C	5.34	121.08	110.40
1	B	215	TYR	CB-CG-CD2	5.24	124.14	121.00
1	A	220	ASP	CB-CG-OD1	5.22	123.00	118.30
1	B	49[A]	ASP	CB-CG-OD2	-5.19	113.63	118.30
1	B	49[B]	ASP	CB-CG-OD2	-5.19	113.63	118.30
1	B	117	ARG	NE-CZ-NH1	-5.16	117.72	120.30
1	B	109[A]	LYS	CD-CE-NZ	-5.08	100.02	111.70
1	B	109[B]	LYS	CD-CE-NZ	-5.08	100.02	111.70
1	A	128[A]	ARG	CB-CA-C	-5.00	100.39	110.40
1	A	128[B]	ARG	CB-CA-C	-5.00	100.39	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	51[A]	ARG	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2474	0	0	14	0
1	B	2501	0	0	30	1
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	463	0	0	3	0
3	B	479	0	0	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5927	0	0	42	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (42) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:A:749:HOH:O	1:B:200[A]:MET:CE	1.95	1.11
1:A:104[B]:MET:CE	1:A:109[B]:LYS:CE	2.28	1.11
1:A:104[B]:MET:CE	1:A:109[B]:LYS:CD	2.32	1.07
1:A:51[A]:ARG:NH2	1:A:51[A]:ARG:CG	2.18	1.03
1:A:51[A]:ARG:CB	1:A:51[A]:ARG:NH2	2.25	0.99
1:B:51[A]:ARG:CG	1:B:51[A]:ARG:NH1	2.23	0.98
1:B:168[B]:ARG:NH1	1:B:173[B]:GLN:OE1	1.98	0.96
1:A:67:CYS:SG	3:A:874:HOH:O	2.22	0.96
1:B:51[A]:ARG:NH1	1:B:51[A]:ARG:N	2.20	0.88
1:B:80[B]:ARG:CZ	3:B:972:HOH:O	2.27	0.82
1:B:51[B]:ARG:O	1:B:51[B]:ARG:CD	2.29	0.80
1:B:48:ARG:CG	3:B:790:HOH:O	2.29	0.79
1:A:104[B]:MET:CE	1:A:109[B]:LYS:NZ	2.46	0.79
1:B:51[B]:ARG:O	1:B:51[B]:ARG:CG	2.30	0.78
1:A:104[B]:MET:SD	1:A:109[B]:LYS:CD	2.74	0.76
1:B:247[B]:PRO:CG	3:B:860:HOH:O	2.35	0.74
1:B:204[C]:TYR:OH	3:B:974:HOH:O	2.09	0.71
1:A:37:GLN:O	1:A:41[B]:ILE:CD1	2.40	0.70
1:B:39:GLU:OE1	3:B:947:HOH:O	2.10	0.69
1:B:264:VAL:CG1	1:B:268:LYS:NZ	2.56	0.68
1:B:46[A]:THR:CG2	3:B:880:HOH:O	2.42	0.68
1:B:304[B]:LYS:CE	3:B:928:HOH:O	2.42	0.67
1:B:262[A]:ASN:ND2	1:B:263:HIS:CD2	2.62	0.67
1:A:50:ASP:OD1	1:A:54:THR:OG1	2.12	0.66
1:B:147[B]:LYS:NZ	3:B:978:HOH:O	2.33	0.62
1:B:80[B]:ARG:NH2	3:B:972:HOH:O	2.34	0.61
1:A:46[B]:THR:CG2	1:B:206[B]:ASP:OD1	2.50	0.59
1:A:50:ASP:O	1:A:51[A]:ARG:CG	2.52	0.58
1:B:104[A]:MET:CG	1:B:109[A]:LYS:NZ	2.69	0.56
1:A:175[A]:ASP:CB	1:B:51[A]:ARG:NH2	2.73	0.52
1:A:41[B]:ILE:CD1	1:A:62[B]:MET:SD	2.98	0.52
1:B:49[B]:ASP:OD1	1:B:53:GLY:C	2.50	0.49
3:A:749:HOH:O	1:B:200[A]:MET:SD	2.59	0.49
1:B:104[A]:MET:CB	1:B:109[A]:LYS:NZ	2.76	0.48
1:B:165[A]:GLU:OE1	1:B:168[A]:ARG:NH2	2.48	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:282[B]:ARG:NE	1:A:301:ASP:OD2	2.52	0.43
1:B:52:THR:O	1:B:52:THR:CG2	2.66	0.43
1:B:51[B]:ARG:C	1:B:51[B]:ARG:CD	2.89	0.41
1:B:51[A]:ARG:NH1	1:B:51[A]:ARG:CB	2.83	0.41
1:B:104[A]:MET:CE	3:B:965:HOH:O	2.68	0.40
1:B:50:ASP:OD2	1:B:54[A]:THR:OG1	2.40	0.40
1:B:34[B]:TYR:OH	1:B:221:MET:O	2.40	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:172:GLU:OE2	1:B:172:GLU:OE2[2_454]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/315 (101%)	308 (97%)	9 (3%)	1 (0%)	50	17
1	B	320/315 (102%)	311 (97%)	8 (2%)	1 (0%)	50	17
All	All	638/630 (101%)	619 (97%)	17 (3%)	2 (0%)	50	17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	137	TYR
1	B	137	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/276 (103%)	276 (97%)	8 (3%)	56	13
1	B	287/276 (104%)	274 (96%)	13 (4%)	38	5
All	All	571/552 (103%)	550 (96%)	21 (4%)	56	7

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51[A]	ARG
1	A	51[B]	ARG
1	A	52	THR
1	A	200[A]	MET
1	A	200[B]	MET
1	A	214	LEU
1	A	310	PRO
1	A	311	MET
1	B	48	ARG
1	B	51[A]	ARG
1	B	51[B]	ARG
1	B	54[A]	THR
1	B	54[B]	THR
1	B	214	LEU
1	B	247[A]	PRO
1	B	247[B]	PRO
1	B	262[A]	ASN
1	B	262[B]	ASN
1	B	310	PRO
1	B	311[A]	MET
1	B	311[B]	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	401	-	4,4,4	1.29	1 (25%)	6,6,6	0.35	0
2	SO4	B	401	-	4,4,4	1.31	0	6,6,6	1.33	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	401	-	-	0/0/0/0	0/0/0/0
2	SO4	B	401	-	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	SO4	O3-S	2.24	1.54	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	SO4	O4-S-O3	-2.53	98.39	109.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	288/315 (91%)	-0.01	11 (3%) 38 42	7, 10, 26, 120	0
1	B	288/315 (91%)	0.06	14 (4%) 28 32	7, 11, 29, 93	0
All	All	576/630 (91%)	0.03	25 (4%) 34 36	7, 11, 29, 120	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	52	THR	19.2
1	A	51[A]	ARG	11.3
1	B	51[A]	ARG	10.8
1	B	25	VAL	9.5
1	B	52	THR	9.2
1	B	24	GLN	9.0
1	B	54[A]	THR	8.4
1	A	53	GLY	6.8
1	A	54	THR	6.8
1	A	50	ASP	5.6
1	B	50	ASP	4.8
1	B	311[A]	MET	4.8
1	B	309[A]	ILE	4.6
1	A	24	GLN	4.5
1	A	310	PRO	4.4
1	A	49	ASP	4.3
1	A	311	MET	3.5
1	B	53	GLY	3.4
1	B	207[A]	ASN	2.8
1	A	207	ASN	2.5
1	A	262	ASN	2.4
1	B	310	PRO	2.3
1	B	49[A]	ASP	2.2
1	B	262[A]	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	55	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	B	401	5/5	0.12	4.54	27,33,41,47	5
2	SO4	A	401	5/5	0.14	3.23	31,33,41,76	5

6.5 Other polymers ⓘ

There are no such residues in this entry.