



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 11:07 PM GMT

PDB ID : 2IZD
Title : APOSTREPTAVIDIN PH 3.0 I222 COMPLEX
Authors : Katz, B.A.
Deposited on : 1997-08-13
Resolution : 1.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2199 (1.60-1.60)
Ramachandran outliers	78287	2126 (1.60-1.60)
Sidechain outliers	78261	2125 (1.60-1.60)
RSRZ outliers	66119	1872 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	B	123	
1	D	123	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	SO4	B	464	-	X

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 4299 atoms, of which 2158 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called STREPTAVIDIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	123	Total	C	H	N	O	78	14	0
			1955	630	950	169	206			
1	D	121	Total	C	H	N	O	39	13	0
			1907	616	922	167	202			

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Cl	0	0
			1	1		

- Molecule 4 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	I	0	0
			1	1		

- Molecule 5 is AMMONIUM ION (three-letter code: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	H	N	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	65	Total	H	O	0	0
			193	128	65		
6	D	77	Total	H	O	0	0
			231	154	77		

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	94.88Å 105.34Å 47.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.50 – 1.60 33.03 – 1.26	Depositor EDS
% Data completeness (in resolution range)	64.0 (7.50-1.60) 39.0 (33.03-1.26)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 1.26Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.206 , 0.252 0.222 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	9.0	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 107.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 24883 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4299	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, IOD, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	1.42	2/1047 (0.2%)	1.78	30/1434 (2.1%)
1	D	1.40	2/1022 (0.2%)	1.81	28/1400 (2.0%)
All	All	1.41	4/2069 (0.2%)	1.79	58/2834 (2.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	4
1	D	0	3
All	All	0	7

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	52	SER	CA-CB	6.42	1.62	1.52
1	D	14	GLU	CA-CB	5.54	1.66	1.53
1	B	45	SER	CA-CB	5.19	1.60	1.52
1	B	79	TRP	CG-CD2	-5.09	1.34	1.43

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	92	TRP	CD1-NE1-CE2	10.89	118.80	109.00
1	D	21	TRP	CD1-NE1-CE2	10.34	118.30	109.00
1	B	75	TRP	CD1-NE1-CE2	9.95	117.96	109.00
1	D	120	TRP	CD1-NE1-CE2	9.93	117.94	109.00
1	B	92	TRP	CD1-NE1-CE2	9.70	117.73	109.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	21	TRP	CD1-NE1-CE2	9.69	117.72	109.00
1	D	79	TRP	CD1-NE1-CE2	9.64	117.68	109.00
1	D	75	TRP	CD1-NE1-CE2	8.91	117.02	109.00
1	B	120	TRP	CD1-NE1-CE2	8.87	116.98	109.00
1	D	108	TRP	CD1-NE1-CE2	8.76	116.89	109.00
1	D	21	TRP	CG-CD1-NE1	-8.55	101.55	110.10
1	D	92	TRP	CG-CD1-NE1	-8.29	101.81	110.10
1	B	92	TRP	CG-CD1-NE1	-8.23	101.87	110.10
1	D	79	TRP	NE1-CE2-CZ2	8.16	139.38	130.40
1	D	21	TRP	NE1-CE2-CZ2	8.03	139.24	130.40
1	B	75	TRP	CG-CD1-NE1	-8.00	102.10	110.10
1	B	79	TRP	CD1-NE1-CE2	7.98	116.18	109.00
1	D	75	TRP	NE1-CE2-CZ2	7.92	139.11	130.40
1	D	92	TRP	NE1-CE2-CZ2	7.88	139.06	130.40
1	B	21	TRP	NE1-CE2-CZ2	7.74	138.92	130.40
1	B	120	TRP	NE1-CE2-CZ2	7.66	138.82	130.40
1	B	108	TRP	CD1-NE1-CE2	7.39	115.65	109.00
1	B	21	TRP	CG-CD1-NE1	-7.33	102.77	110.10
1	D	92	TRP	NE1-CE2-CD2	-7.32	99.98	107.30
1	B	120	TRP	CG-CD1-NE1	-7.14	102.95	110.10
1	D	108	TRP	CG-CD1-NE1	-6.99	103.11	110.10
1	D	78	ALA	N-CA-C	-6.97	92.18	111.00
1	B	59	ARG	NE-CZ-NH2	-6.92	116.84	120.30
1	D	75	TRP	CG-CD1-NE1	-6.90	103.20	110.10
1	D	79	TRP	CG-CD1-NE1	-6.84	103.26	110.10
1	B	108	TRP	CG-CD1-NE1	-6.75	103.35	110.10
1	B	78	ALA	N-CA-C	-6.54	93.34	111.00
1	D	120	TRP	CG-CD1-NE1	-6.53	103.57	110.10
1	B	79	TRP	CG-CD1-NE1	-6.52	103.58	110.10
1	D	21	TRP	NE1-CE2-CD2	-6.51	100.79	107.30
1	B	21	TRP	NE1-CE2-CD2	-6.50	100.80	107.30
1	D	120	TRP	NE1-CE2-CD2	-6.50	100.80	107.30
1	D	120	TRP	NE1-CE2-CZ2	6.38	137.42	130.40
1	D	84	ARG	NE-CZ-NH2	-6.37	117.12	120.30
1	B	79	TRP	NE1-CE2-CZ2	6.12	137.13	130.40
1	B	61[A]	ASP	CB-CG-OD2	6.08	123.78	118.30
1	B	61[B]	ASP	CB-CG-OD2	6.08	123.78	118.30
1	D	79	TRP	NE1-CE2-CD2	-6.05	101.25	107.30
1	D	108	TRP	NE1-CE2-CZ2	6.02	137.02	130.40
1	B	114	THR	OG1-CB-CG2	-5.95	96.32	110.00
1	D	75	TRP	NE1-CE2-CD2	-5.82	101.48	107.30
1	B	107	GLN	CA-CB-CG	-5.70	100.86	113.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	75	TRP	NE1-CE2-CD2	-5.59	101.71	107.30
1	D	108	TRP	NE1-CE2-CD2	-5.50	101.80	107.30
1	B	133	VAL	CA-CB-CG1	5.42	119.03	110.90
1	B	120	TRP	NE1-CE2-CD2	-5.41	101.89	107.30
1	D	107[A]	GLN	N-CA-C	-5.35	96.56	111.00
1	D	107[B]	GLN	N-CA-C	-5.35	96.56	111.00
1	B	75	TRP	NE1-CE2-CZ2	5.32	136.25	130.40
1	B	92	TRP	NE1-CE2-CZ2	5.31	136.24	130.40
1	B	54	TYR	CB-CG-CD2	-5.19	117.89	121.00
1	B	92	TRP	NE1-CE2-CD2	-5.18	102.12	107.30
1	B	84	ARG	NE-CZ-NH2	-5.12	117.74	120.30

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	103	ARG	Sidechain
1	B	134	LYS	Peptide
1	B	53	ARG	Sidechain
1	B	63[A]	ALA	Peptide
1	D	103	ARG	Sidechain
1	D	53	ARG	Sidechain
1	D	63[A]	ALA	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1005	950	0	24	4
1	D	985	922	0	18	0
2	B	5	0	0	0	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	D	1	0	0	3	0
5	D	1	4	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	65	128	0	13	4
6	D	77	154	0	1	3
All	All	2141	2158	0	37	6

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (37) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:22[A]:TYR:CD2	6:B:1460:HOH:O	1.67	1.27
1:B:22[B]:TYR:CD1	6:B:852:HOH:O	2.00	1.13
1:B:22[B]:TYR:HD1	6:B:852:HOH:O	1.33	1.09
1:B:22[A]:TYR:CZ	6:B:491:HOH:O	2.14	1.01
1:B:22[A]:TYR:OH	6:B:491:HOH:O	1.79	0.99
1:B:22[A]:TYR:CE2	6:B:491:HOH:O	2.23	0.91
1:B:40[A]:THR:HG21	6:B:606:HOH:H2	1.21	0.87
1:B:40[A]:THR:CG2	6:B:606:HOH:H2	1.70	0.86
1:B:22[B]:TYR:CD2	6:B:1460:HOH:O	2.28	0.86
1:D:115:THR:HG21	4:D:463:IOD:I	2.52	0.80
1:D:62[B]:SER:HG	6:D:688:HOH:H2	1.26	0.80
1:B:88:SER:HG	6:B:1419:HOH:H2	1.30	0.79
1:B:22[B]:TYR:CE1	1:B:131:THR:HB	2.23	0.74
1:B:114:THR:H	1:D:95:GLN:HE21	1.47	0.61
1:D:21:TRP:CZ2	1:D:132:LYS:HE3	2.40	0.57
1:B:95:GLN:HE21	1:D:114:THR:H	1.51	0.57
1:B:51:GLU:O	1:B:52:SER:CB	2.53	0.56
1:D:51:GLU:HB2	1:D:84:ARG:NE	2.26	0.51
1:D:51:GLU:O	1:D:52:SER:CB	2.58	0.50
1:B:22[A]:TYR:CE1	1:B:28:THR:HG21	2.47	0.49
1:B:37:GLY:HA2	1:B:60[A]:TYR:CE1	2.49	0.48
1:B:61[A]:ASP:OD2	1:D:85:ASN:OD1	2.25	0.48
1:D:49:ASN:OD1	1:D:84:ARG:HD3	2.13	0.48
1:D:114:THR:HA	4:D:463:IOD:I	2.84	0.47
1:D:51:GLU:O	1:D:52:SER:HB2	2.13	0.47
1:D:13:ALA:O	1:D:15:ALA:N	2.48	0.46
1:D:120:TRP:CZ2	1:D:121:LYS:HD2	2.50	0.46
1:B:22[B]:TYR:CE1	1:B:131:THR:CB	2.97	0.46
1:B:40[A]:THR:HG21	6:B:606:HOH:O	2.00	0.45
1:B:40[A]:THR:CG2	6:B:606:HOH:O	2.50	0.45
1:B:63[B]:ALA:O	1:D:85:ASN:ND2	2.50	0.44
1:B:61[A]:ASP:HD2	1:D:85:ASN:CG	2.14	0.44
1:D:105:ASN:ND2	1:D:129:THR:HG21	2.33	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:115:THR:CG2	4:D:463:IOD:I	3.33	0.43
1:B:135:PRO:C	6:B:652:HOH:H1	2.22	0.42
1:B:39:LEU:HD11	1:B:60[B]:TYR:CD1	2.55	0.41
1:D:30:ILE:O	1:D:41:GLY:HA3	2.22	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
6:B:468:HOH:O	6:B:468:HOH:O[4_555]	0.43	1.77
1:B:22[A]:TYR:CE2	6:D:1465:HOH:O[6_554]	1.58	0.62
1:B:22[B]:TYR:CE1	6:B:852:HOH:O[3_655]	1.66	0.54
6:B:491:HOH:O	6:D:1465:HOH:O[8_555]	1.78	0.42
1:B:22[A]:TYR:CE2	6:B:491:HOH:O[3_655]	2.11	0.09
1:B:22[A]:TYR:CZ	6:D:1465:HOH:O[8_555]	2.13	0.07

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	135/123 (110%)	125 (93%)	6 (4%)	4 (3%)	7	0
1	D	132/123 (107%)	123 (93%)	7 (5%)	2 (2%)	15	2
All	All	267/246 (108%)	248 (93%)	13 (5%)	6 (2%)	11	1

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	52	SER
1	D	52	SER
1	D	14	GLU
1	B	64[A]	PRO
1	B	64[B]	PRO
1	B	36	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	101/90 (112%)	92 (91%)	9 (9%)	14 2
1	D	98/90 (109%)	89 (91%)	9 (9%)	13 2
All	All	199/180 (111%)	181 (91%)	18 (9%)	20 2

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	39	LEU
1	B	44	GLU
1	B	62[A]	SER
1	B	62[B]	SER
1	B	66[A]	THR
1	B	66[B]	THR
1	B	73[A]	LEU
1	B	73[B]	LEU
1	B	133	VAL
1	D	31	VAL
1	D	36	ASP
1	D	52	SER
1	D	53	ARG
1	D	73[A]	LEU
1	D	73[B]	LEU
1	D	107[A]	GLN
1	D	107[B]	GLN
1	D	131	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	82	ASN
1	B	95	GLN
1	D	95	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is modelled with single atom and 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	464	-	4,4,4	1.67	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	B	464	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	122/123 (99%)	0.90	22 (18%) 2 2	8, 25, 46, 66	18 (14%)
1	D	121/123 (98%)	1.10	24 (19%) 2 1	9, 27, 59, 85	25 (20%)
All	All	243/246 (98%)	1.00	46 (18%) 2 2	8, 26, 53, 85	43 (17%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	65[A]	ALA	13.5
1	B	66[A]	THR	12.5
1	D	66[A]	THR	12.4
1	B	49	ASN	11.4
1	B	135	PRO	9.6
1	D	15	ALA	7.9
1	B	47	VAL	7.6
1	B	65[A]	ALA	7.0
1	D	97	VAL	5.9
1	D	133	VAL	5.9
1	B	15	ALA	5.6
1	B	67[A]	ASP	5.3
1	B	50	ALA	4.9
1	B	98	GLY	4.9
1	D	62[A]	SER	4.9
1	D	13	ALA	4.9
1	B	62[A]	SER	4.5
1	D	82	ASN	4.5
1	D	83	TYR	4.2
1	B	22[A]	TYR	4.0
1	D	49	ASN	4.0
1	D	64[A]	PRO	3.8
1	D	69[A]	SER	3.7
1	B	63[A]	ALA	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	64[A]	PRO	3.4
1	D	67[A]	ASP	3.4
1	D	100	ALA	3.4
1	B	133	VAL	3.3
1	D	68[A]	GLY	3.2
1	B	48	GLY	3.2
1	B	13	ALA	3.0
1	B	101	GLU	2.9
1	D	47	VAL	2.9
1	B	97	VAL	2.7
1	D	25	LEU	2.7
1	B	30	ILE	2.7
1	D	30	ILE	2.5
1	D	50	ALA	2.4
1	D	70	GLY	2.4
1	B	100	ALA	2.3
1	B	46	ALA	2.2
1	D	48	GLY	2.1
1	D	51	GLU	2.1
1	D	116	GLU	2.1
1	B	117	ALA	2.1
1	D	102	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	B	464	5/5	0.20	3.58	29,32,33,38	5
5	NH4	D	466	1/1	0.19	1.94	42,42,43,43	0
3	CL	D	467	1/1	0.10	-0.30	55,55,55,55	0
3	CL	B	465	1/1	0.09	-0.42	40,40,40,40	0
4	IOD	D	463	1/1	0.05	-0.96	53,53,53,53	1

6.5 Other polymers ⓘ

There are no such residues in this entry.