



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 10:11 PM GMT

PDB ID : 4IZM  
Title : Crystal structure of GltPh L66C-S300C mutant crosslinked with divalent mercury  
Authors : Reyes, N.; Boudker, O.  
Deposited on : 2013-01-30  
Resolution : 4.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 4.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1029 (5.50-3.50)
Clashscore	79885	1300 (5.50-3.50)
Ramachandran outliers	78287	1222 (5.50-3.50)
Sidechain outliers	78261	1203 (5.50-3.50)
RSRZ outliers	66119	1028 (5.50-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	421	
1	B	421	
1	C	421	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NA	A	502	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8727 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 425aa long hypothetical proton glutamate symport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	0	0	0
			2897	1903	465	511	18			
1	B	404	Total	C	N	O	S	0	0	0
			2897	1903	465	511	18			
1	C	404	Total	C	N	O	S	0	0	0
			2897	1903	465	511	18			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	37	HIS	ASP	ENGINEERED MUTATION	UNP O59010
A	40	HIS	LYS	ENGINEERED MUTATION	UNP O59010
A	66	CYS	LEU	ENGINEERED MUTATION	UNP O59010
A	125	HIS	LYS	ENGINEERED MUTATION	UNP O59010
A	132	HIS	LYS	ENGINEERED MUTATION	UNP O59010
A	223	HIS	LYS	ENGINEERED MUTATION	UNP O59010
A	264	HIS	LYS	ENGINEERED MUTATION	UNP O59010
A	300	CYS	SER	ENGINEERED MUTATION	UNP O59010
A	321	ALA	CYS	ENGINEERED MUTATION	UNP O59010
A	368	HIS	GLU	ENGINEERED MUTATION	UNP O59010
A	418	THR	-	EXPRESSION TAG	UNP O59010
A	419	LEU	-	EXPRESSION TAG	UNP O59010
A	420	VAL	-	EXPRESSION TAG	UNP O59010
A	421	PRO	-	EXPRESSION TAG	UNP O59010
B	37	HIS	ASP	ENGINEERED MUTATION	UNP O59010
B	40	HIS	LYS	ENGINEERED MUTATION	UNP O59010
B	66	CYS	LEU	ENGINEERED MUTATION	UNP O59010
B	125	HIS	LYS	ENGINEERED MUTATION	UNP O59010
B	132	HIS	LYS	ENGINEERED MUTATION	UNP O59010
B	223	HIS	LYS	ENGINEERED MUTATION	UNP O59010
B	264	HIS	LYS	ENGINEERED MUTATION	UNP O59010
B	300	CYS	SER	ENGINEERED MUTATION	UNP O59010
B	321	ALA	CYS	ENGINEERED MUTATION	UNP O59010

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	368	HIS	GLU	ENGINEERED MUTATION	UNP O59010
B	418	THR	-	EXPRESSION TAG	UNP O59010
B	419	LEU	-	EXPRESSION TAG	UNP O59010
B	420	VAL	-	EXPRESSION TAG	UNP O59010
B	421	PRO	-	EXPRESSION TAG	UNP O59010
C	37	HIS	ASP	ENGINEERED MUTATION	UNP O59010
C	40	HIS	LYS	ENGINEERED MUTATION	UNP O59010
C	66	CYS	LEU	ENGINEERED MUTATION	UNP O59010
C	125	HIS	LYS	ENGINEERED MUTATION	UNP O59010
C	132	HIS	LYS	ENGINEERED MUTATION	UNP O59010
C	223	HIS	LYS	ENGINEERED MUTATION	UNP O59010
C	264	HIS	LYS	ENGINEERED MUTATION	UNP O59010
C	300	CYS	SER	ENGINEERED MUTATION	UNP O59010
C	321	ALA	CYS	ENGINEERED MUTATION	UNP O59010
C	368	HIS	GLU	ENGINEERED MUTATION	UNP O59010
C	418	THR	-	EXPRESSION TAG	UNP O59010
C	419	LEU	-	EXPRESSION TAG	UNP O59010
C	420	VAL	-	EXPRESSION TAG	UNP O59010
C	421	PRO	-	EXPRESSION TAG	UNP O59010

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Na 2 2	0	0
2	A	2	Total Na 2 2	0	0
2	C	2	Total Na 2 2	0	0

- Molecule 3 is ASPARTIC ACID (three-letter code: ASP) (formula: C<sub>4</sub>H<sub>7</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	4	1	4		
3	B	1	Total	C	N	O	0	0
			9	4	1	4		
3	C	1	Total	C	N	O	0	0
			9	4	1	4		

- Molecule 4 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Hg	0	0
			1	1		
4	A	1	Total	Hg	0	0
			1	1		
4	C	1	Total	Hg	0	0
			1	1		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 425aa long hypothetical proton glutamate symport protein

Chain A:




- Molecule 1: 425aa long hypothetical proton glutamate symport protein

Chain B:



- Molecule 1: 425aa long hypothetical proton glutamate symport protein

Chain C:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.71Å 119.71Å 333.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 4.50 29.93 – 4.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-4.50) 99.9 (29.93-4.50)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.95 (at 4.42Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.250 , 0.299 0.250 , 0.297	Depositor DCC
$R_{free}$ test set	794 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	150.8	Xtriage
Anisotropy	0.485	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.18 , 112.2	EDS
Estimated twinning fraction	0.098 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 15960 reflections	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	8727	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	195.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2952	0.57	1/4040 (0.0%)
1	B	0.36	0/2952	0.52	1/4040 (0.0%)
1	C	0.37	0/2952	0.54	0/4040
All	All	0.37	0/8856	0.54	2/12120 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	146	LEU	CA-CB-CG	5.86	128.78	115.30
1	A	146	LEU	CA-CB-CG	5.74	128.50	115.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	GLN	Peptide
1	B	114	HIS	Peptide

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	C	202	MET	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2897	0	0	18	0
1	B	2897	0	0	4	0
1	C	2897	0	0	6	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
3	A	9	0	0	1	0
3	B	9	0	0	0	0
3	C	9	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
All	All	8727	0	0	27	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (27) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:355:VAL:CG1	1:A:356:PRO:CD	2.75	0.64
1:B:132:HIS:O	1:B:132:HIS:CG	2.62	0.53
1:A:123:GLN:CB	1:A:124:PRO:CD	2.88	0.52
1:A:296:GLU:CA	1:A:299:TYR:CE1	2.93	0.51
1:A:204:TYR:CG	1:A:204:TYR:O	2.63	0.51
1:A:219:GLU:O	1:A:220:GLN:CB	2.60	0.50
1:A:76:ALA:O	1:A:79:GLY:N	2.46	0.49
1:C:215:TYR:O	1:C:217:MET:N	2.46	0.48
1:C:122:PHE:O	1:C:123:GLN:CB	2.62	0.47
1:A:81:VAL:CG1	1:A:82:GLY:N	2.77	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:191:ALA:O	1:B:195:TYR:CD1	2.68	0.47
1:A:44:LYS:N	1:A:45:PRO:CD	2.79	0.46
1:A:394:ASP:OD1	3:A:503:ASP:N	2.49	0.45
1:A:205:ALA:N	1:A:206:PRO:CD	2.79	0.45
1:C:254:TYR:O	1:C:414:LYS:NZ	2.50	0.45
1:A:267:ASP:N	1:A:267:ASP:OD1	2.50	0.45
1:C:25:ILE:CG2	1:C:25:ILE:O	2.65	0.45
1:A:345:ALA:O	1:A:349:SER:N	2.50	0.44
1:A:321:ALA:C	1:A:323:PHE:N	2.71	0.44
1:A:197:ILE:O	1:A:200:GLY:N	2.51	0.43
1:A:228:LEU:O	1:A:232:THR:OG1	2.37	0.43
1:A:126:GLN:O	1:A:127:ALA:CB	2.67	0.42
1:A:121:GLN:O	1:A:123:GLN:N	2.52	0.42
1:C:76:ALA:O	1:C:79:GLY:N	2.54	0.41
1:A:215:TYR:O	1:A:217:MET:N	2.53	0.41
1:B:110:GLY:N	1:B:227:GLU:OE1	2.54	0.41
1:B:52:ARG:NH1	1:C:138:VAL:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/421 (96%)	325 (81%)	60 (15%)	17 (4%)	4	48
1	B	402/421 (96%)	344 (86%)	48 (12%)	10 (2%)	9	62
1	C	402/421 (96%)	342 (85%)	41 (10%)	19 (5%)	4	45
All	All	1206/1263 (96%)	1011 (84%)	149 (12%)	46 (4%)	5	51

All (46) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	66	CYS
1	A	122	PHE

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	123	GLN
1	A	256	ILE
1	B	152	LEU
1	C	66	CYS
1	C	77	ARG
1	C	123	GLN
1	C	152	LEU
1	C	203	GLN
1	A	152	LEU
1	A	203	GLN
1	A	220	GLN
1	A	334	THR
1	B	66	CYS
1	B	73	ILE
1	A	77	ARG
1	A	222	VAL
1	A	245	LEU
1	B	38	ALA
1	B	123	GLN
1	C	78	LEU
1	C	217	MET
1	C	323	PHE
1	C	334	THR
1	B	222	VAL
1	B	323	PHE
1	B	334	THR
1	C	44	LYS
1	C	45	PRO
1	A	42	TYR
1	C	73	ILE
1	C	216	VAL
1	C	220	GLN
1	C	223	HIS
1	B	151	VAL
1	C	79	GLY
1	C	222	VAL
1	A	127	ALA
1	C	127	ALA
1	A	44	LYS
1	A	151	VAL
1	A	225	VAL
1	C	298	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	73	ILE
1	B	79	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/328 (88%)	234 (82%)	53 (18%)	2	18
1	B	287/328 (88%)	233 (81%)	54 (19%)	2	17
1	C	287/328 (88%)	234 (82%)	53 (18%)	2	18
All	All	861/984 (88%)	701 (81%)	160 (19%)	2	18

All (160) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	TYR
1	A	48	ASP
1	A	59	MET
1	A	78	LEU
1	A	98	THR
1	A	102	ILE
1	A	105	ARG
1	A	106	LEU
1	A	117	VAL
1	A	126	GLN
1	A	130	LEU
1	A	133	ILE
1	A	136	ASP
1	A	137	ILE
1	A	151	VAL
1	A	152	LEU
1	A	155	ILE
1	A	174	GLU
1	A	185	ASP
1	A	198	VAL
1	A	199	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	204	TYR
1	A	212	LEU
1	A	217	MET
1	A	236	TYR
1	A	244	LEU
1	A	245	LEU
1	A	250	LEU
1	A	256	ILE
1	A	267	ASP
1	A	271	THR
1	A	274	VAL
1	A	287	ARG
1	A	290	LYS
1	A	292	MET
1	A	296	GLU
1	A	308	THR
1	A	311	MET
1	A	316	LEU
1	A	334	THR
1	A	343	LEU
1	A	347	LEU
1	A	352	THR
1	A	363	LEU
1	A	365	MET
1	A	370	VAL
1	A	372	LEU
1	A	374	LEU
1	A	385	MET
1	A	387	LEU
1	A	397	ARG
1	A	407	THR
1	A	416	GLU
1	B	13	LEU
1	B	22	LEU
1	B	25	ILE
1	B	29	ILE
1	B	32	HIS
1	B	42	TYR
1	B	43	VAL
1	B	48	ASP
1	B	59	MET
1	B	66	CYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	81	VAL
1	B	98	THR
1	B	106	LEU
1	B	120	GLN
1	B	122	PHE
1	B	133	ILE
1	B	137	ILE
1	B	146	LEU
1	B	151	VAL
1	B	154	THR
1	B	159	ILE
1	B	175	LYS
1	B	182	THR
1	B	184	LEU
1	B	185	ASP
1	B	192	GLU
1	B	198	VAL
1	B	199	ASN
1	B	203	GLN
1	B	217	MET
1	B	245	LEU
1	B	259	ILE
1	B	271	THR
1	B	279	SER
1	B	281	THR
1	B	284	VAL
1	B	290	LYS
1	B	291	GLU
1	B	296	GLU
1	B	299	TYR
1	B	308	THR
1	B	311	MET
1	B	316	LEU
1	B	322	THR
1	B	327	ASN
1	B	334	THR
1	B	340	THR
1	B	342	VAL
1	B	361	ILE
1	B	365	MET
1	B	372	LEU
1	B	389	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	402	VAL
1	B	415	THR
1	C	29	ILE
1	C	32	HIS
1	C	37	HIS
1	C	48	ASP
1	C	59	MET
1	C	65	SER
1	C	81	VAL
1	C	105	ARG
1	C	106	LEU
1	C	120	GLN
1	C	121	GLN
1	C	130	LEU
1	C	136	ASP
1	C	137	ILE
1	C	151	VAL
1	C	152	LEU
1	C	155	ILE
1	C	175	LYS
1	C	179	SER
1	C	184	LEU
1	C	185	ASP
1	C	192	GLU
1	C	194	MET
1	C	198	VAL
1	C	199	ASN
1	C	212	LEU
1	C	213	ILE
1	C	223	HIS
1	C	237	VAL
1	C	245	LEU
1	C	256	ILE
1	C	270	LEU
1	C	271	THR
1	C	279	SER
1	C	284	VAL
1	C	290	LYS
1	C	291	GLU
1	C	308	THR
1	C	311	MET
1	C	316	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	334	THR
1	C	339	LEU
1	C	349	SER
1	C	355	VAL
1	C	365	MET
1	C	368	HIS
1	C	372	LEU
1	C	374	LEU
1	C	375	THR
1	C	385	MET
1	C	402	VAL
1	C	407	THR
1	C	415	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ASP	A	503	-	8,8,8	0.90	0	10,10,10	1.36	1 (10%)
3	ASP	B	503	-	8,8,8	0.91	0	10,10,10	1.37	1 (10%)
3	ASP	C	503	-	8,8,8	0.94	0	10,10,10	1.35	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASP	A	503	-	-	0/8/8/8	0/0/0/0
3	ASP	B	503	-	-	0/8/8/8	0/0/0/0
3	ASP	C	503	-	-	0/8/8/8	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	503	ASP	OXT-C-O	-2.95	117.39	124.07
3	A	503	ASP	OXT-C-O	-2.82	117.70	124.07
3	B	503	ASP	OXT-C-O	-2.74	117.87	124.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	404/421 (95%)	-0.14	1 (0%) 93 89	128, 153, 214, 285	0
1	B	404/421 (95%)	0.17	17 (4%) 35 30	143, 212, 283, 308	0
1	C	404/421 (95%)	0.11	16 (3%) 36 32	142, 206, 293, 328	0
All	All	1212/1263 (95%)	0.05	34 (2%) 50 41	128, 191, 280, 328	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	33	TYR	4.9
1	B	416	GLU	4.7
1	C	292	MET	3.3
1	C	326	ALA	3.2
1	B	332	HIS	3.1
1	C	370	VAL	3.0
1	B	199	ASN	2.9
1	B	125	HIS	2.8
1	B	265	ALA	2.7
1	B	333	LEU	2.7
1	C	223	HIS	2.7
1	C	324	PHE	2.7
1	B	267	ASP	2.7
1	C	293	GLY	2.6
1	B	266	LYS	2.6
1	B	76	ALA	2.5
1	C	227	GLU	2.5
1	B	290	LYS	2.4
1	C	343	LEU	2.3
1	A	254	TYR	2.3
1	B	230	LYS	2.3
1	B	79	GLY	2.3
1	C	322	THR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	291	GLU	2.2
1	C	222	VAL	2.2
1	B	355	VAL	2.2
1	B	264	HIS	2.2
1	C	331	SER	2.2
1	C	203	GLN	2.1
1	C	332	HIS	2.1
1	C	118	GLY	2.1
1	B	331	SER	2.1
1	C	325	ILE	2.1
1	B	203	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
2	NA	A	502	1/1	0.56	2.02	124,124,124,124	0
2	NA	C	502	1/1	0.41	1.11	184,184,184,184	0
2	NA	B	502	1/1	0.35	0.76	187,187,187,187	0
3	ASP	C	503	9/9	0.40	0.44	177,182,185,188	0
3	ASP	A	503	9/9	0.29	0.08	130,131,131,131	0
2	NA	A	501	1/1	0.13	-0.50	127,127,127,127	0
3	ASP	B	503	9/9	0.30	-0.68	187,189,190,192	0
2	NA	B	501	1/1	0.12	-0.96	178,178,178,178	0
4	HG	B	504	1/1	0.04	-1.37	208,208,208,208	0
2	NA	C	501	1/1	0.13	-1.43	182,182,182,182	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	HG	C	504	1/1	0.03	-3.47	215,215,215,215	0
4	HG	A	504	1/1	0.06	-6.14	172,172,172,172	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.