



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 01:01 PM GMT

PDB ID : 2J04
Title : THE TAU60-TAU91 SUBCOMPLEX OF YEAST TRANSCRIPTION FACTOR IIIC
Authors : Mylona, A.; Fernandez-Tornero, C.; Legrand, P.; Muller, C.W.
Deposited on : 2006-07-31
Resolution : 3.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1824 (3.30-3.10)
Clashscore	79885	1078 (3.26-3.14)
Ramachandran outliers	78287	1059 (3.26-3.14)
Sidechain outliers	78261	1058 (3.26-3.14)
RSRZ outliers	66119	1825 (3.30-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	588	
1	C	588	
2	B	524	
2	D	524	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16935 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HYPOTHETICAL PROTEIN YPL007C.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	587	Total	C	N	O	S	0	0	0
			4760	3058	777	902	23			
1	C	586	Total	C	N	O	S	0	0	0
			4751	3052	775	901	23			


- Molecule 2 is a protein called YDR362CP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	472	Total	C	N	O	S	0	0	1
			3715	2384	613	703	15			
2	D	469	Total	C	N	O	S	0	0	1
			3704	2379	609	701	15			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	O	0	0
			2	2		
3	B	1	Total	O	0	0
			1	1		
3	C	1	Total	O	0	0
			1	1		
3	D	1	Total	O	0	0
			1	1		





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.42Å 125.80Å 210.47Å 90.00° 94.49° 90.00°	Depositor
Resolution (Å)	208.51 – 3.20 34.91 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.7 (208.51-3.20) 98.8 (34.91-3.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 3.18Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.212 , 0.257 0.208 , 0.254	Depositor DCC
R_{free} test set	2619 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	77.5	Xtriage
Anisotropy	0.356	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 52096 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	16935	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	5/4872 (0.1%)	0.94	15/6610 (0.2%)
1	C	0.80	5/4863 (0.1%)	0.95	16/6599 (0.2%)
2	B	0.76	4/3798 (0.1%)	0.98	21/5153 (0.4%)
2	D	0.77	4/3788 (0.1%)	0.95	15/5140 (0.3%)
All	All	0.79	18/17321 (0.1%)	0.95	67/23502 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	3
2	B	0	8
2	D	0	7
All	All	0	20

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	413	LYS	CD-CE	18.55	1.97	1.51
2	B	413	LYS	CD-CE	14.62	1.87	1.51
1	A	249	LYS	CD-CE	13.65	1.85	1.51
1	A	297	GLU	CD-OE1	11.57	1.38	1.25
1	C	249	LYS	CD-CE	11.35	1.79	1.51

The worst 5 of 67 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	413	LYS	CD-CE-NZ	-8.44	92.29	111.70
2	B	413	LYS	CD-CE-NZ	-8.38	92.42	111.70
2	D	413	LYS	CG-CD-CE	-7.55	89.24	111.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	6	ASP	CB-CG-OD2	7.20	124.78	118.30
1	A	231	ASP	CB-CG-OD2	7.00	124.60	118.30

There are no chirality outliers.

5 of 20 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	243	CYS	Peptide
1	A	98	ASP	Peptide
2	B	277	ALA	Peptide
2	B	293	THR	Peptide
2	B	379	HIS	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4760	0	4722	220	0
1	C	4751	0	4709	244	0
2	B	3715	0	3674	258	0
2	D	3704	0	3660	238	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
All	All	16935	0	16765	942	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 28.

The worst 5 of 942 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:249:LYS:CD	1:C:249:LYS:CE	1.79	1.60
2:B:413:LYS:CD	2:B:413:LYS:CE	1.87	1.52
1:A:249:LYS:CE	1:A:249:LYS:CD	1.85	1.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:413:LYS:NZ	2:B:413:LYS:CE	1.74	1.47
2:D:413:LYS:CE	2:D:413:LYS:NZ	1.76	1.45

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	585/588 (100%)	475 (81%)	74 (13%)	36 (6%)	2	19
1	C	584/588 (99%)	475 (81%)	69 (12%)	40 (7%)	2	15
2	B	462/524 (88%)	362 (78%)	68 (15%)	32 (7%)	2	14
2	D	459/524 (88%)	356 (78%)	76 (17%)	27 (6%)	2	20
All	All	2090/2224 (94%)	1668 (80%)	287 (14%)	135 (6%)	2	17

5 of 135 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	LEU
1	A	124	LYS
1	A	159	ARG
1	A	164	ASN
1	A	166	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	543/544 (100%)	434 (80%)	109 (20%)	2	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	542/544 (100%)	434 (80%)	108 (20%)	2	9
2	B	415/465 (89%)	340 (82%)	75 (18%)	2	12
2	D	414/465 (89%)	336 (81%)	78 (19%)	2	11
All	All	1914/2018 (95%)	1544 (81%)	370 (19%)	2	10

5 of 370 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	549	LEU
1	C	159	ARG
2	D	455	VAL
2	B	586	LEU
1	C	22	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	641	ASN
1	C	106	ASN
2	D	565	HIS
1	C	41	GLN
1	C	138	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	587/588 (99%)	0.11	1 (0%) 93 66	61, 71, 71, 81	0
1	C	586/588 (99%)	0.12	3 (0%) 88 46	60, 71, 71, 87	0
2	B	472/524 (90%)	0.12	0 100 100	70, 71, 71, 72	0
2	D	469/524 (89%)	0.12	2 (0%) 90 51	71, 71, 71, 75	0
All	All	2114/2224 (95%)	0.11	6 (0%) 91 58	60, 71, 71, 87	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	637	ALA	2.6
1	C	163	GLU	2.5
1	A	253	LYS	2.5
2	D	305	ASP	2.4
1	C	160	LYS	2.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.