



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 10:09 PM GMT

PDB ID : 2J0Y
Title : L-FICOLIN COMPLEXED TO B-1,3-D-GLUCAN
Authors : Garlatti, V.; Gaboriaud, C.
Deposited on : 2006-08-08
Resolution : 2.35 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3327 (2.40-2.32)
Clashscore	79885	1064 (2.38-2.34)
Ramachandran outliers	78287	1048 (2.38-2.34)
Sidechain outliers	78261	1049 (2.38-2.34)
RSRZ outliers	66119	3330 (2.40-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	218	
1	B	218	
1	C	218	
1	D	218	
1	E	218	
1	F	218	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
7	BGC	F	1289	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 10651 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FICOLIN-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	193	Total	C	N	O	S	0	0	0
			1566	988	273	297	8			
1	B	218	Total	C	N	O	S	0	0	0
			1744	1096	307	332	9			
1	C	215	Total	C	N	O	S	0	0	0
			1723	1084	303	328	8			
1	D	202	Total	C	N	O	S	0	1	0
			1631	1024	288	310	9			
1	E	218	Total	C	N	O	S	0	0	0
			1744	1096	307	332	9			
1	F	215	Total	C	N	O	S	0	1	0
			1734	1093	304	329	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	168	THR	VAL	CONFLICT	UNP Q15485
A	247	THR	VAL	CONFLICT	UNP Q15485
B	168	THR	VAL	CONFLICT	UNP Q15485
B	247	THR	VAL	CONFLICT	UNP Q15485
C	168	THR	VAL	CONFLICT	UNP Q15485
C	247	THR	VAL	CONFLICT	UNP Q15485
D	168	THR	VAL	CONFLICT	UNP Q15485
D	247	THR	VAL	CONFLICT	UNP Q15485
E	168	THR	VAL	CONFLICT	UNP Q15485
E	247	THR	VAL	CONFLICT	UNP Q15485
F	168	THR	VAL	CONFLICT	UNP Q15485
F	247	THR	VAL	CONFLICT	UNP Q15485

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		
3	C	2	Total	Ca	0	0
			2	2		
3	F	2	Total	Ca	0	0
			2	2		
3	E	1	Total	Ca	0	0
			1	1		

- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	3	Total	C	N	O	0	0
			39	22	2	15		
4	E	3	Total	C	N	O	0	0
			39	22	2	15		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	168	THR	VAL	CONFLICT	UNP Q15485

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	247	THR	VAL	CONFLICT	UNP Q15485
E	168	THR	VAL	CONFLICT	UNP Q15485
E	247	THR	VAL	CONFLICT	UNP Q15485


- Molecule 5 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	4	Total	C	O	0	0
			44	24	20		
5	F	4	Total	C	O	0	0
			44	24	20		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	168	THR	VAL	CONFLICT	UNP Q15485
C	247	THR	VAL	CONFLICT	UNP Q15485
F	168	THR	VAL	CONFLICT	UNP Q15485
F	247	THR	VAL	CONFLICT	UNP Q15485

- Molecule 6 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	C	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is SUGAR (BETA-D-GLUCOSE) (three-letter code: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			11	6	5		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	32	Total	O	0	0
			32	32		
8	B	61	Total	O	0	0
			61	61		
8	C	57	Total	O	0	0
			57	57		
8	D	24	Total	O	0	0
			24	24		
8	E	66	Total	O	0	0
			66	66		
8	F	54	Total	O	0	0
			54	54		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: FICOLIN-2

Chain A:




• Molecule 1: FICOLIN-2

Chain B:




• Molecule 1: FICOLIN-2


Chain C:



• Molecule 1: FICOLIN-2


Chain D:






● Molecule 1: FICOLIN-2

Chain E:



● Molecule 1: FICOLIN-2

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	96.08Å 96.08Å 140.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.95 – 2.35 19.95 – 2.35	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.95-2.35) 98.8 (19.95-2.35)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 2.35Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.242 0.210 , 0.231	Depositor DCC
R_{free} test set	2990 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 31.5	EDS
Estimated twinning fraction	0.028 for -h,-k,l 0.470 for h,-h-k,-l 0.029 for -k,-h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 59794 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	10651	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, CA, BMA, NAG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	1/1605 (0.1%)	0.60	1/2164 (0.0%)
1	B	0.54	0/1792	0.66	0/2425
1	C	0.53	0/1770	0.65	0/2394
1	D	0.70	1/1671 (0.1%)	0.56	0/2253
1	E	0.52	0/1792	0.65	0/2425
1	F	0.54	0/1782	0.66	0/2410
All	All	0.57	2/10412 (0.0%)	0.63	1/14071 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	273	GLY	C-O	21.08	1.57	1.23
1	A	208	ASP	CG-OD1	8.77	1.45	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	208	ASP	CB-CG-OD2	-5.43	113.42	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1566	0	1441	55	0
1	B	1744	0	1611	31	1
1	C	1723	0	1595	37	0
1	D	1631	0	1510	64	1
1	E	1744	0	1611	33	0
1	F	1734	0	1603	32	0
2	B	4	0	3	0	0
3	B	1	0	0	0	0
3	C	2	0	0	0	0
3	E	1	0	0	0	0
3	F	2	0	0	0	0
4	B	39	0	34	0	0
4	E	39	0	34	0	0
5	C	44	0	37	5	0
5	F	44	0	37	6	0
6	C	14	0	13	0	0
6	F	14	0	13	0	0
7	F	11	0	10	8	0
8	A	32	0	0	12	0
8	B	61	0	0	7	0
8	C	57	0	0	6	0
8	D	24	0	0	24	0
8	E	66	0	0	9	0
8	F	54	0	0	3	0
All	All	10651	0	9552	258	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 258 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:183:ARG:HG3	1:D:183:ARG:HH11	1.06	1.10
1:D:206:ALA:HB1	8:D:2022:HOH:O	1.57	1.05
7:F:1289:BGC:H6C1	7:F:1289:BGC:H2	1.45	0.98
1:D:220:THR:HG22	8:D:2018:HOH:O	1.63	0.98
1:B:71:ASN:N	1:B:72:PRO:HD3	1.83	0.94

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:183:ARG:NH1	1:D:288:ALA:O[3_455]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/218 (83%)	162 (90%)	17 (9%)	2 (1%)	21	21
1	B	216/218 (99%)	205 (95%)	10 (5%)	1 (0%)	38	45
1	C	213/218 (98%)	199 (93%)	13 (6%)	1 (0%)	38	45
1	D	193/218 (88%)	171 (89%)	19 (10%)	3 (2%)	14	13
1	E	216/218 (99%)	203 (94%)	12 (6%)	1 (0%)	38	45
1	F	214/218 (98%)	201 (94%)	12 (6%)	1 (0%)	38	45
All	All	1233/1308 (94%)	1141 (92%)	83 (7%)	9 (1%)	30	34

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	ASN
1	D	206	ALA
1	D	195	ASN
1	A	124	VAL
1	B	124	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/183 (90%)	153 (93%)	12 (7%)	20	22
1	B	183/183 (100%)	172 (94%)	11 (6%)	27	32
1	C	180/183 (98%)	168 (93%)	12 (7%)	23	26
1	D	172/183 (94%)	160 (93%)	12 (7%)	21	23
1	E	183/183 (100%)	170 (93%)	13 (7%)	21	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	181/183 (99%)	167 (92%)	14 (8%)	18	20
All	All	1064/1098 (97%)	990 (93%)	74 (7%)	22	23

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	231	ASN
1	D	150	LEU
1	F	193	LYS
1	C	250	LEU
1	D	101[A]	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	216	GLN
1	D	195	ASN
1	F	195	ASN
1	C	231	ASN
1	D	139	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	B	1291	1,4	12,14,15	0.89	1 (8%)	15,19,21	1.67	3 (20%)
4	NAG	B	1292	4	12,14,15	0.74	1 (8%)	15,19,21	1.18	2 (13%)
4	BMA	B	1293	4	10,11,12	0.85	1 (10%)	11,15,17	1.03	0
5	BGC	C	1289	5	10,11,12	0.70	0	11,15,17	1.70	4 (36%)
5	BGC	C	1290	5	10,11,12	0.70	0	11,15,17	2.78	4 (36%)
5	BGC	C	1291	5	10,11,12	0.84	0	11,15,17	1.88	4 (36%)
5	BGC	C	1292	5	10,11,12	0.67	0	11,15,17	2.42	5 (45%)
4	NAG	E	1290	1,4	12,14,15	0.86	1 (8%)	15,19,21	1.20	1 (6%)
4	NAG	E	1291	4	12,14,15	0.66	0	15,19,21	1.12	1 (6%)
4	BMA	E	1292	4	10,11,12	0.53	0	11,15,17	2.13	3 (27%)
5	BGC	F	1290	5	10,11,12	0.75	0	11,15,17	1.97	5 (45%)
5	BGC	F	1291	5	10,11,12	0.70	0	11,15,17	2.65	4 (36%)
5	BGC	F	1292	5	10,11,12	0.82	0	11,15,17	1.77	3 (27%)
5	BGC	F	1293	5	10,11,12	0.79	1 (10%)	11,15,17	2.03	4 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1291	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	1292	4	-	0/6/23/26	0/1/1/1
4	BMA	B	1293	4	-	0/2/19/22	0/1/1/1
5	BGC	C	1289	5	-	0/2/19/22	0/1/1/1
5	BGC	C	1290	5	-	0/2/19/22	0/1/1/1
5	BGC	C	1291	5	-	0/2/19/22	0/1/1/1
5	BGC	C	1292	5	-	0/2/19/22	0/1/1/1
4	NAG	E	1290	1,4	-	0/6/23/26	0/1/1/1
4	NAG	E	1291	4	-	0/6/23/26	0/1/1/1
4	BMA	E	1292	4	-	0/2/19/22	0/1/1/1
5	BGC	F	1290	5	-	0/2/19/22	0/1/1/1
5	BGC	F	1291	5	-	0/2/19/22	0/1/1/1
5	BGC	F	1292	5	-	0/2/19/22	0/1/1/1
5	BGC	F	1293	5	-	0/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1291	NAG	O5-C5	-2.68	1.40	1.45
4	E	1290	NAG	O5-C5	-2.61	1.40	1.45
4	B	1292	NAG	O5-C5	-2.26	1.41	1.45
4	B	1293	BMA	O5-C5	-2.10	1.41	1.45
5	F	1293	BGC	O5-C5	-2.01	1.41	1.45

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1290	BGC	C4-C3-C2	-6.47	101.82	110.50
5	F	1291	BGC	O5-C5-C6	6.14	113.42	106.98
5	F	1291	BGC	C4-C3-C2	-5.25	103.46	110.50
5	C	1290	BGC	O5-C5-C6	5.19	112.42	106.98
4	E	1292	BMA	O5-C5-C4	4.79	116.74	110.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	B	1289	-	1,3,3	1.17	0	0,3,3	0.00	-
6	NAG	C	1295	1	12,14,15	0.74	1 (8%)	15,19,21	0.89	0
7	BGC	F	1289	-	10,11,12	0.62	0	11,15,17	1.42	2 (18%)
6	NAG	F	1296	1	12,14,15	0.81	1 (8%)	15,19,21	1.02	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	B	1289	-	-	0/0/0/0	0/0/0/0
6	NAG	C	1295	1	-	0/6/23/26	0/1/1/1
7	BGC	F	1289	-	-	0/2/19/22	1/1/1/1
6	NAG	F	1296	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	1295	NAG	O5-C5	-2.29	1.41	1.45
6	F	1296	NAG	O5-C5	-2.27	1.41	1.45

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	F	1289	BGC	O5-C5-C6	3.01	110.14	106.98
6	F	1296	NAG	C3-C2-N2	-2.19	108.43	111.76
7	F	1289	BGC	O4-C4-C5	2.07	114.73	109.28

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	F	1289	BGC	C1-C2-C3-C4-C5-O5

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	193/218 (88%)	1.56	62 (32%) 1 1	42, 46, 56, 59	0
1	B	218/218 (100%)	0.42	2 (0%) 81 84	38, 45, 54, 62	0
1	C	215/218 (98%)	0.33	1 (0%) 88 91	39, 45, 56, 67	0
1	D	202/218 (92%)	1.67	76 (37%) 1 0	42, 45, 50, 53	0
1	E	218/218 (100%)	0.40	0 100 100	38, 45, 54, 61	0
1	F	215/218 (98%)	0.44	5 (2%) 57 61	39, 45, 55, 68	0
All	All	1261/1308 (96%)	0.78	146 (11%) 6 6	38, 45, 54, 68	0

The worst 5 of 146 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	268	TRP	7.3
1	D	233	ALA	6.9
1	D	255	LEU	6.6
1	D	191	ALA	6.4
1	D	271	GLY	5.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	BMA	B	1293	11/12	0.47	20.19	63,65,66,66	0
4	NAG	B	1291	14/15	0.31	12.83	45,47,50,50	0
4	NAG	E	1290	14/15	0.20	5.30	36,38,40,41	0
5	BGC	F	1291	11/12	0.22	4.01	50,52,54,55	0
4	NAG	B	1292	14/15	0.29	3.95	52,53,55,59	0
5	BGC	C	1291	11/12	0.19	3.18	52,53,55,56	0
5	BGC	C	1289	11/12	0.17	2.69	47,51,52,55	0
5	BGC	F	1292	11/12	0.18	2.23	50,52,54,55	0
5	BGC	F	1290	11/12	0.21	2.17	47,51,52,53	0
5	BGC	C	1292	11/12	0.14	0.29	54,57,59,61	0
5	BGC	C	1290	11/12	0.14	0.13	47,50,53,55	0
5	BGC	F	1293	11/12	0.12	-0.12	55,56,57,59	0
4	NAG	E	1291	14/15	0.16	-0.33	40,41,44,49	0
4	BMA	E	1292	11/12	0.17	-	54,58,60,61	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	BGC	F	1289	11/12	0.32	9.09	70,76,77,78	0
2	ACT	B	1289	4/4	0.17	1.47	56,57,58,58	0
3	CA	F	1295	1/1	0.13	-0.38	48,48,48,48	0
3	CA	C	1293	1/1	0.10	-1.24	42,42,42,42	0
3	CA	F	1294	1/1	0.08	-1.84	38,38,38,38	0
3	CA	E	1289	1/1	0.09	-3.28	28,28,28,28	0
3	CA	B	1290	1/1	0.08	-5.08	27,27,27,27	0
3	CA	C	1294	1/1	0.09	-6.16	46,46,46,46	0
6	NAG	F	1296	14/15	0.18	-	62,65,68,68	0
6	NAG	C	1295	14/15	0.14	-	61,65,67,67	0

6.5 Other polymers

There are no such residues in this entry.