



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2014 – 03:48 PM GMT

PDB ID : 4J06
Title : Crystal structure of hcv ns5b polymerase in complex with 2-{[(5-BROMOTH
IOPHEN-2-YL)SULFONYL]AMINO}-4-CHLOROBENZOICACID
Authors : Coulombe,R.
Deposited on : 2013-01-30
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	576	
1	B	576	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9285 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	559	Total	C	N	O	S	0	0	0
			4358	2745	770	811	32			
1	B	558	Total	C	N	O	S	0	0	0
			4346	2737	768	809	32			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	571	HIS	-	EXPRESSION TAG	UNP O92972
A	572	HIS	-	EXPRESSION TAG	UNP O92972
A	573	HIS	-	EXPRESSION TAG	UNP O92972
A	574	HIS	-	EXPRESSION TAG	UNP O92972
A	575	HIS	-	EXPRESSION TAG	UNP O92972
A	576	HIS	-	EXPRESSION TAG	UNP O92972
B	571	HIS	-	EXPRESSION TAG	UNP O92972
B	572	HIS	-	EXPRESSION TAG	UNP O92972
B	573	HIS	-	EXPRESSION TAG	UNP O92972
B	574	HIS	-	EXPRESSION TAG	UNP O92972
B	575	HIS	-	EXPRESSION TAG	UNP O92972
B	576	HIS	-	EXPRESSION TAG	UNP O92972

- Molecule 2 is 2-{[(5-BROMOTHIOPHEN-2-YL)SULFONYL]AMINO}-4-CHLOROBENZOIC ACID (three-letter code: 1JG) (formula: C₁₁H₇BrClNO₄S₂).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	Br	C	Cl	N	O	S	0	0
			20	1	11	1	1	4	2		
2	B	1	Total	Br	C	Cl	N	O	S	0	0
			20	1	11	1	1	4	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	315	Total	O	0	0
			315	315		
3	B	226	Total	O	0	0
			226	226		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Genome polyprotein

Chain A: 



- Molecule 1: Genome polyprotein

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.50Å 108.01Å 133.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.00 39.06 – 2.01	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.00-2.00) 88.7 (39.06-2.01)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 2.01Å)	Xtriage
Refinement program	cnx 2002	Depositor
R, R_{free}	0.208 , 0.235 0.207 , 0.233	Depositor DCC
R_{free} test set	9163 reflections (11.11%)	DCC
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.347	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.2	EDS
Estimated twinning fraction	0.014 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 91646 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9285	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1JG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/4453	0.58	0/6044
1	B	0.27	0/4440	0.56	0/6025
All	All	0.28	0/8893	0.57	0/12069

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4358	0	4371	40	0
1	B	4346	0	4359	63	0
2	A	20	0	6	0	0
2	B	20	0	6	0	0
3	A	315	0	0	3	0
3	B	226	0	0	4	0
All	All	9285	0	8742	102	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (102) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:313:MET:HG2	1:B:322:VAL:HG22	1.55	0.87
1:A:313:MET:HG2	1:A:322:VAL:HG22	1.67	0.76
1:B:337:ARG:O	1:B:341:GLU:HG3	1.86	0.75
1:B:106:LYS:NZ	1:B:110:ASN:HD21	1.93	0.67
1:B:515:GLY:HA2	1:B:519:ALA:HB2	1.79	0.65
1:A:394:ARG:O	1:A:398:GLU:HG3	1.97	0.64
1:B:85:ILE:HD13	1:B:116:VAL:HG13	1.79	0.64
1:B:531:ARG:H	1:B:531:ARG:HH11	1.48	0.61
1:A:381:VAL:HG11	1:A:474:LEU:CD2	2.32	0.60
1:B:144:VAL:HB	1:B:394:ARG:HG2	1.85	0.59
1:B:531:ARG:HD2	1:B:532:THR:H	1.68	0.58
1:B:336:LEU:HD23	1:B:356:PRO:HD3	1.86	0.58
1:B:372:VAL:HG21	1:B:380:ARG:NH1	2.19	0.58
1:B:390:THR:HB	1:B:391:PRO:HD3	1.87	0.56
1:B:32:ARG:HG2	1:B:493:GLY:O	2.05	0.56
1:B:314:LEU:HB3	1:B:321:VAL:CG1	2.36	0.56
1:A:381:VAL:HG11	1:A:474:LEU:HD22	1.88	0.55
1:B:531:ARG:HD2	1:B:532:THR:N	2.22	0.55
1:B:346:TYR:O	1:B:347:SER:HB3	2.07	0.55
1:A:515:GLY:HA2	1:A:519:ALA:HB2	1.89	0.54
1:B:106:LYS:HZ3	1:B:110:ASN:HD21	1.56	0.53
1:A:314:LEU:HB3	1:A:321:VAL:CG1	2.40	0.52
1:B:27:SER:HB2	1:B:400:ALA:HB2	1.92	0.52
1:A:212:LYS:HE2	3:A:976:HOH:O	2.10	0.52
1:B:48:ARG:HG2	1:B:159:LEU:HG	1.91	0.51
1:A:107:ASP:HB3	1:A:112:SER:OG	2.11	0.51
1:A:212:LYS:HD2	1:A:325:GLU:OE1	2.10	0.50
1:A:309:GLN:O	1:A:324:CYS:HB2	2.12	0.50
1:B:498:ARG:O	1:B:501:ARG:HG2	2.13	0.49
1:B:501:ARG:O	1:B:505:ARG:HG3	2.11	0.49
1:B:465:ARG:HH11	1:B:545:LEU:HB3	1.78	0.49
1:A:306:ALA:O	1:A:307:LYS:HB2	2.11	0.49
1:B:488:CYS:HB2	3:B:917:HOH:O	2.12	0.49
1:A:24:ASN:HB3	1:A:27:SER:OG	2.13	0.49
1:A:247:PRO:HG3	1:B:234:ARG:HD3	1.95	0.48
1:B:84:SER:OG	1:B:87:GLU:HG3	2.13	0.48
1:B:508:ARG:HD2	1:B:526:PHE:O	2.14	0.48
1:B:470:SER:O	1:B:474:LEU:HG	2.13	0.48
1:B:531:ARG:HD3	1:B:532:THR:HG23	1.94	0.48
1:A:176:TYR:OH	1:A:562:HIS:HE1	1.95	0.48
1:B:191:TYR:O	1:B:194:GLN:HG2	2.13	0.47
1:B:233:ILE:HD13	1:B:261:TYR:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:273:ASN:ND2	1:B:275:GLY:H	2.13	0.47
1:B:106:LYS:HZ2	1:B:110:ASN:HD21	1.63	0.47
1:A:505:ARG:HH21	1:A:531:ARG:NH1	2.13	0.46
1:B:309:GLN:O	1:B:324:CYS:HB2	2.15	0.46
1:A:485:VAL:O	1:A:489:LEU:HG	2.15	0.46
1:A:119:ILE:HD13	1:A:169:VAL:HG11	1.96	0.46
1:A:57:LEU:HD23	1:A:57:LEU:C	2.35	0.46
1:A:106:LYS:HB2	1:A:106:LYS:HZ2	1.79	0.46
1:A:440:GLU:HG2	1:A:457:LEU:CD1	2.46	0.46
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.98	0.45
1:B:183:PRO:HG3	1:B:289:CYS:SG	2.56	0.45
1:B:93:PRO:HG3	1:B:561:TYR:HB2	1.99	0.45
1:A:390:THR:HB	1:A:391:PRO:HD3	1.99	0.45
1:B:545:LEU:HG	1:B:547:LEU:HG	1.99	0.45
1:B:119:ILE:HD13	1:B:169:VAL:HG11	1.98	0.45
1:B:182:LEU:HD23	1:B:182:LEU:C	2.37	0.45
1:A:300:THR:HA	1:A:313:MET:HE1	1.98	0.44
1:B:523:ARG:NH2	1:B:536:LEU:HA	2.32	0.44
1:A:184:GLN:HB2	3:A:768:HOH:O	2.17	0.44
1:B:359:ASP:HB3	1:B:362:LEU:HD12	1.98	0.44
1:B:83:LEU:HB2	1:B:173:MET:HA	2.00	0.44
1:A:346:TYR:O	1:A:347:SER:HB3	2.17	0.44
1:A:466:LEU:HD22	1:A:551:PHE:HE2	1.83	0.44
1:A:82:LEU:HD13	1:A:249:ALA:HB2	2.00	0.44
1:B:268:ASN:HB3	1:B:274:CYS:SG	2.59	0.43
1:B:466:LEU:HD22	1:B:551:PHE:HE2	1.83	0.43
1:A:233:ILE:HD13	1:A:261:TYR:O	2.19	0.43
1:B:398:GLU:HG2	1:B:403:THR:OG1	2.19	0.43
1:A:183:PRO:HG3	1:A:289:CYS:SG	2.59	0.43
1:B:465:ARG:NH1	1:B:545:LEU:HB3	2.34	0.43
1:A:144:VAL:HB	1:A:394:ARG:HG2	2.00	0.43
1:A:115:ALA:O	1:A:119:ILE:HG13	2.18	0.43
1:A:184:GLN:HG3	3:A:852:HOH:O	2.18	0.42
1:B:465:ARG:HG3	1:B:545:LEU:HD23	2.00	0.42
1:B:178:VAL:HG23	3:B:876:HOH:O	2.18	0.42
1:B:85:ILE:CD1	1:B:116:VAL:HG13	2.47	0.42
1:B:455:GLU:HB3	3:B:916:HOH:O	2.19	0.42
1:A:300:THR:HA	1:A:313:MET:CE	2.49	0.42
1:A:93:PRO:HG3	1:A:561:TYR:HB2	2.01	0.42
1:B:421:ALA:O	1:B:426:MET:HG3	2.20	0.42
1:B:388:PRO:HG2	1:B:488:CYS:SG	2.60	0.42
1:B:40:THR:HB	1:B:157:ALA:HB2	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:398:GLU:OE1	1:B:408:TRP:HD1	2.03	0.41
1:A:461:GLN:HB2	1:A:545:LEU:HD11	2.02	0.41
1:B:160:ILE:HA	1:B:282:SER:OG	2.19	0.41
1:A:93:PRO:HA	1:A:94:PRO:HD3	1.94	0.41
1:B:27:SER:C	1:B:29:SER:H	2.23	0.41
1:B:217:PHE:CE1	1:B:322:VAL:HB	2.55	0.41
1:B:21:LEU:HD12	1:B:34:HIS:HB2	2.03	0.41
1:B:505:ARG:NH1	1:B:531:ARG:NH1	2.68	0.41
1:B:22:PRO:HG2	1:B:400:ALA:HB1	2.02	0.41
1:A:79:LYS:HG3	1:A:244:ASP:HB3	2.01	0.41
1:A:92:THR:HA	1:A:93:PRO:HD3	1.92	0.41
1:B:132:THR:HA	1:B:133:PRO:HD3	1.98	0.41
1:A:5:THR:O	1:A:275:GLY:HA3	2.21	0.41
1:A:336:LEU:HD12	1:A:356:PRO:HD3	2.03	0.41
1:B:539:ILE:HA	1:B:540:PRO:HD3	1.91	0.41
1:B:47:LEU:HG	3:B:880:HOH:O	2.21	0.40
1:B:372:VAL:HG22	1:B:373:ALA:N	2.37	0.40
1:A:257:THR:O	1:A:262:ILE:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	555/576 (96%)	546 (98%)	8 (1%)	1 (0%)	56	51
1	B	554/576 (96%)	538 (97%)	15 (3%)	1 (0%)	56	51
All	All	1109/1152 (96%)	1084 (98%)	23 (2%)	2 (0%)	56	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	154	ARG
1	A	544	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	477/491 (97%)	474 (99%)	3 (1%)	92	94
1	B	475/491 (97%)	473 (100%)	2 (0%)	95	97
All	All	952/982 (97%)	947 (100%)	5 (0%)	94	96

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	LYS
1	A	184	GLN
1	A	336	LEU
1	B	273	ASN
1	B	531	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	251	GLN
1	A	273	ASN
1	A	374	HIS
1	A	438	GLN
1	A	544	GLN
1	A	562	HIS
1	B	33	HIS
1	B	110	ASN
1	B	251	GLN
1	B	273	ASN
1	B	309	GLN
1	B	406	ASN
1	B	436	GLN
1	B	544	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1JG	A	601	-	20,21,21	2.33	11 (55%)	26,31,31	1.64	4 (15%)
2	1JG	B	601	-	20,21,21	2.07	10 (50%)	26,31,31	1.36	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1JG	A	601	-	-	0/9/15/15	0/2/2/2
2	1JG	B	601	-	-	0/9/15/15	0/2/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	1JG	S7-N10	4.52	1.71	1.63
2	B	601	1JG	C3-S2	-4.24	1.69	1.73
2	B	601	1JG	S7-N10	4.09	1.70	1.63
2	A	601	1JG	O19-C17	4.05	1.35	1.23
2	A	601	1JG	C3-S2	-3.55	1.70	1.73
2	A	601	1JG	O9-S7	3.43	1.47	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	1JG	O9-S7	3.19	1.47	1.43
2	A	601	1JG	C13-C12	2.85	1.44	1.39
2	A	601	1JG	O8-S7	2.65	1.46	1.43
2	A	601	1JG	C12-C11	2.65	1.45	1.41
2	B	601	1JG	C13-C12	2.63	1.44	1.39
2	B	601	1JG	C16-C11	2.51	1.43	1.39
2	A	601	1JG	O18-C17	-2.44	1.22	1.30
2	A	601	1JG	C13-C14	2.24	1.43	1.38
2	A	601	1JG	C16-C11	2.15	1.43	1.39
2	A	601	1JG	C14-C15	2.14	1.42	1.38
2	B	601	1JG	O8-S7	2.10	1.45	1.43
2	B	601	1JG	C13-C14	2.08	1.42	1.38
2	B	601	1JG	C12-C11	2.07	1.44	1.41
2	B	601	1JG	C16-C15	2.07	1.41	1.38
2	B	601	1JG	C14-C15	2.01	1.41	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	1JG	C6-S7-N10	3.37	112.21	107.11
2	A	601	1JG	C6-S7-N10	3.33	112.14	107.11
2	A	601	1JG	O19-C17-C12	-3.32	113.48	121.82
2	A	601	1JG	O18-C17-C12	3.30	125.32	115.47
2	A	601	1JG	C4-C3-S2	2.87	115.06	110.21
2	B	601	1JG	C4-C3-S2	2.68	114.75	110.21
2	B	601	1JG	O8-S7-C6	-2.12	104.33	108.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	559/576 (97%)	-0.14	12 (2%)	60 61	17, 28, 51, 80	0
1	B	558/576 (96%)	0.20	33 (5%)	22 21	18, 34, 63, 83	0
All	All	1117/1152 (96%)	0.03	45 (4%)	36 36	17, 31, 57, 83	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	23	ILE	6.5
1	B	153	GLY	5.6
1	B	26	LEU	5.4
1	B	543	SER	4.7
1	B	24	ASN	4.6
1	A	149	PRO	4.4
1	B	27	SER	4.3
1	B	548	SER	4.2
1	B	545	LEU	4.2
1	B	29	SER	4.1
1	B	546	ASP	3.9
1	A	531	ARG	3.7
1	A	545	LEU	3.7
1	B	532	THR	3.7
1	A	543	SER	3.6
1	A	14	CYS	3.5
1	B	25	PRO	3.5
1	B	511	LEU	3.5
1	B	531	ARG	3.4
1	A	544	GLN	3.1
1	B	533	LYS	3.1
1	B	534	LEU	3.0
1	B	535	LYS	2.8
1	A	402	HIS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	14	CYS	2.7
1	B	563	SER	2.6
1	B	540	PRO	2.5
1	B	330	GLN	2.5
1	B	544	GLN	2.5
1	A	95	HIS	2.4
1	B	16	ALA	2.4
1	A	15	ALA	2.4
1	B	515	GLY	2.3
1	A	547	LEU	2.3
1	A	535	LYS	2.3
1	B	434	LEU	2.2
1	B	212	LYS	2.2
1	B	57	LEU	2.2
1	B	541	ALA	2.2
1	B	523	ARG	2.1
1	B	106	LYS	2.1
1	B	213	CYS	2.1
1	B	402	HIS	2.1
1	A	548	SER	2.1
1	B	433	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	1JG	B	601	20/20	0.09	-0.66	44,46,47,50	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	1JG	A	601	20/20	0.08	-1.26	29,34,39,40	0

6.5 Other polymers ⓘ

There are no such residues in this entry.