



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 02:58 PM GMT

PDB ID : 4J16
Title : Crystal structure of Thermus thermophilus transhydrogenase heterotrimeric complex of the Alpha1 subunit dimer with the NADP binding domain (domain III) of the Beta subunit
Authors : Yamaguchi, M.; Leung, J.; Schurig Briccio, L.A.; Gennis, R.B.; Stout, C.D.
Deposited on : 2013-02-01
Resolution : 2.41 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2496 (2.44-2.40)
Clashscore	79885	3124 (2.44-2.40)
Ramachandran outliers	78287	3067 (2.44-2.40)
Sidechain outliers	78261	3068 (2.44-2.40)
RSRZ outliers	66119	2499 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	381	
1	B	381	
2	C	185	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NAD	A	500	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7186 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD/NADP transhydrogenase alpha subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	0	0
			2794	1772	498	512	12			
1	B	369	Total	C	N	O	S	0	0	0
			2769	1756	495	506	12			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-4	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-3	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-2	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	-1	HIS	-	EXPRESSION TAG	UNP Q72GR8
A	0	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-5	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-4	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-3	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-2	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	-1	HIS	-	EXPRESSION TAG	UNP Q72GR8
B	0	HIS	-	EXPRESSION TAG	UNP Q72GR8

- Molecule 2 is a protein called NAD(P) transhydrogenase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	177	Total	C	N	O	S	0	0	0
			1343	860	229	248	6			

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	52	Total	O	0	0
			52	52		
8	B	35	Total	O	0	0
			35	35		
8	C	34	Total	O	0	0
			34	34		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.62Å 75.04Å 198.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.98 – 2.41 39.98 – 2.41	Depositor EDS
% Data completeness (in resolution range)	96.3 (39.98-2.41) 96.3 (39.98-2.41)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.201 , 0.253 0.201 , 0.254	Depositor DCC
R_{free} test set	1924 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	58.4	Xtriage
Anisotropy	0.366	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 38562 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7186	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAP, PGE, NAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2840	0.66	2/3852 (0.1%)
1	B	0.35	0/2813	0.59	0/3814
2	C	0.39	0/1366	0.61	0/1848
All	All	0.37	0/7019	0.62	2/9514 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	274	ARG	NE-CZ-NH1	7.42	124.01	120.30
1	A	353	ARG	NE-CZ-NH2	6.36	123.48	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2794	0	0	3	0
1	B	2769	0	0	12	0
2	C	1343	0	0	3	0
3	A	44	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	44	0	0	1	0
4	B	1	0	0	0	0
5	B	10	0	14	1	0
6	B	6	0	8	0	0
6	C	6	0	8	0	0
7	C	48	0	0	0	0
8	A	52	0	0	0	0
8	B	35	0	0	0	0
8	C	34	0	0	1	0
All	All	7186	0	30	19	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (19) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:C:278:LYS:NZ	2:C:426:ALA:O	2.40	0.54
1:B:285:ALA:O	1:B:289:GLY:O	2.28	0.52
1:B:216:LEU:CB	1:B:217:PRO:CA	2.91	0.48
1:A:278:GLY:N	1:A:305:GLY:O	2.48	0.46
1:B:314:LEU:N	1:B:315:PRO:CD	2.78	0.46
1:B:199:ARG:NH1	3:B:401:NAD:O3D	2.49	0.46
1:B:331:TYR:O	1:B:334:SER:OG	2.35	0.45
5:B:403:PGE:H42	5:B:403:PGE:H2	1.72	0.44
1:B:35:ARG:NH1	1:B:63:GLU:O	2.51	0.44
1:A:268:THR:OG1	1:A:270:ASP:OD1	2.36	0.43
1:B:33:ARG:NH2	1:B:57:GLU:OE1	2.52	0.43
3:A:500:NAD:O2A	3:A:500:NAD:O3B	2.38	0.42
2:C:428:ASN:N	8:C:629:HOH:O	2.52	0.42
1:B:234:GLU:O	1:B:236:ARG:N	2.53	0.42
1:B:365:HIS:ND1	1:B:368:THR:OG1	2.54	0.41
1:B:268:THR:OG1	1:B:270:ASP:OD1	2.39	0.41
1:A:218:ILE:O	1:A:238:GLN:NE2	2.53	0.41
1:B:150:LEU:O	1:B:307:ARG:NH1	2.54	0.41
1:B:190:GLY:O	2:C:352:TYR:OH	2.38	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/381 (97%)	352 (95%)	18 (5%)	0	100	100
1	B	365/381 (96%)	345 (94%)	15 (4%)	5 (1%)	16	21
2	C	175/185 (95%)	172 (98%)	3 (2%)	0	100	100
All	All	910/947 (96%)	869 (96%)	36 (4%)	5 (0%)	38	52

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	216	LEU
1	B	228	ARG
1	B	235	LYS
1	B	217	PRO
1	B	215	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/293 (98%)	257 (90%)	30 (10%)	10	14
1	B	285/293 (97%)	259 (91%)	26 (9%)	14	19
2	C	139/146 (95%)	130 (94%)	9 (6%)	24	36
All	All	711/732 (97%)	646 (91%)	65 (9%)	14	19

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	4	VAL
1	A	60	GLU
1	A	71	LEU
1	A	79	GLU
1	A	101	LEU
1	A	114	VAL
1	A	115	ILE
1	A	123	ILE
1	A	124	THR
1	A	149	ARG
1	A	158	LEU
1	A	200	LYS
1	A	203	LEU
1	A	223	GLU
1	A	229	GLU
1	A	230	LEU
1	A	232	GLU
1	A	235	LYS
1	A	261	ARG
1	A	274	ARG
1	A	276	LYS
1	A	303	VAL
1	A	304	ARG
1	A	314	LEU
1	A	333	LEU
1	A	336	LEU
1	A	348	GLU
1	A	353	ARG
1	A	362	GLU
1	B	1	MET
1	B	10	ARG
1	B	49	GLU
1	B	71	LEU
1	B	86	GLU
1	B	98	HIS
1	B	99	LYS
1	B	105	ARG
1	B	123	ILE
1	B	125	ARG
1	B	142	LEU
1	B	149	ARG
1	B	175	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	203	LEU
1	B	212	LYS
1	B	228	ARG
1	B	230	LEU
1	B	231	THR
1	B	242	LEU
1	B	246	VAL
1	B	262	ARG
1	B	314	LEU
1	B	333	LEU
1	B	336	LEU
1	B	350	GLU
1	B	364	LEU
2	C	277	LEU
2	C	325	GLU
2	C	355	LEU
2	C	374	ILE
2	C	397	LEU
2	C	427	GLU
2	C	432	LEU
2	C	437	GLN
2	C	447	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAD	A	500	-	48,48,48	1.85	7 (14%)	73,73,73	2.02	9 (12%)
3	NAD	B	401	-	48,48,48	1.71	6 (12%)	73,73,73	2.05	14 (19%)
5	PGE	B	403	-	9,9,9	0.65	0	8,8,8	0.44	0
6	GOL	B	404	-	5,5,5	0.39	0	5,5,5	0.97	0
7	NAP	C	501	-	52,52,52	1.81	8 (15%)	80,80,80	2.30	21 (26%)
6	GOL	C	502	-	5,5,5	0.35	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	500	-	-	0/30/62/62	0/3/5/5
3	NAD	B	401	-	-	0/30/62/62	0/3/5/5
5	PGE	B	403	-	-	0/7/7/7	0/0/0/0
6	GOL	B	404	-	-	0/4/4/4	0/0/0/0
7	NAP	C	501	-	-	0/35/67/67	0/3/5/5
6	GOL	C	502	-	-	0/4/4/4	0/0/0/0

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	NAD	PN-O2N	7.06	1.65	1.48
7	C	501	NAP	PN-O2N	6.97	1.64	1.48
3	A	500	NAD	PN-O1N	6.97	1.65	1.48
3	B	401	NAD	PN-O2N	6.87	1.64	1.48
7	C	501	NAP	PN-O1N	6.69	1.65	1.48
3	B	401	NAD	PN-O1N	6.47	1.64	1.48
7	C	501	NAP	P2B-O1X	4.09	1.64	1.51
3	A	500	NAD	PA-O1A	3.83	1.66	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	NAD	PA-O1A	3.51	1.64	1.51
3	A	500	NAD	PA-O3	3.33	1.65	1.59
7	C	501	NAP	PA-O1A	2.92	1.62	1.51
7	C	501	NAP	P2B-O3X	2.71	1.64	1.54
3	B	401	NAD	PA-O3	2.69	1.64	1.59
7	C	501	NAP	P2B-O2X	2.52	1.63	1.54
3	A	500	NAD	PN-O3	2.36	1.65	1.60
3	A	500	NAD	PA-O2A	2.33	1.65	1.55
7	C	501	NAP	PA-O2A	2.24	1.65	1.55
3	B	401	NAD	PN-O3	2.09	1.64	1.60
3	B	401	NAD	PA-O2A	2.09	1.64	1.55
3	A	500	NAD	O4B-C1B	2.06	1.44	1.41
7	C	501	NAP	PN-O3	2.06	1.64	1.60

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	501	NAP	N3A-C2A-N1A	-10.52	119.92	128.71
3	A	500	NAD	O4B-C1B-N9A	9.61	117.38	108.44
3	B	401	NAD	O4B-C1B-N9A	8.28	116.14	108.44
3	A	500	NAD	N3A-C2A-N1A	-8.27	121.79	128.71
3	B	401	NAD	N3A-C2A-N1A	-7.74	122.24	128.71
7	C	501	NAP	C3N-C7N-N7N	7.55	126.36	117.77
7	C	501	NAP	O4B-C1B-C2B	-5.70	101.61	106.95
3	B	401	NAD	O4D-C1D-N1N	5.45	113.53	107.95
3	B	401	NAD	N3A-C4A-N9A	4.96	134.39	125.43
3	A	500	NAD	N3A-C4A-N9A	4.95	134.36	125.43
7	C	501	NAP	C4D-O4D-C1D	4.59	114.74	109.75
7	C	501	NAP	O3B-C3B-C4B	-4.03	99.21	111.08
7	C	501	NAP	O7N-C7N-C3N	-4.00	115.06	119.58
7	C	501	NAP	N3A-C4A-N9A	3.89	132.46	125.43
3	A	500	NAD	C4A-C5A-N7A	-3.85	106.22	109.52
3	A	500	NAD	C5A-C4A-N3A	-3.76	117.52	125.70
3	B	401	NAD	C4A-C5A-N7A	-3.73	106.33	109.52
3	B	401	NAD	C5A-C4A-N3A	-3.53	118.02	125.70
7	C	501	NAP	C4A-C5A-N7A	-3.34	106.66	109.52
7	C	501	NAP	O4D-C1D-C2D	-3.28	101.74	106.77
3	B	401	NAD	O4D-C1D-C2D	-3.27	101.77	106.77
7	C	501	NAP	O4B-C4B-C3B	-3.18	98.72	105.17
7	C	501	NAP	O3D-C3D-C4D	-3.18	101.71	111.08
3	A	500	NAD	PN-O3-PA	-3.16	119.39	132.95
3	B	401	NAD	PN-O3-PA	-2.91	120.46	132.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	NAD	C3D-C2D-C1D	2.89	105.43	100.91
7	C	501	NAP	P2B-O2B-C2B	2.88	128.01	121.96
3	B	401	NAD	C2B-C1B-N9A	-2.81	106.04	113.27
7	C	501	NAP	O4D-C1D-N1N	-2.79	105.10	107.95
7	C	501	NAP	O7N-C7N-N7N	-2.78	118.57	122.59
3	A	500	NAD	C2A-N3A-C4A	2.71	121.72	114.01
7	C	501	NAP	C5A-C4A-N3A	-2.65	119.93	125.70
3	B	401	NAD	C4D-O4D-C1D	-2.65	106.87	109.75
7	C	501	NAP	C2B-C3B-C4B	2.63	108.18	101.94
7	C	501	NAP	C2A-N3A-C4A	2.54	121.25	114.01
3	B	401	NAD	C2A-N3A-C4A	2.40	120.83	114.01
7	C	501	NAP	O2D-C2D-C3D	-2.29	104.37	111.83
7	C	501	NAP	C6N-N1N-C2N	-2.23	119.52	122.04
3	B	401	NAD	O4B-C4B-C5B	-2.21	101.46	109.36
3	A	500	NAD	C2B-C1B-N9A	-2.11	107.85	113.27
3	B	401	NAD	C2D-C1D-N1N	-2.09	110.32	113.86
3	B	401	NAD	C5D-C4D-C3D	-2.06	106.95	115.21
7	C	501	NAP	C2B-C1B-N9A	-2.01	106.10	113.74
7	C	501	NAP	PN-O5D-C5D	2.01	127.18	120.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	372/381 (97%)	0.48	29 (7%)	13 11	39, 70, 103, 120	0
1	B	369/381 (96%)	0.36	18 (4%)	28 25	42, 77, 114, 146	0
2	C	177/185 (95%)	-0.16	2 (1%)	77 77	40, 59, 79, 96	0
All	All	918/947 (96%)	0.31	49 (5%)	25 23	39, 69, 107, 146	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	230	LEU	5.6
1	A	33	ARG	5.4
1	A	371	LEU	3.3
1	A	145	ILE	3.2
1	A	304	ARG	3.1
1	A	370	ALA	3.0
1	A	185	THR	2.9
1	B	99	LYS	2.9
1	A	112	ALA	2.9
1	B	262	ARG	2.9
1	B	189	LEU	2.8
1	B	215	GLU	2.8
1	A	138	VAL	2.6
1	A	1	MET	2.6
1	A	141	TYR	2.6
1	A	159	THR	2.5
1	A	188	ARG	2.5
1	A	261	ARG	2.5
1	A	189	LEU	2.5
1	A	157	MET	2.5
1	B	260	GLY	2.5
1	A	101	LEU	2.5
1	A	38	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	186	ALA	2.4
1	B	373	GLY	2.4
1	B	219	SER	2.4
1	A	360	GLU	2.4
1	A	158	LEU	2.4
1	A	109	ALA	2.4
1	A	184	ALA	2.4
1	B	144	ALA	2.3
1	B	228	ARG	2.3
1	B	141	TYR	2.3
1	B	145	ILE	2.3
1	A	140	GLY	2.3
1	A	363	VAL	2.3
1	A	342	ALA	2.2
1	B	259	PRO	2.2
1	A	156	PRO	2.2
1	B	242	LEU	2.1
1	A	137	THR	2.1
1	B	368	THR	2.1
1	A	62	GLY	2.1
1	A	269	GLU	2.1
2	C	396	ILE	2.1
1	B	142	LEU	2.0
1	B	322	ALA	2.0
2	C	337	MET	2.0
1	B	190	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAD	A	500	44/44	0.40	4.77	103,120,160,164	0
4	CL	B	402	1/1	0.27	1.36	83,83,83,83	0
6	GOL	C	502	6/6	0.18	0.52	83,89,90,92	0
3	NAD	B	401	44/44	0.18	-0.06	67,76,91,94	0
7	NAP	C	501	48/48	0.14	-0.08	43,47,60,63	0
5	PGE	B	403	10/10	0.16	-0.21	65,72,75,78	0
6	GOL	B	404	6/6	0.22	-0.26	53,57,60,61	0

6.5 Other polymers ⓘ

There are no such residues in this entry.