



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 08:55 PM GMT

PDB ID : 2J2F
Title : THE T199D MUTANT OF STEAROYL ACYL CARRIER PROTEIN DESATURASE FROM RICINUS COMMUNIS (CASTOR BEAN)
Authors : Guy, J.E.; Abreu, I.A.; Moche, M.; Lindqvist, Y.; Whittle, E.; Shanklin, J.
Deposited on : 2006-08-16
Resolution : 2.65 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance







The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2232 (2.70-2.62)
Clashscore	79885	2700 (2.70-2.62)
Ramachandran outliers	78287	2657 (2.70-2.62)
Sidechain outliers	78261	2657 (2.70-2.62)
RSRZ outliers	66119	2234 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	363	
1	B	363	
1	C	363	
1	D	363	
1	E	363	
1	F	363	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 17055 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACYL-[ACYL-CARRIER-PROTEIN]DESATURASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	348	Total	C	N	O	S	0	0	0
			2824	1791	490	529	14			
1	B	347	Total	C	N	O	S	0	0	0
			2815	1785	488	528	14			
1	C	346	Total	C	N	O	S	0	0	0
			2808	1780	487	527	14			
1	D	347	Total	C	N	O	S	0	0	0
			2815	1785	488	528	14			
1	E	347	Total	C	N	O	S	0	0	0
			2815	1785	488	528	14			
1	F	346	Total	C	N	O	S	0	0	0
			2808	1780	487	527	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	ASP	THR	ENGINEERED MUTATION	UNP P22337
B	199	ASP	THR	ENGINEERED MUTATION	UNP P22337
C	199	ASP	THR	ENGINEERED MUTATION	UNP P22337
D	199	ASP	THR	ENGINEERED MUTATION	UNP P22337
E	199	ASP	THR	ENGINEERED MUTATION	UNP P22337
F	199	ASP	THR	ENGINEERED MUTATION	UNP P22337

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	2	Total	Fe	0	0
			2	2		
2	E	2	Total	Fe	0	0
			2	2		
2	B	2	Total	Fe	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	2	Total 2	Fe 2	0	0
2	A	2	Total 2	Fe 2	0	0
2	F	2	Total 2	Fe 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	20	Total 20	O 20	0	0
3	B	38	Total 38	O 38	0	0
3	C	27	Total 27	O 27	0	0
3	D	33	Total 33	O 33	0	0
3	E	20	Total 20	O 20	0	0
3	F	20	Total 20	O 20	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACYL-[ACYL-CARRIER-PROTEIN]DESATURASE




• Molecule 1: ACYL-[ACYL-CARRIER-PROTEIN]DESATURASE




• Molecule 1: ACYL-[ACYL-CARRIER-PROTEIN]DESATURASE






- Molecule 1: ACYL-[ACYL-CARRIER-PROTEIN]DESATURASE

Chain D:




- Molecule 1: ACYL-[ACYL-CARRIER-PROTEIN]DESATURASE

Chain E:



- Molecule 1: ACYL-[ACYL-CARRIER-PROTEIN]DESATURASE

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.05Å 145.77Å 193.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.65 33.67 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.4 (40.00-2.65) 97.9 (33.67-2.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.240 , 0.272 0.233 , 0.263	Depositor DCC
R_{free} test set	3388 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	43.2	Xtriage
Anisotropy	0.476	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 9.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 66623 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	17055	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 16.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.65	0/2893	0.69	4/3916 (0.1%)
1	B	0.80	0/2884	0.78	4/3904 (0.1%)
1	C	0.75	0/2876	0.75	5/3893 (0.1%)
1	D	0.79	1/2884 (0.0%)	0.77	2/3904 (0.1%)
1	E	0.71	0/2884	0.71	0/3904
1	F	0.69	0/2876	0.72	5/3893 (0.1%)
All	All	0.73	1/17297 (0.0%)	0.74	20/23414 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	82	GLU	CG-CD	5.80	1.60	1.51

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	86	ARG	NE-CZ-NH2	-9.38	115.61	120.30
1	C	86	ARG	NE-CZ-NH2	-9.13	115.73	120.30
1	B	86	ARG	NE-CZ-NH1	8.75	124.67	120.30
1	C	84	ARG	NE-CZ-NH2	-8.62	115.99	120.30
1	F	274	ARG	NE-CZ-NH1	7.90	124.25	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens

added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2824	0	2765	83	2
1	B	2815	0	2753	70	1
1	C	2808	0	2745	80	0
1	D	2815	0	2753	74	0
1	E	2815	0	2753	79	0
1	F	2808	0	2745	71	1
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	20	0	0	1	0
3	B	38	0	0	2	0
3	C	27	0	0	3	0
3	D	33	0	0	2	0
3	E	20	0	0	5	0
3	F	20	0	0	9	0
All	All	17055	0	16514	439	3

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 439 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:16:LYS:HG2	1:A:19:MET:HE3	1.39	1.04
1:A:16:LYS:HG2	1:A:19:MET:CE	1.90	1.02
1:E:217:LYS:HE2	3:E:2018:HOH:O	1.67	0.95
1:D:114:MET:CE	1:D:177:MET:HB3	1.96	0.95
1:B:59:GLU:HG3	1:C:18:PHE:CD2	2.03	0.94

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:25:HIS:CD2	1:A:49:GLU:OE2[4_555]	2.15	0.05
1:A:25:HIS:CD2	1:A:49:GLU:OE1[4_555]	2.18	0.02
1:B:339:GLU:OE1	1:F:35:GLN:OE1[4_555]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/363 (95%)	317 (92%)	23 (7%)	6 (2%)	14	30
1	B	345/363 (95%)	324 (94%)	17 (5%)	4 (1%)	19	42
1	C	344/363 (95%)	321 (93%)	19 (6%)	4 (1%)	19	42
1	D	345/363 (95%)	316 (92%)	25 (7%)	4 (1%)	19	42
1	E	345/363 (95%)	317 (92%)	24 (7%)	4 (1%)	19	42
1	F	344/363 (95%)	317 (92%)	22 (6%)	5 (2%)	15	34
All	All	2069/2178 (95%)	1912 (92%)	130 (6%)	27 (1%)	18	39

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	17	PRO
1	A	213	HIS
1	A	340	ARG
1	B	213	HIS
1	B	340	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/315 (96%)	286 (95%)	16 (5%)	32	60
1	B	301/315 (96%)	284 (94%)	17 (6%)	30	56
1	C	300/315 (95%)	279 (93%)	21 (7%)	21	44
1	D	301/315 (96%)	284 (94%)	17 (6%)	30	56
1	E	301/315 (96%)	281 (93%)	20 (7%)	24	47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	F	300/315 (95%)	284 (95%)	16 (5%)	32 60
All	All	1805/1890 (96%)	1698 (94%)	107 (6%)	28 54

5 of 107 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	340	ARG
1	D	177	MET
1	F	270	MET
1	C	342	GLN
1	D	31	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	342	GLN
1	D	50	ASN
1	F	268	HIS
1	D	35	GLN
1	D	268	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	348/363 (95%)	1.52	110 (31%) 1 1	31, 33, 34, 50	0
1	B	347/363 (95%)	0.94	46 (13%) 4 3	31, 33, 34, 48	0
1	C	346/363 (95%)	0.98	45 (13%) 4 3	22, 33, 34, 36	0
1	D	347/363 (95%)	0.91	39 (11%) 6 5	31, 33, 34, 44	0
1	E	347/363 (95%)	1.18	75 (21%) 1 1	31, 33, 34, 50	0
1	F	346/363 (95%)	1.78	130 (37%) 1 1	31, 33, 34, 45	0
All	All	2081/2178 (95%)	1.22	445 (21%) 1 1	22, 33, 34, 50	0

The worst 5 of 445 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	345	ALA	11.6
1	B	345	ALA	9.2
1	D	344	ARG	9.0
1	F	341	ALA	8.9
1	E	345	ALA	8.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FE	D	370	1/1	0.13	-1.76	48,48,48,48	0
2	FE	E	370	1/1	0.13	-2.02	48,48,48,48	0
2	FE	B	370	1/1	0.12	-2.65	48,48,48,48	0
2	FE	C	371	1/1	0.06	-2.79	50,50,50,50	0
2	FE	D	371	1/1	0.06	-3.26	50,50,50,50	0
2	FE	B	371	1/1	0.11	-3.40	49,49,49,49	0
2	FE	C	370	1/1	0.07	-3.95	48,48,48,48	0
2	FE	F	370	1/1	0.06	-4.03	48,48,48,48	0
2	FE	E	371	1/1	0.02	-4.04	49,49,49,49	0
2	FE	A	370	1/1	0.07	-4.07	48,48,48,48	0
2	FE	A	371	1/1	0.05	-4.13	50,50,50,50	0
2	FE	F	371	1/1	0.04	-5.28	49,49,49,49	0

6.5 Other polymers

There are no such residues in this entry.