



wwPDB X-ray Structure Validation Summary Report i

Feb 26, 2014 – 09:03 PM GMT

PDB ID : 4J5M
Title : Structure of the Cargo Binding Domain from Human Myosin Vb
Authors : Nascimento, A.F.Z.; Trindade, D.M.; Mahajan, P.; Berridge, G.; Krojer, T.; Vollmar, M.; Tonoli, C.C.C.; Assis, L.H.P.; Burgess-Brown, N.; von Delft, F.; Murakami, M.T.
Deposited on : 2013-02-08
Resolution : 2.07 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance


The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1224 (2.08-2.04)
Clashscore	79885	1390 (2.08-2.04)
Ramachandran outliers	78287	1381 (2.08-2.04)
Sidechain outliers	78261	1381 (2.08-2.04)
RSRZ outliers	66119	1225 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	396	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NO3	A	1901	X	-
2	NO3	A	1902	X	X

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 3181 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-Vb.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	363	2967	1888	502	551	26	2	6	0

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	N	O		
2	A	1	4	1	3	0	0
2	A	1	4	1	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	206	Total	O	0	0
			206	206		

i

- Molecule 1: Unconventional myosin-Vb

Protein	Protein ID	Protein Name	Protein Type	Protein Class	Protein Subclass	Protein Family	Protein Subfamily	Protein Domain	Protein Subdomain	Protein Motif	Protein Signature	Protein Annotation
LEU	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN
ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN
ARG	GLN	GLN	GLN	GLN	GLN	GLN	GLN	GLN	GLN	GLN	GLN	GLN
GLN	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL	VAL
VAL	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR	THR
THR	Q1460	Q1460	Q1460	Q1460	Q1460	Q1460	Q1460	Q1460	Q1460	Q1460	Q1460	Q1460
Q1460	E1463	E1463	E1463	E1463	E1463	E1463	E1463	E1463	E1463	E1463	E1463	E1463
E1463	K1464	K1464	K1464	K1464	K1464	K1464	K1464	K1464	K1464	K1464	K1464	K1464
K1464	D1465	D1465	D1465	D1465	D1465	D1465	D1465	D1465	D1465	D1465	D1465	D1465
D1465	F1466	F1466	F1466	F1466	F1466	F1466	F1466	F1466	F1466	F1466	F1466	F1466
F1466	E1477	E1477	E1477	E1477	E1477	E1477	E1477	E1477	E1477	E1477	E1477	E1477
E1477	L1493	L1493	L1493	L1493	L1493	L1493	L1493	L1493	L1493	L1493	L1493	L1493
L1493	S1494	S1494	S1494	S1494	S1494	S1494	S1494	S1494	S1494	S1494	S1494	S1494
S1494	G1495	G1495	G1495	G1495	G1495	G1495	G1495	G1495	G1495	G1495	G1495	G1495
G1495	T1496	T1496	T1496	T1496	T1496	T1496	T1496	T1496	T1496	T1496	T1496	T1496
T1496	L1500	L1500	L1500	L1500	L1500	L1500	L1500	L1500	L1500	L1500	L1500	L1500
L1500	L1505	L1505	L1505	L1505	L1505	L1505	L1505	L1505	L1505	L1505	L1505	L1505
L1505	L1519	L1519	L1519	L1519	L1519	L1519	L1519	L1519	L1519	L1519	L1519	L1519
L1519	K1520	K1520	K1520	K1520	K1520	K1520	K1520	K1520	K1520	K1520	K1520	K1520
K1520	V1521	V1521	V1521	V1521	V1521	V1521	V1521	V1521	V1521	V1521	V1521	V1521
V1521	H1522	H1522	H1522	H1522	H1522	H1522	H1522	H1522	H1522	H1522	H1522	H1522
H1522	S1523	S1523	S1523	S1523	S1523	S1523	S1523	S1523	S1523	S1523	S1523	S1523
S1523	Q1572	Q1572	Q1572	Q1572	Q1572	Q1572	Q1572	Q1572	Q1572	Q1572	Q1572	Q1572
Q1572	K1576	K1576	K1576	K1576	K1576	K1576	K1576	K1576	K1576	K1576	K1576	K1576
K1576	Y1590	Y1590	Y1590	Y1590	Y1590	Y1590	Y1590	Y1590	Y1590	Y1590	Y1590	Y1590
Y1590	R1591	R1591	R1591	R1591	R1591	R1591	R1591	R1591	R1591	R1591	R1591	R1591
R1591	Q1592	Q1592	Q1592	Q1592	Q1592	Q1592	Q1592	Q1592	Q1592	Q1592	Q1592	Q1592
Q1592	E1623	E1623	E1623	E1623	E1623	E1623	E1623	E1623	E1623	E1623	E1623	E1623
E1623	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN	ASN
ASN	GLU	GLU	GLU	GLU	GLU	GLU	GLU	GLU	GLU	GLU	GLU	GLU
GLU	SER	SER	SER	SER								

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	69.09Å 78.20Å 151.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.01 – 2.07 49.01 – 2.07	Depositor EDS
% Data completeness (in resolution range)	97.1 (49.01-2.07) 99.5 (49.01-2.07)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.07Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.189 , 0.232 0.182 , 0.221	Depositor DCC
R_{free} test set	1295 reflections (5.10%)	DCC
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 25404 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3181	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3036	0.50	0/4104

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2967	0	0	12	0
2	A	8	0	0	0	0
3	A	206	0	0	5	0
All	All	3181	0	0	12	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

The worst 5 of 12 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1727:ASN:O	1:A:1730:GLN:NE2	2.28	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1576:LYS:NZ	1:A:1832:ASP:OD2	2.32	0.62
1:A:1805:ASN:N	1:A:1805:ASN:OD1	2.37	0.57
1:A:1786:GLU:OE1	3:A:2050:HOH:O	2.18	0.57
1:A:1659:ARG:NH2	3:A:2143:HOH:O	2.38	0.56

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	365/396 (92%)	357 (98%)	5 (1%)	3 (1%)	27	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1832	ASP
1	A	1807	PRO
1	A	1806	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/363 (94%)	329 (96%)	12 (4%)	48	40

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1519[B]	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1592[A]	GLN
1	A	1782	LEU
1	A	1519[A]	LEU
1	A	1657	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NO3	A	1901	-	3,3,3	3.22	3 (100%)	3,3,3	0.25	0
2	NO3	A	1902	-	3,3,3	3.19	3 (100%)	3,3,3	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NO3	A	1901	-	-	0/0/0/0	0/0/0/0
2	NO3	A	1902	-	-	0/0/0/0	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1902	NO3	O1-N	3.85	1.40	1.24
2	A	1901	NO3	O1-N	3.78	1.40	1.24
2	A	1901	NO3	O3-N	2.97	1.41	1.25
2	A	1902	NO3	O2-N	2.89	1.40	1.25
2	A	1901	NO3	O2-N	2.82	1.40	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/396 (91%)	-0.37	6 (1%) 67 70	15, 32, 74, 100	12 (3%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1460	GLN	4.6
1	A	1459	VAL	4.4
1	A	1464	LYS	2.8
1	A	1463	GLU	2.7
1	A	1807	PRO	2.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NO3	A	1902	4/4	0.17	7.95	45,50,52,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NO3	A	1901	4/4	0.14	1.45	36,39,43,45	0

6.5 Other polymers ⓘ

There are no such residues in this entry.