



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 02:29 PM GMT

PDB ID : 4J5T  
Title : Crystal structure of Processing alpha-Glucosidase I  
Authors : Barker, M.K.; Rose, D.R.  
Deposited on : 2013-02-09  
Resolution : 2.04 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1380 (2.06-2.02)
Clashscore	79885	1577 (2.06-2.02)
Ramachandran outliers	78287	1565 (2.06-2.02)
Sidechain outliers	78261	1565 (2.06-2.02)
RSRZ outliers	66119	1381 (2.06-2.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	811	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NAG	A	1002	-	X
2	NAG	A	1004	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6838 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mannosyl-oligosaccharideglucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	788	6497	4163	1074	1233	27	0	7	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	ALA	-	EXPRESSION TAG	UNP P53008
A	-1	GLU	-	EXPRESSION TAG	UNP P53008
A	0	PHE	-	EXPRESSION TAG	UNP P53008
A	1	MET	-	EXPRESSION TAG	UNP P53008
A	801	ARG	-	EXPRESSION TAG	UNP P53008
A	802	SER	-	EXPRESSION TAG	UNP P53008
A	803	HIS	-	EXPRESSION TAG	UNP P53008
A	804	HIS	-	EXPRESSION TAG	UNP P53008
A	805	HIS	-	EXPRESSION TAG	UNP P53008
A	806	HIS	-	EXPRESSION TAG	UNP P53008
A	807	HIS	-	EXPRESSION TAG	UNP P53008
A	808	HIS	-	EXPRESSION TAG	UNP P53008

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	285	Total	O	0	0
			285	285		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mannosyl-oligosaccharideglucosidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.80Å 101.80Å 103.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.00 – 2.04 27.86 – 2.04	Depositor EDS
% Data completeness (in resolution range)	97.8 (28.00-2.04) 97.8 (27.86-2.04)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 2.04Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.226 , 0.235 0.225 , 0.249	Depositor DCC
$R_{free}$ test set	3176 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	36.5	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 49.4	EDS
Estimated twinning fraction	0.002 for -h,l,k	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 62638 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6838	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/6675	0.47	0/9047

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6497	0	6192	49	0
2	A	56	0	52	4	0
3	A	285	0	0	1	0
All	All	6838	0	6244	49	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (49) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:9:ASN:HD21	2:A:1001:NAG:C1	1.46	1.29

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:89:ASN:HD21	2:A:1003:NAG:C1	1.68	1.06
1:A:581:LEU:HD12	1:A:638:ILE:HD11	1.57	0.87
1:A:9:ASN:ND2	2:A:1001:NAG:O5	2.13	0.80
1:A:392:ASP:OD1	1:A:789:TRP:CZ3	2.45	0.69
1:A:532:GLU:OE1	1:A:579:ALA:HB2	1.96	0.66
1:A:428:ARG:HB3	1:A:453:ASN:OD1	1.99	0.63
1:A:151:LYS:HD3	1:A:165:THR:HG22	1.82	0.62
1:A:89:ASN:HD21	2:A:1003:NAG:C2	2.13	0.61
1:A:537:LEU:HD13	1:A:546:ILE:HD13	1.83	0.60
1:A:458:LEU:HD21	1:A:515:LEU:HD22	1.85	0.57
1:A:530:ILE:O	1:A:532:GLU:O	2.24	0.56
1:A:25:MET:HG3	1:A:37:MET:HE1	1.88	0.55
1:A:37:MET:HG3	1:A:126:PHE:CE1	2.43	0.53
1:A:530:ILE:HB	1:A:551:VAL:HG21	1.91	0.52
1:A:397:LEU:HD13	1:A:412[A]:LEU:HG	1.92	0.51
1:A:532:GLU:OE1	1:A:579:ALA:CB	2.59	0.51
1:A:425:TRP:CE2	1:A:449:PRO:HA	2.46	0.51
1:A:573:ALA:HB2	1:A:651:VAL:HG21	1.94	0.50
1:A:161:GLU:OE1	1:A:278:LEU:CD2	2.60	0.50
1:A:532:GLU:O	1:A:533:TYR:HB2	2.12	0.48
1:A:519:TYR:CZ	1:A:523:ARG:HD2	2.50	0.47
1:A:405:PHE:CE2	1:A:409:PHE:HE1	2.33	0.47
1:A:405:PHE:CE2	1:A:409:PHE:CE1	3.03	0.46
1:A:198:LEU:CD1	1:A:200:ILE:HG12	2.45	0.46
1:A:390:TYR:OH	1:A:518:HIS:NE2	2.38	0.45
1:A:458:LEU:HD12	1:A:592:MET:HE3	1.98	0.45
1:A:198:LEU:HD11	1:A:200:ILE:HD11	1.97	0.45
1:A:563:LEU:N	1:A:564:PRO:CD	2.80	0.44
1:A:161:GLU:OE1	1:A:278:LEU:HD22	2.17	0.44
1:A:532:GLU:HB3	1:A:640:ILE:HD11	2.00	0.44
1:A:458:LEU:HD12	1:A:592:MET:CE	2.48	0.43
1:A:458:LEU:HD12	1:A:592:MET:HG2	1.99	0.43
1:A:412[A]:LEU:HD23	1:A:457:LEU:HD22	2.01	0.43
1:A:37:MET:HG3	1:A:126:PHE:CD1	2.53	0.43
1:A:233:PRO:HD2	1:A:236:LEU:HD12	2.01	0.43
1:A:232:ARG:N	1:A:233:PRO:CD	2.82	0.43
1:A:419:ILE:HD11	1:A:518:HIS:HB2	2.00	0.42
1:A:716:MET:HE2	1:A:759:ILE:CD1	2.49	0.42
1:A:726[B]:MET:HE1	1:A:748:TYR:N	2.35	0.41
1:A:730:TYR:O	1:A:733:VAL:HG22	2.20	0.41
1:A:583:VAL:HG21	1:A:635:TYR:HB3	2.02	0.41
1:A:219:SER:HA	1:A:246:LEU:HD22	2.02	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:25:MET:HE2	1:A:25:MET:HB3	1.72	0.41
1:A:453:ASN:ND2	3:A:3272:HOH:O	2.49	0.41
1:A:244:ARG:CZ	1:A:246:LEU:HD13	2.51	0.41
1:A:109:GLY:HA3	1:A:269:PHE:CZ	2.56	0.41
1:A:409:PHE:HD2	1:A:507:TYR:CZ	2.40	0.40
1:A:124:LEU:HD11	1:A:271:PHE:CE1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	789/811 (97%)	770 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	708/726 (98%)	703 (99%)	5 (1%)	91	91

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	391	TRP
1	A	443[A]	GLU
1	A	443[B]	GLU
1	A	768	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	789	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	9	ASN
1	A	89	ASN
1	A	807	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	A	1001	1,2	12,14,15	0.76	1 (8%)	15,19,21	1.72	2 (13%)
2	NAG	A	1002	2	12,14,15	0.72	1 (8%)	15,19,21	0.90	0
2	NAG	A	1003	1,2	12,14,15	0.73	1 (8%)	15,19,21	1.14	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	1004	2	12,14,15	0.65	0	15,19,21	1.10	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1001	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1002	2	-	0/6/23/26	0/1/1/1
2	NAG	A	1003	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1004	2	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1002	NAG	O5-C5	-2.13	1.41	1.45
2	A	1003	NAG	O5-C5	-2.11	1.41	1.45
2	A	1001	NAG	O5-C5	-2.04	1.41	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	NAG	C3-C4-C5	4.64	118.49	110.20
2	A	1001	NAG	O5-C5-C4	3.60	115.22	110.65
2	A	1003	NAG	C3-C4-C5	3.06	115.66	110.20
2	A	1004	NAG	O5-C5-C4	2.50	113.82	110.65
2	A	1004	NAG	C3-C4-C5	2.39	114.47	110.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	788/811 (97%)	0.73	99 (12%) 4 4	26, 42, 73, 97	0

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	806	HIS	8.0
1	A	0	PHE	5.7
1	A	247	TYR	5.6
1	A	803	HIS	5.5
1	A	740	GLU	5.4
1	A	178	GLU	5.1
1	A	-1	GLU	4.6
1	A	644	ASP	4.5
1	A	232	ARG	4.4
1	A	802	SER	4.3
1	A	-2	ALA	4.3
1	A	234	ALA	4.3
1	A	246	LEU	4.3
1	A	224	LEU	4.1
1	A	145	ASP	4.1
1	A	122	VAL	3.9
1	A	422	ASP	3.9
1	A	807	HIS	3.9
1	A	474	ASP	3.9
1	A	114	PRO	3.8
1	A	180	GLU	3.8
1	A	181	THR	3.7
1	A	123	VAL	3.7
1	A	646	GLU	3.6
1	A	39	ILE	3.6
1	A	804	HIS	3.6
1	A	647	ILE	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	221	ARG	3.4
1	A	266	LYS	3.3
1	A	220	ILE	3.3
1	A	124	LEU	3.3
1	A	107	VAL	3.3
1	A	113	ASP	3.3
1	A	539	ASP	3.3
1	A	475	PHE	3.2
1	A	535	GLU	3.1
1	A	147	PRO	3.1
1	A	645	ASP	3.1
1	A	384	PHE	3.0
1	A	126	PHE	3.0
1	A	544	ASP	3.0
1	A	60	GLN	3.0
1	A	493	MET	3.0
1	A	92	VAL	3.0
1	A	23	PHE	3.0
1	A	738	ALA	2.9
1	A	47	GLN	2.9
1	A	319	GLU	2.9
1	A	7	PHE	2.9
1	A	133	ILE	2.9
1	A	257	ILE	2.9
1	A	225	GLU	2.8
1	A	105	VAL	2.8
1	A	24	GLY	2.8
1	A	273	ILE	2.7
1	A	292	GLU	2.7
1	A	223	ILE	2.7
1	A	736	ASP	2.7
1	A	685	GLU	2.6
1	A	281	THR	2.6
1	A	739	GLY	2.5
1	A	541	GLY	2.5
1	A	529	LEU	2.5
1	A	125	TYR	2.5
1	A	706	ASP	2.5
1	A	805	HIS	2.5
1	A	643	GLU	2.5
1	A	707	GLU	2.5
1	A	421	ASP	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	16	PRO	2.4
1	A	48	ASP	2.4
1	A	37	MET	2.3
1	A	41	TRP	2.3
1	A	765	GLU	2.3
1	A	25	MET	2.3
1	A	659	VAL	2.3
1	A	267	ASP	2.3
1	A	61	ASP	2.3
1	A	51	HIS	2.3
1	A	248	ASN	2.3
1	A	652	CYS	2.3
1	A	121	SER	2.3
1	A	693	LEU	2.3
1	A	177	PRO	2.2
1	A	38	GLY	2.2
1	A	249	PHE	2.2
1	A	15	ALA	2.2
1	A	222	ASP	2.2
1	A	538	GLU	2.2
1	A	531	ASP	2.2
1	A	84	GLU	2.2
1	A	146	GLY	2.1
1	A	377	THR	2.1
1	A	2	GLU	2.1
1	A	768	TYR	2.1
1	A	587	ALA	2.0
1	A	764	GLU	2.0
1	A	705	LYS	2.0
1	A	351	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	1004	14/15	0.35	6.25	72,84,92,93	0
2	NAG	A	1002	14/15	0.28	4.85	64,72,78,80	0
2	NAG	A	1003	14/15	0.21	0.94	51,64,71,75	0
2	NAG	A	1001	14/15	0.16	-0.10	43,48,57,58	0

## 6.5 Other polymers

There are no such residues in this entry.