



Full wwPDB X-ray Structure Validation Report

Mar 1, 2014 – 01:40 AM GMT

PDB ID : 4JDV
Title : Crystal structure of germ-line precursor of NIH45-46 Fab
Authors : Diskin, R.; Scharf, L.; Bjorkman, P.J.
Deposited on : 2013-02-25
Resolution : 1.65 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1404 (1.68-1.64)
Clashscore	79885	1001 (1.66-1.66)
Ramachandran outliers	78287	1581 (1.68-1.64)
Sidechain outliers	78261	1580 (1.68-1.64)
RSRZ outliers	66119	1404 (1.68-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	232	
1	H	232	
2	B	210	
2	L	210	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7597 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	227	Total	C	N	O	S	0	0	0
			1737	1095	299	333	10			
1	H	215	Total	C	N	O	S	0	0	0
			1652	1046	282	315	9			

- Molecule 2 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	208	Total	C	N	O	S	0	0	0
			1604	1007	269	324	4			
2	L	209	Total	C	N	O	S	0	0	0
			1613	1012	270	327	4			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).

Image for chem-comp SO4 is not available.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).

Image for chem-comp GOL is not available.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is TRIS-HYDROXYMETHYL-METHYL-AMMONIUM (three-letter code: 144) (formula: $C_4H_{12}NO_3$).

Image for chem-comp 144 is not available.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		
5	H	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 6 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).

Image for chem-comp 1PE is not available.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	H	1	Total	C	O	0	0
			16	10	6		
6	L	1	Total	C	O	0	0
			16	10	6		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	281	Total	O	0	0
			281	281		
7	B	238	Total	O	0	0
			238	238		
7	H	173	Total	O	0	0
			173	173		
7	L	219	Total	O	0	0
			219	219		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab heavy chain

Chain A: 



- Molecule 1: Fab heavy chain

Chain H: 



- Molecule 2: Fab light chain

Chain B: 



- Molecule 2: Fab light chain

Chain L: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.96Å 70.14Å 225.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.89 – 1.65 37.89 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.4 (37.89-1.65) 95.2 (37.89-1.65)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 1.65Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1069)	Depositor
R, R_{free}	0.174 , 0.203 0.174 , 0.203	Depositor DCC
R_{free} test set	1730 reflections (1.69%)	DCC
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.211	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 106926 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7597	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 144, SO4, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1783	0.58	1/2426 (0.0%)
1	H	0.39	0/1696	0.55	0/2310
2	B	0.44	1/1639 (0.1%)	0.62	0/2226
2	L	0.38	0/1648	0.58	0/2238
All	All	0.40	1/6766 (0.0%)	0.58	1/9200 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	80	PRO	N-CD	5.57	1.55	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	ARG	NE-CZ-NH1	5.64	123.12	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1737	0	27	4	0
1	H	1652	0	37	10	0
2	B	1604	0	62	12	0
2	L	1613	0	56	12	0
3	A	10	0	0	0	0
3	B	5	0	0	1	0
3	L	5	0	0	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	8	0	12	2	0
5	H	8	0	12	0	0
6	H	16	0	22	6	0
6	L	16	0	22	10	0
7	A	281	0	0	6	0
7	B	238	0	0	6	0
7	H	173	0	0	5	0
7	L	219	0	0	2	0
All	All	7597	0	266	47	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (47) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
6:H:302:1PE:H161	7:H:515:HOH:O	1.60	1.01
1:H:98:CYS:SG	1:H:99(B):ARG:CG	2.63	0.87
1:H:58:ASN:OD1	6:H:302:1PE:H151	1.74	0.86
1:A:220:CYS:SG	7:B:606:HOH:O	2.32	0.85
6:H:302:1PE:H132	7:L:549:HOH:O	1.78	0.83
2:L:45:ARG:NH1	6:L:302:1PE:H222	1.96	0.81
2:B:125:THR:N	7:B:619:HOH:O	2.20	0.74
2:B:24:ARG:NH1	2:B:70:ASP:OD1	2.24	0.71
2:B:54:ARG:NH1	7:B:584:HOH:O	2.25	0.69
2:L:185:HIS:ND1	7:L:592:HOH:O	2.28	0.65
1:H:82(B):ARG:NH1	7:H:558:HOH:O	2.30	0.65
5:A:304:144:H32	7:A:615:HOH:O	1.95	0.65
1:H:102:TRP:NE1	6:H:302:1PE:H162	2.12	0.64
2:L:179:LYS:NZ	2:L:183:GLU:OE2	2.32	0.63
2:B:29:VAL:CB	2:B:90:GLN:OE1	2.48	0.62
2:L:61:ARG:NH2	6:L:302:1PE:H141	2.17	0.60
6:L:302:1PE:OH4	6:L:302:1PE:OH2	2.20	0.60
2:B:2:ILE:CG2	2:B:93:PHE:CZ	2.86	0.59
7:A:642:HOH:O	6:L:302:1PE:H152	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:186:LYS:NZ	3:B:301:SO4:O3	2.37	0.57
2:L:45:ARG:NH1	6:L:302:1PE:C22	2.67	0.56
1:H:98:CYS:SG	1:H:99(B):ARG:CD	2.93	0.56
2:L:59:PRO:CG	6:L:302:1PE:H252	2.37	0.55
2:L:81:GLU:CG	6:L:302:1PE:H142	2.37	0.54
2:B:24:ARG:NH1	2:B:69:THR:OG1	2.40	0.54
2:B:24:ARG:CZ	7:B:601:HOH:O	2.57	0.53
2:B:2:ILE:CD1	2:B:29:VAL:CG2	2.87	0.53
2:L:59:PRO:CB	6:L:302:1PE:H161	2.39	0.52
2:B:24:ARG:NH2	7:B:601:HOH:O	2.44	0.51
1:A:214:ARG:NH2	7:A:673:HOH:O	2.44	0.49
2:L:138:ARG:NH2	2:L:159:VAL:HG21	2.27	0.49
5:A:304:144:C3	7:A:615:HOH:O	2.58	0.49
2:L:134:ASN:N	2:L:170:SER:HG	2.11	0.48
1:H:102:TRP:CZ2	6:H:302:1PE:H251	2.49	0.48
2:B:24:ARG:NH1	2:B:70:ASP:CG	2.67	0.47
6:H:302:1PE:H261	7:H:449:HOH:O	2.14	0.46
1:A:208:ASN:OD1	7:A:676:HOH:O	2.21	0.46
2:B:39:LYS:NZ	7:B:526:HOH:O	2.49	0.45
1:A:205:LYS:N	1:A:206:PRO:CD	2.80	0.45
7:A:612:HOH:O	6:L:302:1PE:H242	2.17	0.44
1:H:151:PRO:C	1:H:152:GLU:O	2.56	0.42
2:L:138:ARG:CZ	2:L:159:VAL:HG21	2.49	0.42
1:H:67:VAL:CG1	7:H:462:HOH:O	2.68	0.42
2:L:1:GLU:CG	2:L:2:ILE:N	2.83	0.41
1:H:205:LYS:N	1:H:206:PRO:CD	2.83	0.41
6:L:302:1PE:C13	6:L:302:1PE:C12	2.98	0.41
1:H:82(B):ARG:NH2	7:H:413:HOH:O	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/232 (96%)	222 (100%)	1 (0%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	209/232 (90%)	207 (99%)	2 (1%)	0	100	100
2	B	206/210 (98%)	201 (98%)	5 (2%)	0	100	100
2	L	207/210 (99%)	203 (98%)	4 (2%)	0	100	100
All	All	845/884 (96%)	833 (99%)	12 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/196 (98%)	191 (100%)	1 (0%)	94	87
1	H	182/196 (93%)	179 (98%)	3 (2%)	75	51
2	B	180/182 (99%)	178 (99%)	2 (1%)	84	66
2	L	181/182 (100%)	181 (100%)	0	100	100
All	All	735/756 (97%)	729 (99%)	6 (1%)	89	78

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	ARG
2	B	2	ILE
2	B	101	GLU
1	H	58	ASN
1	H	71	ARG
1	H	154	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	301	-	4,4,4	0.33	0	6,6,6	0.18	0
3	SO4	A	302	-	4,4,4	0.27	0	6,6,6	0.12	0
4	GOL	A	303	-	5,5,5	0.29	0	5,5,5	0.43	0
5	144	A	304	-	7,7,7	2.16	3 (42%)	9,9,9	1.06	0
3	SO4	B	301	-	4,4,4	0.25	0	6,6,6	0.06	0
4	GOL	B	302	-	5,5,5	0.40	0	5,5,5	0.15	0
5	144	H	301	-	7,7,7	2.04	3 (42%)	9,9,9	0.76	0
6	1PE	H	302	-	15,15,15	0.47	0	14,14,14	0.86	0
3	SO4	L	301	-	4,4,4	0.25	0	6,6,6	0.11	0
6	1PE	L	302	-	15,15,15	0.53	0	14,14,14	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	301	-	-	0/0/0/0	0/0/0/0
3	SO4	A	302	-	-	0/0/0/0	0/0/0/0
4	GOL	A	303	-	-	0/4/4/4	0/0/0/0
5	144	A	304	-	-	0/0/9/9	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	B	301	-	-	0/0/0/0	0/0/0/0
4	GOL	B	302	-	-	0/4/4/4	0/0/0/0
5	144	H	301	-	-	0/0/9/9	0/0/0/0
6	1PE	H	302	-	-	0/13/13/13	0/0/0/0
3	SO4	L	301	-	-	0/0/0/0	0/0/0/0
6	1PE	L	302	-	-	0/13/13/13	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	304	144	C3-N	-3.28	1.46	1.52
5	A	304	144	C4-N	-3.16	1.47	1.52
5	H	301	144	C4-N	-3.09	1.47	1.52
5	A	304	144	C2-N	-3.06	1.47	1.52
5	H	301	144	C3-N	-3.03	1.47	1.52
5	H	301	144	C2-N	-2.88	1.47	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	227/232 (97%)	-0.29	3 (1%) 74 76	8, 20, 42, 53	1 (0%)
1	H	215/232 (92%)	0.09	20 (9%) 9 7	15, 26, 54, 67	1 (0%)
2	B	208/210 (99%)	-0.35	1 (0%) 88 92	14, 21, 35, 52	0
2	L	209/210 (99%)	-0.39	1 (0%) 88 92	14, 20, 40, 61	0
All	All	859/884 (97%)	-0.23	25 (2%) 48 50	8, 22, 45, 67	2 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	114	THR	14.1
1	H	114	THR	13.1
2	B	29	VAL	4.5
1	A	224	HIS	3.6
1	H	212	ASP	3.5
1	H	162	ALA	3.5
1	H	131	SER	3.4
1	H	193	LEU	3.2
1	H	163	LEU	3.0
1	H	197	THR	3.0
1	H	205	LYS	2.9
1	H	191	SER	2.8
1	H	217	PRO	2.8
1	H	198	TYR	2.7
1	H	158	TRP	2.7
2	L	209	GLU	2.6
1	H	139	THR	2.5
1	A	133	LYS	2.3
1	H	211	VAL	2.2
1	H	196	GLN	2.2
1	H	189	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	150	PHE	2.1
1	H	190	SER	2.0
1	H	160	SER	2.0
1	H	199	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	A	303	6/6	0.21	-	10,11,16,18	6
3	SO4	A	301	5/5	0.20	-	10,12,13,13	5
6	1PE	L	302	16/16	0.30	-	14,28,34,38	16
3	SO4	L	301	5/5	0.29	-	36,38,41,42	5
5	144	A	304	8/8	0.28	-	13,21,23,30	8
4	GOL	B	302	6/6	0.10	-	24,26,29,30	0
3	SO4	A	302	5/5	0.20	-	10,14,16,17	5
6	1PE	H	302	16/16	0.40	-	22,32,36,39	16
5	144	H	301	8/8	0.22	-	23,29,34,36	8
3	SO4	B	301	5/5	0.23	-	37,39,40,44	5

6.5 Other polymers ⓘ

There are no such residues in this entry.