



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 04:50 AM GMT

PDB ID : 2JGP
Title : Structure of the TycC5-6 PCP-C bidomain of the tyrocidine synthetase TycC
Authors : Samel, S.A.; Schoenafinger, G.; Knappe, T.A.; Marahiel, M.A.; Essen, L.-O.
Deposited on : 2007-02-13
Resolution : 1.85 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1269 (1.86-1.86)
Clashscore	79885	1470 (1.86-1.86)
Ramachandran outliers	78287	1451 (1.86-1.86)
Sidechain outliers	78261	1451 (1.86-1.86)
RSRZ outliers	66119	1269 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	520	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	SO4	A	1523	-	X
3	DIO	A	1526	-	X
4	NA	A	1527	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4477 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TYROCIDINE SYNTHETASE 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	520	4161	2655	695	795	16	35	2	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	183	GLU	GLN	CONFLICT	UNP O30409

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is 1,4-DIETHYLENE DIOXIDE (three-letter code: DIO) (formula: C₄H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	4	2		
3	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	293	Total	O	0	0
			293	293		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TYROCIDINE SYNTHETASE 3

Chain A:



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.97Å 84.97Å 164.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.81 – 1.85 19.82 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.6 (19.81-1.85) 98.6 (19.82-1.85)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.208 , 0.238 0.215 , 0.244	Depositor DCC
R_{free} test set	1766 reflections (3.55%)	DCC
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 59.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 51770 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4477	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, DIO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	1/4259 (0.0%)	0.74	6/5783 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	460	GLY	C-N	-7.94	1.15	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	457	ASP	CB-CG-OD2	-28.52	92.63	118.30
1	A	455	MET	O-C-N	-22.29	87.04	122.70
1	A	457	ASP	CB-CG-OD1	15.23	132.00	118.30
1	A	460	GLY	C-N-CA	6.67	138.37	121.70
1	A	455	MET	CA-C-N	5.61	129.54	117.20
1	A	460	GLY	O-C-N	-5.40	114.06	122.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	455	MET	Mainchain
1	A	457	ASP	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4161	0	4084	41	0
2	A	10	0	0	0	0
3	A	12	0	16	1	0
4	A	1	0	0	0	0
5	A	293	0	0	1	0
All	All	4477	0	4100	41	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (41) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:508:ILE:HD12	1:A:509:VAL:N	2.01	0.76
1:A:117:VAL:HG12	1:A:120:SER:HB3	1.72	0.71
1:A:117:VAL:CG1	1:A:120:SER:HB3	2.21	0.70
1:A:323:LEU:CD2	1:A:508:ILE:HD13	2.23	0.69
1:A:117:VAL:HG12	1:A:117:VAL:O	1.95	0.66
1:A:220:THR:HG21	3:A:1525:DIO:H22	1.82	0.61
1:A:252:ARG:NH1	1:A:253:ILE:HD11	2.15	0.61
1:A:420:ILE:HG23	1:A:422:ARG:HB2	1.83	0.60
1:A:175:LEU:C	1:A:175:LEU:HD23	2.24	0.57
1:A:521:MET:O	1:A:522:LEU:HB2	2.05	0.57
1:A:446:LEU:HD13	1:A:451:LEU:HD11	1.85	0.57
1:A:508:ILE:HD12	1:A:508:ILE:C	2.26	0.56
1:A:3:ARG:NH2	1:A:28:PRO:O	2.38	0.55
1:A:3:ARG:NH2	1:A:27:VAL:HG12	2.22	0.53
1:A:126:THR:HG21	1:A:189:VAL:HG11	1.90	0.53
1:A:175:LEU:HD23	1:A:176:ALA:N	2.25	0.52
1:A:19:GLN:NE2	5:A:2020:HOH:O	2.43	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:342:LEU:HD13	1:A:497:PHE:CZ	2.47	0.49
1:A:3:ARG:NH2	1:A:27:VAL:CG1	2.77	0.48
1:A:175:LEU:HD23	1:A:176:ALA:C	2.34	0.48
1:A:126:THR:HG21	1:A:189:VAL:CG1	2.44	0.47
1:A:321:HIS:CE1	1:A:331:LEU:HD22	2.49	0.47
1:A:13:TRP:CH2	1:A:59:VAL:HG21	2.50	0.47
1:A:50:LEU:HD22	1:A:61:ILE:O	2.14	0.47
1:A:381:ASN:ND2	1:A:399:ASN:HD22	2.14	0.45
1:A:117:VAL:O	1:A:117:VAL:CG1	2.63	0.45
1:A:446:LEU:CD1	1:A:451:LEU:HD11	2.46	0.45
1:A:50:LEU:HD11	1:A:63:LEU:HA	2.00	0.44
1:A:301:GLN:NE2	1:A:464:LYS:O	2.50	0.44
1:A:34:ASP:HB3	1:A:39:LEU:HD11	2.00	0.44
1:A:367:GLU:N	1:A:368:PRO:CD	2.82	0.43
1:A:323:LEU:HD22	1:A:508:ILE:HD13	2.00	0.42
1:A:181:THR:HG22	1:A:182:GLU:N	2.35	0.41
1:A:213:GLU:N	1:A:213:GLU:CD	2.73	0.41
1:A:175:LEU:C	1:A:175:LEU:CD2	2.88	0.41
1:A:46:ALA:HB1	1:A:66:LEU:HD23	2.02	0.41
1:A:440:THR:HG22	1:A:440:THR:O	2.20	0.41
1:A:181:THR:HG22	1:A:183:GLU:H	1.85	0.40
1:A:357:PRO:HD3	1:A:433:VAL:HG13	2.02	0.40
1:A:93:VAL:HG11	1:A:260:VAL:HB	2.04	0.40
1:A:235:LEU:CD1	1:A:370:VAL:HG21	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	520/520 (100%)	489 (94%)	25 (5%)	6 (1%)	19 5

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	213	GLU
1	A	440	THR
1	A	441	GLU
1	A	133	LEU
1	A	475	ARG
1	A	423	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	452/450 (100%)	439 (97%)	13 (3%)	55 36

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	ARG
1	A	94	GLN
1	A	133	LEU
1	A	169	GLU
1	A	187	GLU
1	A	198	LEU
1	A	235	LEU
1	A	325	GLN
1	A	341	VAL
1	A	342	LEU
1	A	426	ARG
1	A	437	GLN
1	A	457	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	143	GLN
1	A	267	GLN
1	A	280	GLN
1	A	325	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	381	ASN
1	A	427	ASN
1	A	500	HIS
1	A	510	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1523	-	4,4,4	0.29	0	6,6,6	0.15	0
2	SO4	A	1524	-	4,4,4	0.30	0	6,6,6	0.15	0
3	DIO	A	1525	-	6,6,6	0.48	0	6,6,6	0.46	0
3	DIO	A	1526	-	6,6,6	0.48	0	6,6,6	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1523	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1524	-	-	0/0/0/0	0/0/0/0
3	DIO	A	1525	-	-	0/0/6/6	0/1/1/1
3	DIO	A	1526	-	-	0/0/6/6	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	515/520 (99%)	1.16	94 (18%) 2 2	43, 62, 75, 87	0

All (94) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	443	VAL	18.2
1	A	442	GLU	8.5
1	A	421	ALA	8.2
1	A	420	ILE	7.8
1	A	439	SER	7.4
1	A	422	ARG	7.2
1	A	440	THR	6.8
1	A	114	PHE	6.4
1	A	475	ARG	6.1
1	A	226	ILE	5.7
1	A	117	VAL	5.5
1	A	285	ASP	5.1
1	A	444	MET	5.1
1	A	447	PRO	5.0
1	A	419	ALA	4.5
1	A	476	GLU	4.5
1	A	441	GLU	4.4
1	A	461	GLN	4.3
1	A	446	LEU	4.2
1	A	255	TYR	4.1
1	A	94	GLN	4.1
1	A	462	HIS	4.1
1	A	445	THR	4.0
1	A	225	SER	4.0
1	A	424	LEU	4.0
1	A	132	LYS	4.0
1	A	371	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	423	ASP	3.9
1	A	369	ILE	3.8
1	A	13	TRP	3.8
1	A	72	ILE	3.8
1	A	520	ASP	3.8
1	A	247	VAL	3.7
1	A	18	ALA	3.7
1	A	21	TRP	3.6
1	A	370	VAL	3.6
1	A	118	GLY	3.6
1	A	17	LEU	3.6
1	A	129	ILE	3.5
1	A	213	GLU	3.4
1	A	115	GLU	3.3
1	A	438	ASN	3.2
1	A	119	ILE	3.2
1	A	385	ARG	3.1
1	A	258	PHE	3.1
1	A	79	ILE	3.0
1	A	32	ALA	3.0
1	A	44	LEU	3.0
1	A	75	LEU	3.0
1	A	455	MET	3.0
1	A	284	ASP	2.9
1	A	64	ARG	2.9
1	A	448	GLU	2.9
1	A	267	GLN	2.9
1	A	217	VAL	2.9
1	A	245	GLY	2.9
1	A	373	PHE	2.8
1	A	386	GLU	2.8
1	A	254[A]	GLN	2.7
1	A	467	LEU	2.7
1	A	30	VAL	2.7
1	A	15	ALA	2.6
1	A	116	GLY	2.6
1	A	212	GLY	2.6
1	A	187	GLU	2.5
1	A	522	LEU	2.5
1	A	57	TYR	2.5
1	A	250	GLU	2.5
1	A	66	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	203	LEU	2.5
1	A	489	PHE	2.4
1	A	208	LEU	2.4
1	A	84[A]	LYS	2.4
1	A	151	SER	2.4
1	A	152	LEU	2.3
1	A	214	ASP	2.3
1	A	188	LEU	2.2
1	A	183	GLU	2.2
1	A	366	LEU	2.2
1	A	105	GLN	2.2
1	A	8	ALA	2.2
1	A	186	ARG	2.1
1	A	9	PRO	2.1
1	A	496	ARG	2.1
1	A	48	ARG	2.1
1	A	372	MET	2.1
1	A	353	ILE	2.1
1	A	7	VAL	2.1
1	A	218	LEU	2.1
1	A	223	HIS	2.0
1	A	137	ARG	2.0
1	A	393	LEU	2.0
1	A	474	GLU	2.0
1	A	3	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NA	A	1527	1/1	0.55	9.86	69,69,69,69	0
3	DIO	A	1526	6/6	0.29	4.51	79,79,79,79	0
2	SO4	A	1523	5/5	0.23	2.82	69,70,70,70	0
2	SO4	A	1524	5/5	0.18	0.37	57,57,58,59	0
3	DIO	A	1525	6/6	0.14	-0.54	55,56,56,56	0

6.5 Other polymers ⓘ

There are no such residues in this entry.