



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 05:31 AM GMT

PDB ID : 4JK3
Title : PylD holoenzyme (SeMet)
Authors : Quitterer, F.; Beck, P.; Bacher, A.; Groll, M.
Deposited on : 2013-03-09
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	259	
1	B	259	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 3939 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PylD.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	Se	0	0	0
			1899	1210	305	374	5	5			
1	B	251	Total	C	N	O	S	Se	0	0	0
			1899	1210	305	374	5	5			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	30	Total	O	0	0
			30	30		
5	B	19	Total	O	0	0
			19	19		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PylD

Chain A: 



• Molecule 1: PylD

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	79.65Å 155.40Å 39.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.50 14.82 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (15.00-2.50) 100.0 (14.82-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.17 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.216 , 0.257 0.214 , 0.258	Depositor DCC
R_{free} test set	882 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	45.0	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 45.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 17685 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3939	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/1929	0.62	0/2611
1	B	0.37	0/1929	0.59	0/2611
All	All	0.39	0/3858	0.60	0/5222

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1899	0	0	3	0
1	B	1899	0	0	3	0
2	A	44	0	0	0	0
2	B	44	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	30	0	0	1	0
5	B	19	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3939	0	0	5	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (5) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:220:VAL:CB	5:A:1028:HOH:O	2.46	0.64
1:B:5:THR:OG1	1:B:6:PRO:CD	2.56	0.54
1:B:27:THR:O	5:B:1016:HOH:O	2.19	0.52
1:A:5:THR:OG1	1:A:6:PRO:CD	2.59	0.51
1:A:206:CYS:SG	1:B:206:CYS:SG	3.14	0.45

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/259 (95%)	238 (96%)	8 (3%)	1 (0%)	43	66
1	B	247/259 (95%)	239 (97%)	7 (3%)	1 (0%)	43	66
All	All	494/518 (95%)	477 (97%)	15 (3%)	2 (0%)	43	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	PRO
1	B	6	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of

similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/208 (100%)	193 (93%)	15 (7%)	21	36
1	B	208/208 (100%)	194 (93%)	14 (7%)	23	40
All	All	416/416 (100%)	387 (93%)	29 (7%)	21	38

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	11	ASN
1	A	38	LEU
1	A	41	THR
1	A	44	SER
1	A	48	VAL
1	A	51	VAL
1	A	53	VAL
1	A	67	LEU
1	A	112	ASN
1	A	137	LEU
1	A	145	LEU
1	A	184	LEU
1	A	214	VAL
1	A	242	LEU
1	B	5	THR
1	B	38	LEU
1	B	41	THR
1	B	48	VAL
1	B	51	VAL
1	B	53	VAL
1	B	67	LEU
1	B	112	ASN
1	B	137	LEU
1	B	140	ASP
1	B	145	LEU
1	B	184	LEU
1	B	214	VAL
1	B	242	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	A	901	4	48,48,48	0.95	3 (6%)	73,73,73	1.61	10 (13%)
2	NAD	B	901	4	48,48,48	0.97	3 (6%)	73,73,73	1.65	12 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	901	4	-	0/30/62/62	0/3/5/5
2	NAD	B	901	4	-	0/30/62/62	0/3/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	NAD	C5A-C4A	3.20	1.47	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	NAD	C4A-N9A	-3.17	1.33	1.37
2	A	901	NAD	C4A-N9A	-3.04	1.33	1.37
2	A	901	NAD	C5A-C4A	2.76	1.46	1.40
2	A	901	NAD	C2A-N3A	2.15	1.36	1.32
2	B	901	NAD	O4D-C1D	2.14	1.44	1.41

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	901	NAD	N3A-C2A-N1A	-6.28	123.45	128.71
2	A	901	NAD	N3A-C4A-N9A	5.53	135.41	125.43
2	A	901	NAD	N3A-C2A-N1A	-5.52	124.09	128.71
2	B	901	NAD	C4A-C5A-N7A	-4.50	105.67	109.52
2	B	901	NAD	N3A-C4A-N9A	4.18	132.98	125.43
2	B	901	NAD	O4D-C1D-N1N	4.14	112.19	107.95
2	A	901	NAD	C4B-O4B-C1B	-3.86	105.56	109.75
2	A	901	NAD	C8A-N9A-C4A	3.53	109.60	106.90
2	A	901	NAD	C4A-C5A-N7A	-3.52	106.51	109.52
2	B	901	NAD	C1B-N9A-C4A	-3.49	120.61	126.64
2	A	901	NAD	O4B-C1B-N9A	3.27	111.48	108.44
2	A	901	NAD	C5A-C4A-N3A	-3.27	118.58	125.70
2	A	901	NAD	PN-O3-PA	-2.72	121.25	132.95
2	B	901	NAD	C5A-C4A-N3A	-2.65	119.93	125.70
2	B	901	NAD	PN-O3-PA	-2.54	122.03	132.95
2	B	901	NAD	C8A-N9A-C4A	2.44	108.76	106.90
2	A	901	NAD	C2A-N3A-C4A	2.31	120.58	114.01
2	B	901	NAD	C2A-N1A-C6A	2.19	122.72	118.77
2	B	901	NAD	C8A-N9A-C1B	2.15	130.62	126.38
2	B	901	NAD	O2A-PA-O1A	2.14	124.15	112.21
2	B	901	NAD	C2A-N3A-C4A	2.14	120.09	114.01
2	A	901	NAD	O2A-PA-O1A	2.02	123.48	112.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/259 (96%)	0.39	20 (7%) 12 11	25, 42, 74, 132	0
1	B	251/259 (96%)	0.95	45 (17%) 2 2	31, 60, 104, 137	0
All	All	502/518 (96%)	0.67	65 (12%) 4 4	25, 49, 92, 137	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	60	ILE	14.1
1	B	60	ILE	11.8
1	A	61	GLY	8.7
1	B	42	PHE	6.9
1	A	7	ASP	6.2
1	B	61	GLY	6.1
1	A	5	THR	5.5
1	B	226	ILE	5.1
1	B	140	ASP	5.1
1	B	218	ASN	4.6
1	B	232	GLU	4.4
1	B	45	SER	4.3
1	B	105	ASP	4.0
1	A	226	ILE	4.0
1	B	75	GLY	4.0
1	B	21	SER	3.9
1	B	9	LEU	3.9
1	B	6	PRO	3.7
1	B	17	GLN	3.5
1	B	55	SER	3.5
1	A	11	ASN	3.4
1	A	10	ILE	3.4
1	B	73	TYR	3.3
1	B	234	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	164	ASP	3.2
1	B	5	THR	3.2
1	B	72	GLN	3.2
1	A	55	SER	3.2
1	B	174	GLU	3.1
1	B	43	SER	3.1
1	B	11	ASN	3.0
1	B	30	ASP	3.0
1	B	14	MSE	2.9
1	B	24	GLN	2.9
1	B	10	ILE	2.9
1	B	94	GLN	2.8
1	B	28	GLY	2.8
1	B	231	SER	2.8
1	B	236	ASP	2.7
1	B	114	LYS	2.7
1	A	140	ASP	2.7
1	A	105	ASP	2.7
1	B	44	SER	2.7
1	B	235	ARG	2.6
1	A	94	GLN	2.6
1	B	7	ASP	2.6
1	B	162	GLN	2.5
1	B	142	LYS	2.5
1	B	41	THR	2.5
1	B	227	PRO	2.4
1	B	217	GLU	2.4
1	A	106	ARG	2.4
1	B	54	THR	2.4
1	A	227	PRO	2.4
1	B	157	ALA	2.4
1	A	62	ASN	2.4
1	A	164	ASP	2.4
1	B	230	ILE	2.3
1	A	232	GLU	2.2
1	A	8	ASP	2.2
1	B	13	ASN	2.2
1	A	6	PRO	2.2
1	B	19	ALA	2.1
1	A	54	THR	2.1
1	A	178	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NA	A	902	1/1	0.16	-0.17	43,43,43,43	0
4	MG	A	903	1/1	0.11	-0.50	49,49,49,49	0
2	NAD	B	901	44/44	0.13	-0.60	38,42,44,45	0
2	NAD	A	901	44/44	0.10	-0.73	34,36,39,42	0
4	MG	B	903	1/1	0.14	-0.75	60,60,60,60	0
3	NA	B	902	1/1	0.13	-0.76	46,46,46,46	0

6.5 Other polymers ⓘ

There are no such residues in this entry.