



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 10:27 PM GMT

PDB ID : 4JMQ  
Title : Crystal structure of pb9: The Dit of bacteriophage T5.  
Authors : Flayhan, A.; Vellieux, F.M.D.; Girard, E.; Maury, O.; Boulanger, P.; Breyton, C.  
Deposited on : 2013-03-14  
Resolution : 1.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	217	
1	B	217	
1	C	217	
1	D	217	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6935 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriophage T5 distal tail protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	190	Total	C	N	O	S	0	1	0
			1505	964	246	293	2			
1	B	189	Total	C	N	O	S	0	1	0
			1493	956	242	293	2			
1	C	201	Total	C	N	O	S	0	1	0
			1608	1029	269	307	3			
1	D	189	Total	C	N	O	S	0	2	0
			1500	961	243	294	2			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	GLU	-	EXPRESSION TAG	UNP Q6QGE8
A	206	ASN	-	EXPRESSION TAG	UNP Q6QGE8
A	207	LEU	-	EXPRESSION TAG	UNP Q6QGE8
A	208	TYR	-	EXPRESSION TAG	UNP Q6QGE8
A	209	PHE	-	EXPRESSION TAG	UNP Q6QGE8
A	210	GLN	-	EXPRESSION TAG	UNP Q6QGE8
A	211	GLY	-	EXPRESSION TAG	UNP Q6QGE8
A	212	HIS	-	EXPRESSION TAG	UNP Q6QGE8
A	213	HIS	-	EXPRESSION TAG	UNP Q6QGE8
A	214	HIS	-	EXPRESSION TAG	UNP Q6QGE8
A	215	HIS	-	EXPRESSION TAG	UNP Q6QGE8
A	216	HIS	-	EXPRESSION TAG	UNP Q6QGE8
A	217	HIS	-	EXPRESSION TAG	UNP Q6QGE8
B	205	GLU	-	EXPRESSION TAG	UNP Q6QGE8
B	206	ASN	-	EXPRESSION TAG	UNP Q6QGE8
B	207	LEU	-	EXPRESSION TAG	UNP Q6QGE8
B	208	TYR	-	EXPRESSION TAG	UNP Q6QGE8
B	209	PHE	-	EXPRESSION TAG	UNP Q6QGE8
B	210	GLN	-	EXPRESSION TAG	UNP Q6QGE8
B	211	GLY	-	EXPRESSION TAG	UNP Q6QGE8
B	212	HIS	-	EXPRESSION TAG	UNP Q6QGE8

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	213	HIS	-	EXPRESSION TAG	UNP Q6QGE8
B	214	HIS	-	EXPRESSION TAG	UNP Q6QGE8
B	215	HIS	-	EXPRESSION TAG	UNP Q6QGE8
B	216	HIS	-	EXPRESSION TAG	UNP Q6QGE8
B	217	HIS	-	EXPRESSION TAG	UNP Q6QGE8
C	205	GLU	-	EXPRESSION TAG	UNP Q6QGE8
C	206	ASN	-	EXPRESSION TAG	UNP Q6QGE8
C	207	LEU	-	EXPRESSION TAG	UNP Q6QGE8
C	208	TYR	-	EXPRESSION TAG	UNP Q6QGE8
C	209	PHE	-	EXPRESSION TAG	UNP Q6QGE8
C	210	GLN	-	EXPRESSION TAG	UNP Q6QGE8
C	211	GLY	-	EXPRESSION TAG	UNP Q6QGE8
C	212	HIS	-	EXPRESSION TAG	UNP Q6QGE8
C	213	HIS	-	EXPRESSION TAG	UNP Q6QGE8
C	214	HIS	-	EXPRESSION TAG	UNP Q6QGE8
C	215	HIS	-	EXPRESSION TAG	UNP Q6QGE8
C	216	HIS	-	EXPRESSION TAG	UNP Q6QGE8
C	217	HIS	-	EXPRESSION TAG	UNP Q6QGE8
D	205	GLU	-	EXPRESSION TAG	UNP Q6QGE8
D	206	ASN	-	EXPRESSION TAG	UNP Q6QGE8
D	207	LEU	-	EXPRESSION TAG	UNP Q6QGE8
D	208	TYR	-	EXPRESSION TAG	UNP Q6QGE8
D	209	PHE	-	EXPRESSION TAG	UNP Q6QGE8
D	210	GLN	-	EXPRESSION TAG	UNP Q6QGE8
D	211	GLY	-	EXPRESSION TAG	UNP Q6QGE8
D	212	HIS	-	EXPRESSION TAG	UNP Q6QGE8
D	213	HIS	-	EXPRESSION TAG	UNP Q6QGE8
D	214	HIS	-	EXPRESSION TAG	UNP Q6QGE8
D	215	HIS	-	EXPRESSION TAG	UNP Q6QGE8
D	216	HIS	-	EXPRESSION TAG	UNP Q6QGE8
D	217	HIS	-	EXPRESSION TAG	UNP Q6QGE8

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	215	Total O 215 215	0	0
2	B	212	Total O 212 212	0	0
2	C	191	Total O 191 191	0	0
2	D	211	Total O 211 211	0	0

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.66Å 70.03Å 71.01Å 91.05° 107.66° 112.03°	Depositor
Resolution (Å)	27.69 – 1.90 66.91 – 1.89	Depositor EDS
% Data completeness (in resolution range)	95.4 (27.69-1.90) 95.4 (66.91-1.89)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	0.19	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1069)	Depositor
R, $R_{free}$	0.203 , 0.252 0.205 , 0.255	Depositor DCC
$R_{free}$ test set	3547 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.8	Xtriage
Anisotropy	0.291	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 70973 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6935	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/1545	0.55	0/2100
1	B	0.38	0/1532	0.56	0/2082
1	C	0.39	0/1656	0.60	0/2251
1	D	0.35	0/1542	0.55	0/2096
All	All	0.37	0/6275	0.57	0/8529

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1505	0	0	12	0
1	B	1493	0	0	13	0
1	C	1608	0	16	11	0
1	D	1500	0	5	9	0
2	A	215	0	0	8	0
2	B	212	0	0	6	0
2	C	191	0	0	3	0
2	D	211	0	0	7	0
All	All	6935	0	21	44	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:108:GLN:NE2	2:A:366:HOH:O	1.94	0.98
1:B:43:ILE:N	2:B:505:HOH:O	1.97	0.97
1:A:43:ILE:N	2:A:464:HOH:O	1.97	0.97
1:C:60:ASP:OD1	2:C:460:HOH:O	1.91	0.86
1:A:170:GLY:O	2:A:495:HOH:O	2.00	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	187/217 (86%)	184 (98%)	3 (2%)	0	100	100
1	B	186/217 (86%)	184 (99%)	2 (1%)	0	100	100
1	C	198/217 (91%)	190 (96%)	7 (4%)	1 (0%)	38	23
1	D	187/217 (86%)	183 (98%)	4 (2%)	0	100	100
All	All	758/868 (87%)	741 (98%)	16 (2%)	1 (0%)	59	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	27	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/193 (88%)	168 (99%)	1 (1%)	92	92
1	B	168/193 (87%)	165 (98%)	3 (2%)	71	66
1	C	179/193 (93%)	175 (98%)	4 (2%)	64	57
1	D	169/193 (88%)	168 (99%)	1 (1%)	92	92
All	All	685/772 (89%)	676 (99%)	9 (1%)	80	77

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	56	GLU
1	D	204	LEU
1	C	72	TYR
1	B	204	LEU
1	C	60	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	190/217 (87%)	-0.12	2 (1%) 77 79	21, 32, 60, 97	0
1	B	189/217 (87%)	-0.13	3 (1%) 68 70	19, 31, 57, 83	0
1	C	201/217 (92%)	0.10	6 (2%) 48 49	19, 33, 79, 105	0
1	D	189/217 (87%)	0.18	11 (5%) 22 22	19, 37, 85, 104	0
All	All	769/868 (88%)	0.01	22 (2%) 49 50	19, 32, 72, 105	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	209	PHE	6.0
1	C	208	TYR	5.1
1	C	186	THR	4.7
1	D	186	THR	4.4
1	D	43	ILE	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.