



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 05:39 PM GMT

PDB ID : 1JRH  
Title : COMPLEX (ANTIBODY/ANTIGEN)  
Authors : Winkler, F.K.; Sogabe, S.  
Deposited on : 1997-09-23  
Resolution : 2.80 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2295 (2.80-2.80)
Ramachandran outliers	78287	2252 (2.80-2.80)
Sidechain outliers	78261	2254 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	213	
2	H	219	
3	I	108	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3465 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANTIBODY A6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	167	Total	C	N	O	S	0	0	0
			1290	809	211	264	6			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	9	SER	ALA	CONFLICT	PIR S01320
L	11	PHE	LEU	CONFLICT	PIR S01320
L	15	LEU	VAL	CONFLICT	PIR S01320
L	17	ASP	GLU	CONFLICT	PIR S01320
L	18	ARG	SER	CONFLICT	PIR S01320
L	24	LYS	ARG	CONFLICT	PIR S01320
L	28	ASP	ASN	CONFLICT	PIR S01320
L	31	ASN	SER	CONFLICT	PIR S01320
L	32	ARG	ASN	CONFLICT	PIR S01320
L	40	PRO	GLN	CONFLICT	PIR S01320
L	42	ASN	LYS	CONFLICT	PIR S01320
L	43	ALA	SER	CONFLICT	PIR S01320
L	45	ARG	GLN	CONFLICT	PIR S01320
L	48	ILE	VAL	CONFLICT	PIR S01320
L	49	SER	TYR	CONFLICT	PIR S01320
L	50	GLY	VAL	CONFLICT	PIR S01320
L	53	SER	LYS	CONFLICT	PIR S01320
L	55	GLU	VAL	CONFLICT	PIR S01320
L	56	THR	ASP	CONFLICT	PIR S01320
L	57	GLU	GLY	CONFLICT	PIR S01320
L	69	LYS	THR	CONFLICT	PIR S01320
L	70	ASP	GLN	CONFLICT	PIR S01320
L	72	THR	SER	CONFLICT	PIR S01320
L	74	SER	LYS	CONFLICT	PIR S01320
L	76	THR	ASN	CONFLICT	PIR S01320
L	80	THR	SER	CONFLICT	PIR S01320
L	83	VAL	PHE	CONFLICT	PIR S01320

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
L	84	ALA	GLY	CONFLICT	PIR S01320
L	85	THR	SER	CONFLICT	PIR S01320
L	90	GLN	HIS	CONFLICT	PIR S01320
L	91	TYR	PHE	CONFLICT	PIR S01320
L	93	SER	ASP	CONFLICT	PIR S01320
L	?	-	PRO	DELETION	PIR S01320
L	96	TRP	PHE	CONFLICT	PIR S01320
L	100	GLY	SER	CONFLICT	PIR S01320
L	106	ILE	MET	CONFLICT	PIR S01320

- Molecule 2 is a protein called ANTIBODY A6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	180	Total	C	N	O	S	0	0	0
			1384	881	230	267	6			

- Molecule 3 is a protein called INTERFERON-GAMMA RECEPTOR ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	95	Total	C	N	O	S	11	0	0
			761	485	126	146	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	105	SER	CYS	ENGINEERED	UNP P15260

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	16	Total	O	0	0
			16	16		
4	I	5	Total	O	0	0
			5	5		
4	L	9	Total	O	0	0
			9	9		

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: ANTIBODY A6



#### • Molecule 2: ANTIBODY A6



#### • Molecule 3: INTERFERON-GAMMA RECEPTOR ALPHA CHAIN



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.74Å 90.80Å 101.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80	Depositor
% Data completeness (in resolution range)	99.3 (20.00-2.80)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.246 , 0.314	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3465	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.55	0/1317	0.82	0/1787
2	H	0.53	0/1419	0.82	2/1932 (0.1%)
3	I	0.62	1/783 (0.1%)	0.80	1/1071 (0.1%)
All	All	0.56	1/3519 (0.0%)	0.81	3/4790 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	11	SER	CB-OG	-5.32	1.35	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	23	TYR	N-CA-C	-5.60	95.88	111.00
2	H	63	LEU	CA-CB-CG	5.14	127.12	115.30
2	H	56	ASP	N-CA-C	-5.07	97.32	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1290	0	1239	43	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1384	0	1342	41	0
3	I	761	0	724	21	0
4	H	16	0	0	2	0
4	I	5	0	0	0	0
4	L	9	0	0	2	0
All	All	3465	0	3305	99	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 15.

The worst 5 of 99 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:H:150:GLU:HG3	2:H:151:PRO:HA	1.47	0.93
1:L:135:PHE:CE2	2:H:190:SER:HB3	2.11	0.85
2:H:30:THR:HG23	2:H:73:THR:HG21	1.60	0.83
1:L:162:SER:HB2	2:H:175:PRO:HG2	1.65	0.77
1:L:141:PRO:HG3	1:L:199:LYS:HD3	1.69	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	159/213 (75%)	142 (89%)	15 (9%)	2 (1%)	18	51
2	H	170/219 (78%)	146 (86%)	21 (12%)	3 (2%)	13	39
3	I	93/108 (86%)	81 (87%)	10 (11%)	2 (2%)	10	32
All	All	422/540 (78%)	369 (87%)	46 (11%)	7 (2%)	14	42

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	171	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	H	116	THR
2	H	179	GLN
2	H	216	SER
3	I	62	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	148/190 (78%)	138 (93%)	10 (7%)	22	54
2	H	155/188 (82%)	145 (94%)	10 (6%)	24	57
3	I	87/96 (91%)	81 (93%)	6 (7%)	22	53
All	All	390/474 (82%)	364 (93%)	26 (7%)	23	55

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	31	THR
2	H	81	LYS
3	I	62	ASN
2	H	38	ARG
2	H	48	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	161	ASN
1	L	198	HIS
3	I	66	HIS
3	I	69	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.