



# Full wwPDB X-ray Structure Validation Report ⓘ

May 1, 2014 – 11:38 PM EDT

PDB ID : 4JRY  
Title : Crystal Structure of SB47 TCR-HLA B\*3505-LPEP complex  
Authors : Liu, Y.C.; Rossjohn, J.; Gras, S.  
Deposited on : 2013-03-22  
Resolution : 2.80 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable22978  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22978

# 1 Overall quality at a glance

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1799 (2.80-2.80)
Clashscore	79885	2295 (2.80-2.80)
Ramachandran outliers	78287	2252 (2.80-2.80)
Sidechain outliers	78261	2254 (2.80-2.80)
RSRZ outliers	66119	1802 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	276	
2	B	100	
3	C	13	
4	D	201	
5	E	242	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
7	MG	B	103	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6830 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	0	0
			2257	1405	414	431	7			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	EXPRESSION TAG	UNP P61769

- Molecule 3 is a protein called Trans-activator protein BZLF1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	0	0	0
			101	66	15	20			

- Molecule 4 is a protein called SB47 TCR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	201	Total	C	N	O	S	0	0	0
			1574	991	261	313	9			

- Molecule 5 is a protein called SB47 TCR beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	242	Total	C	N	O	S	1	0	0
			1934	1216	340	373	5			

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	Na 1	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	2	Total 2	Mg 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	20	Total 20	O 20	0	0
8	B	14	Total 14	O 14	0	0
8	C	2	Total 2	O 2	0	0
8	D	32	Total 32	O 32	0	0
8	E	56	Total 56	O 56	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MHC class I antigen




- Molecule 2: Beta-2-microglobulin



- Molecule 3: Trans-activator protein BZLF1




- Molecule 4: SB47 TCR alpha chain



- Molecule 5: SB47 TCR beta chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	237.60Å 237.60Å 61.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 2.80 168.01 – 2.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (100.00-2.80) 99.9 (168.01-2.80)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.11 (at 2.82Å)	Xtriage
Refinement program	BUSTER	Depositor
R, $R_{free}$	0.200 , 0.233 0.203 , 0.236	Depositor DCC
$R_{free}$ test set	2147 reflections (5.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	48.8	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 27.3	EDS
Estimated twinning fraction	0.022 for -k,-h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 42581 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/2320	0.76	1/3153 (0.0%)
2	B	0.53	0/860	0.72	0/1162
3	C	0.48	0/104	0.69	0/142
4	D	0.67	0/1607	0.78	1/2178 (0.0%)
5	E	0.63	0/1989	0.76	0/2710
All	All	0.62	0/6880	0.76	2/9345 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	ASP	N-CA-C	-6.23	94.18	111.00
4	D	138	LEU	CA-CB-CG	5.06	126.94	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2257	0	2123	40	0
2	B	837	0	803	12	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	101	0	102	2	0
4	D	1574	0	1522	37	0
5	E	1934	0	1817	23	0
6	B	1	0	0	0	0
7	B	2	0	0	0	0
8	A	20	0	0	0	0
8	B	14	0	0	0	0
8	C	2	0	0	0	0
8	D	32	0	0	1	0
8	E	56	0	0	0	0
All	All	6830	0	6367	109	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (109) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
4:D:146:LYS:N	4:D:147:SER:HA	1.51	1.18
4:D:146:LYS:H	4:D:147:SER:CA	1.59	1.16
1:A:273:ARG:HG3	1:A:273:ARG:HH11	1.17	1.06
5:E:9:THR:HG23	5:E:10:HIS:HD2	1.23	0.97
4:D:13:MET:HG3	4:D:124:ILE:HG13	1.52	0.91
5:E:9:THR:HG23	5:E:10:HIS:CD2	2.07	0.90
4:D:146:LYS:N	4:D:147:SER:CA	2.20	0.86
4:D:146:LYS:H	4:D:147:SER:HA	0.74	0.85
4:D:204:ASN:O	4:D:205:ASN:HB3	1.80	0.82
4:D:209:PRO:O	4:D:210:GLU:CG	2.30	0.80
4:D:209:PRO:O	4:D:210:GLU:CB	2.29	0.77
1:A:273:ARG:HG3	1:A:273:ARG:NH1	1.90	0.76
1:A:17:ARG:HH11	1:A:17:ARG:HG2	1.54	0.72
4:D:209:PRO:O	4:D:210:GLU:HG3	1.90	0.69
4:D:208:ILE:HG13	4:D:212:THR:HG21	1.75	0.69
4:D:204:ASN:O	4:D:205:ASN:CB	2.41	0.68
1:A:129:ASP:OD1	1:A:129:ASP:C	2.32	0.67
5:E:214:GLN:HA	5:E:254:ARG:O	1.95	0.66
1:A:17:ARG:CG	1:A:17:ARG:HH11	2.09	0.66
4:D:205:ASN:O	4:D:205:ASN:ND2	2.30	0.65
1:A:255:GLN:HA	1:A:255:GLN:OE1	1.97	0.65
1:A:273:ARG:HH11	1:A:273:ARG:CG	2.02	0.64
5:E:9:THR:CG2	5:E:10:HIS:HD2	2.04	0.63
4:D:59:ASN:HD21	4:D:83:LYS:HE2	1.65	0.61
4:D:209:PRO:O	4:D:210:GLU:HB2	1.99	0.61

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:103:LEU:HD11	1:A:107:GLY:HA2	1.83	0.60
4:D:210:GLU:H	4:D:212:THR:HG23	1.66	0.60
4:D:94:ALA:HB1	4:D:124:ILE:HG12	1.84	0.60
1:A:28:VAL:O	1:A:29:ASP:HB2	2.04	0.58
4:D:15:GLU:HG2	4:D:126:PRO:HA	1.85	0.58
5:E:179:HIS:O	5:E:182:VAL:HG22	2.05	0.56
1:A:121:LYS:HE3	2:B:1:ILE:HD11	1.87	0.56
1:A:260:HIS:CE1	1:A:271:THR:HG23	2.41	0.56
4:D:213:PHE:CE2	4:D:215:PRO:HG3	2.41	0.55
4:D:209:PRO:HB2	4:D:212:THR:HG22	1.88	0.55
4:D:215:PRO:O	4:D:216:SER:HB2	2.07	0.55
5:E:63:GLU:HA	5:E:80:GLN:HB3	1.89	0.54
1:A:250:PRO:HB2	1:A:253:GLU:HG3	1.91	0.53
4:D:146:LYS:N	4:D:147:SER:CB	2.72	0.53
2:B:1:ILE:HD12	2:B:1:ILE:H	1.72	0.53
1:A:223:ASP:HB3	1:A:225:THR:HG22	1.90	0.53
5:E:192:GLN:C	5:E:194:ALA:H	2.13	0.52
5:E:18:GLN:HG3	5:E:92:ASN:HB3	1.92	0.52
1:A:203:CYS:CB	1:A:259:CYS:SG	2.98	0.51
5:E:20:THR:C	5:E:21:LEU:HD23	2.31	0.51
5:E:229:LEU:HD12	5:E:242:PRO:HD2	1.92	0.51
4:D:209:PRO:C	4:D:210:GLU:CG	2.77	0.50
5:E:10:HIS:CG	5:E:227:TYR:HB3	2.46	0.50
4:D:196:ASP:C	4:D:196:ASP:OD1	2.48	0.50
2:B:24:ASN:HB3	2:B:65:LEU:HD11	1.94	0.50
1:A:129:ASP:OD2	1:A:132:SER:OG	2.30	0.50
1:A:103:LEU:HD13	1:A:165:VAL:HG22	1.94	0.50
1:A:129:ASP:OD1	1:A:129:ASP:O	2.30	0.49
4:D:210:GLU:H	4:D:212:THR:CG2	2.24	0.49
1:A:129:ASP:OD1	1:A:131:SER:N	2.40	0.49
5:E:105:ALA:HA	5:E:116:TYR:O	2.12	0.49
1:A:127:ASN:O	1:A:129:ASP:O	2.29	0.49
4:D:196:ASP:OD1	4:D:196:ASP:O	2.30	0.49
1:A:81:LEU:HD11	3:C:13:TYR:HB3	1.94	0.49
5:E:39:VAL:HG13	5:E:104:CYS:SG	2.53	0.49
4:D:43:ARG:HB2	4:D:53:LEU:HD11	1.95	0.48
1:A:187:THR:HB	1:A:272:LEU:HD11	1.94	0.48
2:B:16:GLU:HB3	2:B:19:LYS:HG3	1.95	0.48
4:D:163:SER:HB2	4:D:170:ILE:CD1	2.43	0.48
1:A:139:ALA:HA	1:A:142:ILE:HD12	1.95	0.47
4:D:13:MET:HB2	4:D:19:TYR:CE2	2.48	0.47
1:A:35:ARG:HG2	1:A:48:ARG:HD3	1.97	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:E:108:ARG:HG2	5:E:111:SER:HB2	1.97	0.47
4:D:43:ARG:HD3	8:D:330:HOH:O	2.13	0.47
5:E:134:PRO:HD3	5:E:242:PRO:HB3	1.97	0.47
4:D:151:PHE:HB2	4:D:203:PHE:CE1	2.50	0.47
5:E:192:GLN:C	5:E:194:ALA:N	2.68	0.46
1:A:261:VAL:HB	1:A:270:LEU:HB2	1.97	0.46
4:D:164:LYS:HE3	4:D:205:ASN:OD1	2.16	0.46
5:E:75:ARG:HH22	5:E:98:ASP:CG	2.20	0.46
1:A:220:ASP:OD1	1:A:256:ARG:HD2	2.16	0.45
4:D:75:ARG:HD2	4:D:98:LEU:HD11	1.99	0.45
5:E:147:ILE:HG23	5:E:210:ALA:HB1	1.98	0.45
5:E:188:PRO:HB2	5:E:200:TYR:HB3	1.98	0.44
5:E:14:THR:HA	5:E:126:THR:O	2.18	0.44
1:A:123:TYR:HD2	1:A:124:ILE:HG22	1.83	0.44
4:D:167:ASP:HB3	4:D:194:LYS:HE3	1.99	0.43
1:A:66:ILE:HA	3:C:5:LEU:HD21	2.01	0.43
1:A:36:PHE:CE2	1:A:43:PRO:HB2	2.53	0.43
1:A:199:ALA:O	1:A:249:VAL:HG22	2.19	0.43
1:A:104:GLY:CA	1:A:110:LEU:HB2	2.49	0.43
1:A:31:THR:HG23	1:A:239:ARG:HD3	2.00	0.42
4:D:21:ILE:HD11	4:D:89:LEU:HD23	2.01	0.42
1:A:1:GLY:HA3	1:A:2:SER:HA	1.74	0.42
1:A:219:ARG:HD3	1:A:257:TYR:OH	2.19	0.42
1:A:8:PHE:CE2	1:A:98:MET:HG3	2.54	0.42
4:D:125:LYS:HB3	4:D:156:SER:HB3	2.02	0.42
5:E:56:TYR:HB3	5:E:80:GLN:HB2	2.01	0.42
1:A:49:ALA:O	1:A:52:ILE:HG22	2.20	0.41
1:A:234:ARG:HG3	2:B:10:TYR:CE2	2.55	0.41
4:D:13:MET:CB	4:D:19:TYR:CE2	3.03	0.41
5:E:105:ALA:HB1	5:E:115:GLN:HG2	2.02	0.41
2:B:23:LEU:HB2	2:B:70:PHE:CD2	2.55	0.41
2:B:51:HIS:HA	2:B:65:LEU:O	2.21	0.41
2:B:49:VAL:HG22	2:B:68:THR:HB	2.03	0.41
1:A:45:THR:HG21	1:A:63:ASN:HB3	2.02	0.41
2:B:23:LEU:O	2:B:67:TYR:HA	2.21	0.41
2:B:39:LEU:HB3	2:B:46:ILE:HD12	2.02	0.41
5:E:192:GLN:O	5:E:194:ALA:N	2.53	0.41
1:A:177:GLU:OE1	1:A:177:GLU:HA	2.20	0.41
1:A:258:THR:HG22	1:A:260:HIS:NE2	2.36	0.41
4:D:44:GLN:HB2	4:D:50:LEU:HD23	2.03	0.41
1:A:234:ARG:HG3	2:B:10:TYR:CZ	2.57	0.40
2:B:27:VAL:HG21	2:B:37:VAL:HG21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/276 (99%)	267 (97%)	7 (3%)	0	100	100
2	B	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
3	C	11/13 (85%)	11 (100%)	0	0	100	100
4	D	199/201 (99%)	196 (98%)	3 (2%)	0	100	100
5	E	240/242 (99%)	233 (97%)	7 (3%)	0	100	100
All	All	822/832 (99%)	802 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/234 (100%)	207 (88%)	27 (12%)	8	23
2	B	95/95 (100%)	85 (90%)	10 (10%)	10	27
3	C	11/11 (100%)	9 (82%)	2 (18%)	2	7
4	D	178/178 (100%)	151 (85%)	27 (15%)	4	12
5	E	210/210 (100%)	195 (93%)	15 (7%)	21	51
All	All	728/728 (100%)	647 (89%)	81 (11%)	9	25

All (81) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	12	MET
1	A	14	ARG
1	A	17	ARG
1	A	23	ILE
1	A	34	VAL
1	A	35	ARG
1	A	75	ARG
1	A	76	GLU
1	A	98	MET
1	A	111	ARG
1	A	132	SER
1	A	134	THR
1	A	146	LYS
1	A	156	ARG
1	A	163	LEU
1	A	165	VAL
1	A	176	LYS
1	A	177	GLU
1	A	194	VAL
1	A	216	THR
1	A	222	GLU
1	A	225	THR
1	A	232	GLU
1	A	248	VAL
1	A	251	SER
1	A	273	ARG
2	B	0	MET
2	B	3	ARG
2	B	16	GLU
2	B	20	SER
2	B	40	LEU
2	B	45	ARG
2	B	47	GLU
2	B	48	LYS
2	B	58	LYS
2	B	70	PHE
3	C	3	GLU
3	C	9	GLN
4	D	5	GLU
4	D	10	PHE
4	D	12	SER
4	D	13	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
4	D	68	GLU
4	D	79	SER
4	D	80	LEU
4	D	122	LEU
4	D	129	GLN
4	D	141	SER
4	D	142	LYS
4	D	143	SER
4	D	144	SER
4	D	162	GLN
4	D	164	LYS
4	D	166	SER
4	D	173	LYS
4	D	174	CYS
4	D	193	ASN
4	D	195	SER
4	D	201	ASN
4	D	208	ILE
4	D	209	PRO
4	D	210	GLU
4	D	211	ASP
4	D	212	THR
4	D	216	SER
5	E	21	LEU
5	E	22	ARG
5	E	40	SER
5	E	46	LEU
5	E	54	PHE
5	E	55	GLN
5	E	75	ARG
5	E	77	SER
5	E	109	THR
5	E	139	VAL
5	E	195	LEU
5	E	205	ARG
5	E	231	GLU
5	E	236	THR
5	E	237	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
4	D	59	ASN
4	D	96	HIS
4	D	162	GLN
5	E	10	HIS
5	E	55	GLN
5	E	92	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	276/276 (100%)	-0.04	0 100 100	31, 50, 70, 87	25 (9%)
2	B	100/100 (100%)	0.02	1 (1%) 79 79	34, 45, 67, 75	6 (6%)
3	C	13/13 (100%)	-0.03	0 100 100	34, 51, 72, 73	0
4	D	201/201 (100%)	0.28	5 (2%) 54 55	27, 42, 78, 91	11 (5%)
5	E	242/242 (100%)	0.08	4 (1%) 67 68	23, 38, 55, 112	14 (5%)
All	All	832/832 (100%)	0.08	10 (1%) 75 76	23, 44, 72, 112	56 (6%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	215	PRO	3.6
4	D	209	PRO	2.6
4	D	216	SER	2.6
5	E	21	LEU	2.4
5	E	46	LEU	2.3
5	E	256	ASP	2.2
4	D	141	SER	2.2
2	B	1	ILE	2.1
4	D	162	GLN	2.1
5	E	11	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.



## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	MG	B	103	1/1	0.25	2.91	53,53,53,53	0
7	MG	B	102	1/1	0.17	-0.79	60,60,60,60	0
6	NA	B	101	1/1	0.14	-0.92	44,44,44,44	0

## 6.5 Other polymers

There are no such residues in this entry.