



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 04:32 AM GMT

PDB ID : 1JYM  
Title : Crystals of Peptide Deformylase from Plasmodium falciparum with Ten Subunits per Asymmetric Unit Reveal Critical Characteristics of the Active Site for Drug Design  
Authors : Kumar, A.; Nguyen, K.T.; Srivathsan, S.; Ornstein, B.; Turley, S.; Hirsh, I.; Pei, D.; Hol, W.G.J.  
Deposited on : 2001-09-12  
Resolution : 2.80 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2295 (2.80-2.80)
Ramachandran outliers	78287	2252 (2.80-2.80)
Sidechain outliers	78261	2254 (2.80-2.80)
RSRZ outliers	66119	1802 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	183	
1	B	183	
1	C	183	
1	D	183	
1	E	183	
1	F	183	
1	G	183	
1	H	183	
1	I	183	
1	J	183	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14584 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptide Deformylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	179	Total	C	N	O	S	Se	0	0	0
			1487	953	263	267	1	3			
1	B	179	Total	C	N	O	S	Se	0	0	0
			1487	953	263	267	1	3			
1	C	168	Total	C	N	O	S	Se	0	0	0
			1394	898	246	246	1	3			
1	D	173	Total	C	N	O	S	Se	0	0	0
			1435	921	253	257	1	3			
1	E	176	Total	C	N	O	S	Se	0	0	0
			1458	936	256	262	1	3			
1	F	172	Total	C	N	O	S	Se	0	0	0
			1426	916	252	254	1	3			
1	G	172	Total	C	N	O	S	Se	0	0	0
			1425	915	252	254	1	3			
1	H	173	Total	C	N	O	S	Se	0	0	0
			1435	921	253	257	1	3			
1	I	172	Total	C	N	O	S	Se	0	0	0
			1425	915	252	254	1	3			
1	J	172	Total	C	N	O	S	Se	0	0	0
			1425	915	252	254	1	3			

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Co	0	0
			1	1		
2	J	1	Total	Co	0	0
			1	1		
2	D	1	Total	Co	0	0
			1	1		
2	E	1	Total	Co	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	1	Total 1	Co 1	0	0
2	B	1	Total 1	Co 1	0	0
2	I	1	Total 1	Co 1	0	0
2	C	1	Total 1	Co 1	0	0
2	A	1	Total 1	Co 1	0	0
2	F	1	Total 1	Co 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	22	Total 22	O 22	0	0
3	B	17	Total 17	O 17	0	0
3	C	17	Total 17	O 17	0	0
3	D	16	Total 16	O 16	0	0
3	E	19	Total 19	O 19	0	0
3	F	26	Total 26	O 26	0	0
3	G	13	Total 13	O 13	0	0
3	H	15	Total 15	O 15	0	0
3	I	16	Total 16	O 16	0	0
3	J	16	Total 16	O 16	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Peptide Deformylase

Chain A:




#### • Molecule 1: Peptide Deformylase

Chain B:




#### • Molecule 1: Peptide Deformylase

Chain C:




#### • Molecule 1: Peptide Deformylase

Chain D:









● Molecule 1: Peptide Deformylase

Chain J:



HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	121.26Å 121.26Å 177.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.80 29.41 – 2.80	Depositor EDS
% Data completeness (in resolution range)	82.5 (30.00-2.80) 82.5 (29.41-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.89 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, $R_{free}$	0.231 , 0.296 0.230 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	43.5	Xtriage
Anisotropy	0.414	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 29.0	EDS
Estimated twinning fraction	0.052 for h,-k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	1 of 51866 reflections (0.002%)	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	14584	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
CO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/1513	0.87	5/2026 (0.2%)
1	B	0.54	0/1513	0.91	8/2026 (0.4%)
1	C	0.63	0/1416	0.92	5/1893 (0.3%)
1	D	0.53	0/1458	0.86	6/1950 (0.3%)
1	E	0.59	0/1482	0.91	5/1984 (0.3%)
1	F	0.71	0/1449	0.97	4/1938 (0.2%)
1	G	0.59	0/1448	0.92	6/1937 (0.3%)
1	H	0.63	0/1458	0.94	8/1950 (0.4%)
1	I	0.55	0/1448	0.91	8/1937 (0.4%)
1	J	0.60	0/1448	0.94	5/1937 (0.3%)
All	All	0.59	0/14633	0.91	60/19578 (0.3%)

There are no bond length outliers.

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	199	ASP	CB-CG-OD2	7.67	125.20	118.30
1	A	199	ASP	CB-CG-OD2	7.13	124.72	118.30
1	C	208	ASP	CB-CG-OD2	7.07	124.67	118.30
1	B	214	ASP	CB-CG-OD2	7.06	124.66	118.30
1	A	85	ASP	CB-CG-OD2	6.95	124.56	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens

added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1487	0	1533	34	0
1	B	1487	0	1533	34	0
1	C	1394	0	1460	26	0
1	D	1435	0	1491	22	0
1	E	1458	0	1513	29	0
1	F	1426	0	1485	26	0
1	G	1425	0	1483	26	0
1	H	1435	0	1491	29	0
1	I	1425	0	1483	35	0
1	J	1425	0	1483	36	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
3	A	22	0	0	0	0
3	B	17	0	0	0	0
3	C	17	0	0	0	0
3	D	16	0	0	0	0
3	E	19	0	0	0	0
3	F	26	0	0	1	0
3	G	13	0	0	0	0
3	H	15	0	0	1	0
3	I	16	0	0	1	0
3	J	16	0	0	0	0
All	All	14584	0	14955	283	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

The worst 5 of 283 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:J:164:VAL:O	1:J:192:ARG:NH2	2.05	0.90

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:I:65:ILE:HA	1:I:98:ILE:HD12	1.57	0.86
1:D:117:ARG:HH22	1:D:202:ASN:HD21	1.26	0.84
1:F:65:ILE:HA	1:F:98:ILE:HD11	1.63	0.80
1:A:231:ALA:HA	1:A:234:SER:HB3	1.64	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/183 (97%)	164 (93%)	12 (7%)	1 (1%)	33	72
1	B	177/183 (97%)	157 (89%)	20 (11%)	0	100	100
1	C	166/183 (91%)	154 (93%)	11 (7%)	1 (1%)	33	72
1	D	171/183 (93%)	153 (90%)	17 (10%)	1 (1%)	33	72
1	E	174/183 (95%)	160 (92%)	11 (6%)	3 (2%)	14	42
1	F	170/183 (93%)	157 (92%)	8 (5%)	5 (3%)	7	23
1	G	170/183 (93%)	149 (88%)	20 (12%)	1 (1%)	33	72
1	H	171/183 (93%)	154 (90%)	15 (9%)	2 (1%)	19	54
1	I	170/183 (93%)	160 (94%)	8 (5%)	2 (1%)	19	54
1	J	170/183 (93%)	157 (92%)	12 (7%)	1 (1%)	33	72
All	All	1716/1830 (94%)	1565 (91%)	134 (8%)	17 (1%)	22	60

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	89	LYS
1	J	159	GLY
1	A	117	ARG
1	C	65	ILE
1	I	70	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/169 (99%)	138 (82%)	30 (18%)	2	7
1	B	168/169 (99%)	140 (83%)	28 (17%)	3	9
1	C	157/169 (93%)	130 (83%)	27 (17%)	3	8
1	D	162/169 (96%)	133 (82%)	29 (18%)	2	7
1	E	165/169 (98%)	129 (78%)	36 (22%)	1	4
1	F	161/169 (95%)	132 (82%)	29 (18%)	2	7
1	G	161/169 (95%)	128 (80%)	33 (20%)	2	5
1	H	162/169 (96%)	133 (82%)	29 (18%)	2	7
1	I	161/169 (95%)	131 (81%)	30 (19%)	2	7
1	J	161/169 (95%)	126 (78%)	35 (22%)	1	4
All	All	1626/1690 (96%)	1320 (81%)	306 (19%)	2	7

5 of 306 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	191	SER
1	F	191	SER
1	J	94	LYS
1	E	211	THR
1	F	79	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	111	GLN
1	F	202	ASN
1	J	132	ASN
1	F	132	ASN
1	C	111	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	179/183 (97%)	0.22	5 (2%)	50	52	32, 54, 96, 104	0
1	B	179/183 (97%)	0.28	10 (5%)	24	23	22, 46, 96, 106	0
1	C	168/183 (91%)	0.11	4 (2%)	56	57	13, 39, 81, 92	0
1	D	173/183 (94%)	0.15	4 (2%)	57	58	21, 47, 92, 102	0
1	E	176/183 (96%)	0.23	9 (5%)	27	27	23, 47, 99, 106	0
1	F	172/183 (93%)	-0.06	2 (1%)	75	76	15, 33, 80, 104	0
1	G	172/183 (93%)	0.19	9 (5%)	26	26	18, 43, 88, 104	0
1	H	173/183 (94%)	0.00	4 (2%)	57	58	17, 39, 82, 106	0
1	I	172/183 (93%)	0.21	7 (4%)	35	36	18, 42, 90, 105	0
1	J	172/183 (93%)	0.12	7 (4%)	35	36	20, 43, 89, 105	0
All	All	1736/1830 (94%)	0.15	61 (3%)	42	42	13, 43, 92, 106	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	63	ASP	15.3
1	E	63	ASP	13.4
1	J	63	ASP	11.7
1	I	63	ASP	10.5
1	G	63	ASP	10.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CO	F	5300	1/1	0.14	-0.69	24,24,24,24	0
2	CO	C	2300	1/1	0.13	-0.95	35,35,35,35	0
2	CO	A	300	1/1	0.10	-1.39	46,46,46,46	0
2	CO	G	6300	1/1	0.11	-1.69	30,30,30,30	0
2	CO	B	1300	1/1	0.11	-1.76	39,39,39,39	0
2	CO	H	7300	1/1	0.11	-1.90	34,34,34,34	0
2	CO	E	4300	1/1	0.11	-1.91	39,39,39,39	0
2	CO	I	8300	1/1	0.09	-2.04	45,45,45,45	0
2	CO	D	3300	1/1	0.10	-2.50	34,34,34,34	0
2	CO	J	9300	1/1	0.10	-3.50	44,44,44,44	0

### 6.5 Other polymers ⓘ

There are no such residues in this entry.