



# wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 08:11 AM GMT

PDB ID : 3JYM  
Title : Crystal Structure of the 3 FKBP domains of wheat FKBP73  
Authors : Dym, O.; Breiman, A.; Israel Structural Proteomics Center (ISPC)  
Deposited on : 2009-09-22  
Resolution : 2.28 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3861 (2.30-2.26)
Clashscore	79885	4801 (2.30-2.26)
Ramachandran outliers	78287	4729 (2.30-2.26)
Sidechain outliers	78261	4728 (2.30-2.26)
RSRZ outliers	66119	3864 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	377	
1	B	377	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4726 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FK506-binding protein (FKBP) from wheat.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	0	0
			2344	1501	378	460	5			
1	B	308	Total	C	N	O	S	0	0	0
			2382	1523	385	469	5			



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	186.96Å 31.01Å 68.98Å 90.00° 93.59° 90.00°	Depositor
Resolution (Å)	46.63 – 2.28 46.65 – 2.28	Depositor EDS
% Data completeness (in resolution range)	98.6 (46.63-2.28) 98.6 (46.65-2.28)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.97 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.270 , 0.343 0.271 , 0.342	Depositor DCC
$R_{free}$ test set	1834 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.6	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 15.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 36734 reflections	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	4726	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.68	32/2382 (1.3%)	1.29	10/3210 (0.3%)
1	B	1.54	19/2419 (0.8%)	1.25	11/3261 (0.3%)
All	All	1.61	51/4801 (1.1%)	1.27	21/6471 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	1
1	B	0	4
All	All	1	5

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	209	CYS	CB-SG	-12.13	1.61	1.82
1	A	209	CYS	CB-SG	-11.82	1.62	1.82
1	A	355	TYR	CD2-CE2	-7.48	1.28	1.39
1	A	244	GLY	N-CA	7.44	1.57	1.46
1	A	283	GLU	CB-CG	-7.09	1.38	1.52

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	247	GLY	N-CA-C	-8.12	92.80	113.10
1	A	240	ARG	NE-CZ-NH1	-7.66	116.47	120.30
1	B	316	ASP	CB-CG-OD1	-7.08	111.93	118.30
1	B	244	GLY	N-CA-C	-6.96	95.69	113.10
1	B	88	LYS	N-CA-C	-6.31	93.97	111.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	268	THR	CB

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	245	GLU	Peptide
1	B	155	LYS	Peptide
1	B	243	ALA	Peptide
1	B	286	GLY	Peptide
1	B	87	PHE	Peptide

## 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2344	0	2347	63	0
1	B	2382	0	2392	85	0
All	All	4726	0	4739	148	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 148 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:273:ASP:OD1	1:A:275:LYS:HD2	1.27	1.33
1:B:273:ASP:OD1	1:B:275:LYS:CD	1.90	1.18
1:B:273:ASP:OD1	1:B:275:LYS:HD2	0.93	1.10
1:B:243:ALA:HB1	1:B:244:GLY:HA3	1.11	1.06
1:A:65:VAL:O	1:A:87:PHE:N	2.00	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/377 (78%)	261 (89%)	17 (6%)	14 (5%)	4	1
1	B	299/377 (79%)	266 (89%)	23 (8%)	10 (3%)	6	3
All	All	591/754 (78%)	527 (89%)	40 (7%)	24 (4%)	4	2

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	109	LYS
1	A	246	GLY
1	B	134	ALA
1	B	135	ASN
1	B	287	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/311 (82%)	226 (89%)	28 (11%)	9	9
1	B	258/311 (83%)	223 (86%)	35 (14%)	5	5
All	All	512/622 (82%)	449 (88%)	63 (12%)	7	6

5 of 63 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	38	ILE
1	B	114	LEU
1	B	348	LEU
1	B	56	THR

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	91	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	HIS
1	A	232	GLN
1	B	207	HIS
1	B	232	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	300/377 (79%)	0.52	29 (9%) 8 12	8, 26, 67, 81	0
1	B	307/377 (81%)	0.44	25 (8%) 12 17	11, 30, 61, 80	0
All	All	607/754 (80%)	0.48	54 (8%) 10 15	8, 28, 63, 81	0

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	100	TRP	8.7
1	A	59	VAL	6.3
1	A	54	TRP	6.0
1	B	96	VAL	5.5
1	B	97	ILE	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

### 6.5 Other polymers

There are no such residues in this entry.