



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 04:15 PM GMT

PDB ID : 1K0R
Title : Crystal Structure of Mycobacterium tuberculosis NusA
Authors : Gopal, B.; Haire, L.F.; Gamblin, S.J.; Dodson, E.J.; Lane, A.N.; Papavinasundaram, K.G.; Colston, M.J.; Dodson, G.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 2001-09-20
Resolution : 1.70 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2456 (1.70-1.70)
Clashscore	79885	2929 (1.70-1.70)
Ramachandran outliers	78287	2878 (1.70-1.70)
Sidechain outliers	78261	2878 (1.70-1.70)
RSRZ outliers	66119	2456 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	366	
1	B	366	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	SO4	B	509	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5367 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NusA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	823	0	0
			2482	1526	476	473	7			
1	B	326	Total	C	N	O	S	6	0	0
			2470	1518	474	471	7			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	EXPRESSION TAG	UNP P0A5M2
A	-17	GLY	-	EXPRESSION TAG	UNP P0A5M2
A	-16	SER	-	EXPRESSION TAG	UNP P0A5M2
A	-15	SER	-	EXPRESSION TAG	UNP P0A5M2
A	-14	HIS	-	EXPRESSION TAG	UNP P0A5M2
A	-13	HIS	-	EXPRESSION TAG	UNP P0A5M2
A	-12	HIS	-	EXPRESSION TAG	UNP P0A5M2
A	-11	HIS	-	EXPRESSION TAG	UNP P0A5M2
A	-10	HIS	-	EXPRESSION TAG	UNP P0A5M2
A	-9	HIS	-	EXPRESSION TAG	UNP P0A5M2
A	-8	SER	-	EXPRESSION TAG	UNP P0A5M2
A	-7	SER	-	EXPRESSION TAG	UNP P0A5M2
A	-6	GLY	-	EXPRESSION TAG	UNP P0A5M2
A	-5	LEU	-	EXPRESSION TAG	UNP P0A5M2
A	-4	VAL	-	EXPRESSION TAG	UNP P0A5M2
A	-3	SER	-	EXPRESSION TAG	UNP P0A5M2
A	-2	ARG	-	EXPRESSION TAG	UNP P0A5M2
A	-1	ARG	-	EXPRESSION TAG	UNP P0A5M2
A	0	HIS	-	EXPRESSION TAG	UNP P0A5M2
B	-18	MET	-	EXPRESSION TAG	UNP P0A5M2
B	-17	GLY	-	EXPRESSION TAG	UNP P0A5M2
B	-16	SER	-	EXPRESSION TAG	UNP P0A5M2
B	-15	SER	-	EXPRESSION TAG	UNP P0A5M2
B	-14	HIS	-	EXPRESSION TAG	UNP P0A5M2
B	-13	HIS	-	EXPRESSION TAG	UNP P0A5M2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	HIS	-	EXPRESSION TAG	UNP P0A5M2
B	-11	HIS	-	EXPRESSION TAG	UNP P0A5M2
B	-10	HIS	-	EXPRESSION TAG	UNP P0A5M2
B	-9	HIS	-	EXPRESSION TAG	UNP P0A5M2
B	-8	SER	-	EXPRESSION TAG	UNP P0A5M2
B	-7	SER	-	EXPRESSION TAG	UNP P0A5M2
B	-6	GLY	-	EXPRESSION TAG	UNP P0A5M2
B	-5	LEU	-	EXPRESSION TAG	UNP P0A5M2
B	-4	VAL	-	EXPRESSION TAG	UNP P0A5M2
B	-3	SER	-	EXPRESSION TAG	UNP P0A5M2
B	-2	ARG	-	EXPRESSION TAG	UNP P0A5M2
B	-1	ARG	-	EXPRESSION TAG	UNP P0A5M2
B	0	HIS	-	EXPRESSION TAG	UNP P0A5M2

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	165	Total 165	O 165	0	0
3	B	230	Total 230	O 230	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NusA

Chain A: 



• Molecule 1: NusA

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.99Å 88.99Å 180.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 1.70 18.38 – 1.70	Depositor EDS
% Data completeness (in resolution range)	86.2 (15.00-1.70) 91.0 (18.38-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 1.70Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.248 , 0.271 0.251 , 0.260	Depositor DCC
R_{free} test set	4160 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 55.4	EDS
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 82635 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5367	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2512	0.53	0/3399
1	B	0.28	0/2500	0.56	0/3381
All	All	0.27	0/5012	0.55	0/6780

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2482	0	2512	25	1
1	B	2470	0	2496	55	0
2	A	5	0	0	0	0
2	B	15	0	0	0	0
3	A	165	0	0	6	0
3	B	230	0	0	12	0
All	All	5367	0	5008	76	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (76) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:185:HIS:HD2	1:A:187:ASN:H	1.22	0.86
1:B:185:HIS:HD2	1:B:187:ASN:H	1.23	0.83
1:A:124:ARG:NH2	1:B:128:ARG:HE	1.81	0.79
1:B:12:GLU:HG3	1:B:13:VAL:HG23	1.64	0.79
1:A:124:ARG:HH22	1:B:128:ARG:HE	1.31	0.76
1:B:278:VAL:HG23	1:B:295:PRO:HD3	1.77	0.67
1:A:207:ILE:HD13	1:A:220:ILE:HD11	1.79	0.64
1:A:185:HIS:CD2	1:A:187:ASN:H	2.12	0.63
1:B:277:VAL:HG13	1:B:293:VAL:O	1.98	0.63
1:B:277:VAL:HG11	1:B:292:VAL:HG13	1.81	0.62
1:A:213:GLU:HG3	1:A:312:LEU:HD21	1.80	0.62
1:B:213:GLU:OE1	1:B:308:GLN:HG3	2.00	0.61
1:B:18:SER:HB2	1:B:21:GLU:HB3	1.82	0.61
1:B:328:PRO:HB2	3:B:710:HOH:O	2.03	0.58
1:A:212:ARG:HG3	1:A:218:SER:OG	2.04	0.58
1:B:327:ALA:HA	3:B:668:HOH:O	2.06	0.55
1:B:185:HIS:CD2	1:B:187:ASN:H	2.14	0.55
1:B:61:GLU:HB2	1:B:69:ILE:HG21	1.89	0.55
1:B:17:ILE:HA	3:B:698:HOH:O	2.06	0.54
1:B:18:SER:CB	1:B:21:GLU:HB3	2.37	0.54
1:B:213:GLU:HG3	1:B:312:LEU:HD21	1.90	0.53
1:B:34:TYR:O	1:B:37:THR:HG22	2.09	0.52
1:B:69:ILE:HD12	1:B:69:ILE:N	2.25	0.51
1:A:143:GLU:OE2	1:B:128:ARG:NH1	2.39	0.51
1:B:277:VAL:HG12	1:B:278:VAL:N	2.27	0.50
1:B:177:PRO:HD3	3:B:694:HOH:O	2.12	0.49
1:B:137:THR:HB	3:B:607:HOH:O	2.13	0.49
1:A:213:GLU:CG	1:A:312:LEU:HD21	2.43	0.49
1:B:60:ARG:HD2	3:B:729:HOH:O	2.11	0.48
1:A:111:GLU:HG3	3:A:561:HOH:O	2.12	0.48
1:B:216:HIS:HB3	1:B:308:GLN:NE2	2.28	0.47
1:A:285:GLN:HG2	3:A:559:HOH:O	2.14	0.47
1:A:119:ILE:HD12	1:A:161:ASN:HB2	1.96	0.47
1:A:110:ARG:HB2	1:A:113:GLU:OE2	2.15	0.47
1:B:35:ARG:HG2	3:B:663:HOH:O	2.14	0.47
1:B:22:LEU:O	1:B:25:THR:HG22	2.14	0.47
1:A:285:GLN:HG3	1:A:286:THR:N	2.30	0.46
1:B:17:ILE:HB	3:B:734:HOH:O	2.15	0.46
1:B:111:GLU:HG2	1:B:169:GLY:HA2	1.96	0.46
1:B:220:ILE:HG13	1:B:220:ILE:O	2.15	0.46
1:B:172:ARG:HG3	1:B:172:ARG:O	2.15	0.46
1:B:295:PRO:HG2	1:B:298:GLN:CG	2.46	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:277:VAL:CG1	1:B:278:VAL:N	2.79	0.45
1:B:69:ILE:O	1:B:70:SER:C	2.53	0.45
1:B:276:LYS:H	1:B:298:GLN:HE21	1.64	0.45
1:A:276:LYS:HE2	1:A:276:LYS:HA	1.97	0.45
1:B:277:VAL:HG11	1:B:292:VAL:CG1	2.46	0.45
1:A:207:ILE:CD1	1:A:220:ILE:HD11	2.45	0.45
1:B:30:LEU:HD21	1:B:83:ALA:HA	1.98	0.45
1:B:94:ARG:HG2	1:B:94:ARG:HH11	1.83	0.44
1:B:61:GLU:CD	1:B:69:ILE:HG21	2.38	0.44
1:A:303:ILE:HG22	1:A:307:GLY:HA2	2.00	0.44
1:B:59:ALA:HB3	1:B:72:TRP:CE2	2.53	0.43
1:B:92:LEU:O	1:B:96:ARG:HG3	2.17	0.43
1:A:108:SER:N	1:A:138:GLU:HB2	2.33	0.43
1:A:124:ARG:HH22	1:B:128:ARG:NE	2.09	0.43
1:B:164:ARG:NE	3:B:728:HOH:O	2.47	0.43
1:B:276:LYS:H	1:B:298:GLN:NE2	2.17	0.43
1:B:326:ASP:C	1:B:328:PRO:HD3	2.40	0.42
1:B:327:ALA:N	1:B:328:PRO:HD3	2.35	0.42
1:B:28:SER:O	1:B:32:THR:HG23	2.20	0.42
1:A:176:GLU:HA	1:A:177:PRO:HD3	1.91	0.42
1:B:212:ARG:HG3	1:B:218:SER:OG	2.20	0.42
1:B:247:MET:HG2	1:B:253:GLU:O	2.20	0.42
1:A:276:LYS:HE2	1:A:276:LYS:CA	2.50	0.42
1:B:153:PRO:HG2	3:B:636:HOH:O	2.20	0.42
1:A:187:ASN:ND2	3:A:588:HOH:O	2.53	0.41
1:A:285:GLN:HB2	3:A:659:HOH:O	2.20	0.41
1:B:61:GLU:HB2	1:B:69:ILE:CG2	2.50	0.41
1:B:19:VAL:HG12	1:B:19:VAL:O	2.21	0.41
1:B:61:GLU:HB2	1:B:69:ILE:CB	2.51	0.41
1:B:175:ARG:HD3	1:B:175:ARG:HA	1.95	0.40
1:A:329:PRO:HD2	3:A:643:HOH:O	2.21	0.40
1:A:161:ASN:ND2	3:A:614:HOH:O	2.54	0.40
1:B:149:ALA:HB3	3:B:611:HOH:O	2.22	0.40
1:B:56:ARG:HD2	3:B:696:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:291:ARG:NH2	1:A:291:ARG:NH2[4_555]	1.95	0.25

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/366 (88%)	285 (88%)	28 (9%)	9 (3%)	8	1
1	B	322/366 (88%)	296 (92%)	19 (6%)	7 (2%)	10	1
All	All	644/732 (88%)	581 (90%)	47 (7%)	16 (2%)	9	1

All (16) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	15	ARG
1	A	22	LEU
1	A	68	LEU
1	B	20	ASN
1	B	21	GLU
1	A	64	GLU
1	A	81	ARG
1	B	61	GLU
1	B	63	ASP
1	A	15	ARG
1	A	16	GLY
1	A	43	ASP
1	A	174	ALA
1	B	13	VAL
1	B	11	ILE
1	A	13	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/294 (88%)	253 (97%)	7 (3%)	57	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	258/294 (88%)	252 (98%)	6 (2%)	63	41
All	All	518/588 (88%)	505 (98%)	13 (2%)	60	37

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	15	ARG
1	A	72	TRP
1	A	94	ARG
1	A	110	ARG
1	A	163	LEU
1	A	276	LYS
1	B	56	ARG
1	B	61	GLU
1	B	94	ARG
1	B	121	ARG
1	B	138	GLU
1	B	172	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	126	ASN
1	A	161	ASN
1	A	185	HIS
1	A	187	ASN
1	A	241	GLN
1	A	298	GLN
1	B	126	ASN
1	B	159	HIS
1	B	161	ASN
1	B	185	HIS
1	B	187	ASN
1	B	298	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	506	-	4,4,4	0.29	0	6,6,6	0.10	0
2	SO4	B	505	-	4,4,4	0.15	0	6,6,6	0.09	0
2	SO4	B	508	-	4,4,4	0.25	0	6,6,6	0.07	0
2	SO4	B	509	-	4,4,4	0.24	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	506	-	-	0/0/0/0	0/0/0/0
2	SO4	B	505	-	-	0/0/0/0	0/0/0/0
2	SO4	B	508	-	-	0/0/0/0	0/0/0/0
2	SO4	B	509	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	222/366 (60%)	0.81	30 (13%) 4 5	19, 28, 47, 73	1 (0%)
1	B	326/366 (89%)	1.98	75 (23%) 1 2	18, 31, 79, 97	2 (0%)
All	All	548/732 (74%)	1.51	105 (19%) 2 3	18, 29, 73, 97	3 (0%)

All (105) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	69	ILE	23.3
1	B	328	PRO	22.6
1	B	14	ASP	20.9
1	B	19	VAL	20.8
1	B	329	PRO	19.7
1	B	-4	VAL	19.5
1	B	65	ALA	18.4
1	B	98	ALA	18.2
1	B	174	ALA	15.8
1	B	62	THR	14.8
1	B	16	GLY	14.7
1	B	13	VAL	14.4
1	B	67	ASN	13.4
1	B	66	GLY	13.2
1	B	15	ARG	12.3
1	A	174	ALA	12.0
1	B	64	GLU	11.4
1	B	63	ASP	11.2
1	B	99	GLU	10.7
1	A	175	ARG	10.6
1	B	327	ALA	10.6
1	B	68	LEU	10.0
1	B	20	ASN	9.7
1	A	173	GLY	9.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	17	ILE	9.5
1	A	329	PRO	9.4
1	B	176	GLU	9.3
1	B	61	GLU	8.3
1	B	175	ARG	8.2
1	B	18	SER	8.0
1	B	173	GLY	7.9
1	B	38	GLN	7.5
1	A	172	ARG	7.4
1	B	172	ARG	7.3
1	B	21	GLU	7.0
1	B	39	GLY	6.6
1	B	43	ASP	6.4
1	B	60	ARG	5.9
1	A	176	GLU	5.7
1	B	37	THR	5.4
1	B	97	ASP	5.4
1	A	121	ARG	5.0
1	B	40	HIS	4.8
1	B	22	LEU	4.7
1	A	177	PRO	4.6
1	B	36	HIS	4.3
1	B	11	ILE	4.2
1	B	121	ARG	4.1
1	B	306	GLU	3.7
1	B	77	GLU	3.7
1	B	32	THR	3.6
1	B	41	GLN	3.5
1	B	12	GLU	3.4
1	B	123	SER	3.4
1	B	89	GLN	3.3
1	A	154	GLY	3.2
1	B	159	HIS	3.2
1	A	138	GLU	3.2
1	B	108	SER	3.2
1	A	241	GLN	3.1
1	B	24	GLU	3.0
1	B	124	ARG	3.0
1	B	305	LYS	3.0
1	A	238	PRO	3.0
1	B	220	ILE	3.0
1	B	207	ILE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	124	ARG	2.9
1	A	285	GLN	2.7
1	A	181	LEU	2.7
1	A	239	MET	2.7
1	A	255	ILE	2.7
1	A	266	ARG	2.7
1	B	321	ILE	2.7
1	A	171	THR	2.7
1	A	321	ILE	2.6
1	B	59	ALA	2.6
1	A	153	PRO	2.5
1	A	110	ARG	2.5
1	A	228	GLY	2.5
1	B	-3	SER	2.5
1	B	165	CYS	2.5
1	B	94	ARG	2.4
1	B	135	ILE	2.4
1	B	222	VAL	2.4
1	B	50	ARG	2.4
1	B	228	GLY	2.4
1	B	47	GLU	2.4
1	B	42	THR	2.3
1	A	306	GLU	2.3
1	B	44	ALA	2.3
1	B	189	VAL	2.3
1	B	310	ALA	2.3
1	A	189	VAL	2.2
1	B	158	GLU	2.2
1	B	181	LEU	2.2
1	A	159	HIS	2.2
1	A	222	VAL	2.1
1	B	110	ARG	2.1
1	B	299	LEU	2.1
1	A	292	VAL	2.1
1	B	266	ARG	2.1
1	A	272	LEU	2.1
1	B	145	VAL	2.0
1	B	8	LEU	2.0
1	A	145	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	B	509	5/5	0.50	4.10	95,95,95,96	0
2	SO4	B	508	5/5	0.43	0.92	87,87,88,88	0
2	SO4	B	505	5/5	0.16	-0.29	54,55,56,56	0
2	SO4	A	506	5/5	0.08	-0.88	45,45,45,47	0

6.5 Other polymers ⓘ

There are no such residues in this entry.