



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 03:57 AM GMT

PDB ID : 3K5B
Title : Crystal structure of the peripheral stalk of *Thermus thermophilus* H⁺-ATPase/synthase
Authors : Lee, L.K.; Stewart, A.G.; Donohoe, M.; Bernal, R.A.; Stock, D.
Deposited on : 2009-10-07
Resolution : 3.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1007 (3.18-3.02)
Clashscore	79885	1078 (3.16-3.04)
Ramachandran outliers	78287	1044 (3.16-3.04)
Sidechain outliers	78261	1044 (3.16-3.04)
RSRZ outliers	66119	1008 (3.18-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	B	104	
1	G	104	
2	A	188	
2	E	188	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4056 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type ATP synthase, subunit (VAPC-THERM).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	100	Total	C	N	O	Se	0	0	0
			747	460	136	150	1			
1	B	100	Total	C	N	O	Se	0	0	0
			720	442	134	143	1			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	17	GLY	-	EXPRESSION TAG	UNP Q5SIT5
B	17	GLY	-	EXPRESSION TAG	UNP Q5SIT5

- Molecule 2 is a protein called V-type ATP synthase subunit E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	184	Total	C	N	O	Se	0	0	0
			1312	815	242	252	3			
2	A	185	Total	C	N	O	Se	0	0	0
			1277	788	240	246	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	134	MSE	LEU	ENGINEERED	UNP P74901
E	171	MSE	LEU	ENGINEERED	UNP P74901
E	178	MSE	LEU	ENGINEERED	UNP P74901
A	134	MSE	LEU	ENGINEERED	UNP P74901
A	171	MSE	LEU	ENGINEERED	UNP P74901
A	178	MSE	LEU	ENGINEERED	UNP P74901

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: V-type ATP synthase, subunit (VAPC-THERM)

Chain G: 




- Molecule 1: V-type ATP synthase, subunit (VAPC-THERM)

Chain B: 




- Molecule 2: V-type ATP synthase subunit E

Chain E: 



- Molecule 2: V-type ATP synthase subunit E

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.78Å 79.75Å 75.80Å 90.00° 97.64° 90.00°	Depositor
Resolution (Å)	29.66 – 3.10 29.65 – 3.10	Depositor EDS
% Data completeness (in resolution range)	94.4 (29.66-3.10) 98.4 (29.65-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.232 , 0.288 0.261 , 0.303	Depositor DCC
R_{free} test set	857 reflections (5.62%)	DCC
Wilson B-factor (Å ²)	106.8	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 57.2	EDS
Estimated twinning fraction	0.536 for H, K, L 0.464 for L, -K, H 0.139 for l,-k,h	Xtriage
Reported twinning fraction	0.536 for H, K, L 0.464 for L, -K, H	Depositor
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 16171 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	4056	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.32	0/720	0.50	0/968
1	G	0.34	0/748	0.48	0/1001
2	A	0.32	0/1287	0.49	0/1748
2	E	0.34	0/1322	0.53	0/1789
All	All	0.33	0/4077	0.50	0/5506

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	720	0	678	17	0
1	G	747	0	726	21	0
2	A	1277	0	1161	29	0
2	E	1312	0	1244	51	0
All	All	4056	0	3809	102	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:E:87:VAL:HG12	2:E:171:MSE:HE2	1.20	1.09
1:G:110:ALA:HB1	2:E:182:VAL:HG23	1.57	0.85
2:E:149:ARG:O	2:E:150:LEU:HD13	1.84	0.76
2:E:41:LEU:O	2:E:41:LEU:HD13	1.86	0.76
2:E:88:ARG:HA	2:E:171:MSE:HE1	1.69	0.73

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	98/104 (94%)	91 (93%)	5 (5%)	2 (2%)	11	49
1	G	98/104 (94%)	90 (92%)	6 (6%)	2 (2%)	11	49
2	A	181/188 (96%)	151 (83%)	22 (12%)	8 (4%)	4	25
2	E	180/188 (96%)	156 (87%)	14 (8%)	10 (6%)	3	19
All	All	557/584 (95%)	488 (88%)	47 (8%)	22 (4%)	5	29

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	22	LEU
2	E	160	LYS
1	B	23	ILE
2	A	140	VAL
2	E	5	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	55/75 (73%)	48 (87%)	7 (13%)	6	24
1	G	62/75 (83%)	59 (95%)	3 (5%)	35	77
2	A	97/137 (71%)	92 (95%)	5 (5%)	32	73
2	E	108/137 (79%)	94 (87%)	14 (13%)	6	23
All	All	322/424 (76%)	293 (91%)	29 (9%)	14	47

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	149	ARG
2	E	184	GLN
2	A	84	ARG
2	E	165	ASN
1	B	30	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	B	100/104 (96%)	-0.34	0	100 100	52, 63, 81, 92	0
1	G	100/104 (96%)	0.21	0	100 100	45, 56, 73, 86	3 (3%)
2	A	185/188 (98%)	-0.21	0	100 100	49, 64, 76, 77	2 (1%)
2	E	184/188 (97%)	0.10	1 (0%)	88 39	36, 58, 74, 79	6 (3%)
All	All	569/584 (97%)	-0.06	1 (0%)	93 61	36, 60, 76, 92	11 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	158	GLU	3.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.