



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 04:27 AM GMT

PDB ID : 4K5D  
Title : Structure of neuronal nitric oxide synthase heme domain in complex with (S)  
-1,2-bis((2-amino-4-methylpyridin-6-yl)-methoxy)-propan-3-amine  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2013-04-14  
Resolution : 2.10 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	ACT	A	803	-	X

## 2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 7119 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	0	0
			3313	2121	566	605	21			
1	B	411	Total	C	N	O	S	0	0	0
			3345	2140	574	610	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 6,6'-{[(2S)-3-AMINOPROPANE-1,2-DIYL]BIS(OXYMETHANEDIYL)}BIS(4-METHYLPYRIDIN-2-AMINE) (three-letter code: 1Q6) (formula: C<sub>17</sub>H<sub>25</sub>N<sub>5</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			24	17	5	2		
5	B	1	Total	C	N	O	0	0
			24	17	5	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	106	Total	O	0	0
			106	106		
7	B	178	Total	O	0	0
			178	178		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nitric oxide synthase, brain

Chain A: 



- Molecule 1: Nitric oxide synthase, brain

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.96Å 110.93Å 164.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.19 – 2.10 43.92 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (92.19-2.10) 99.6 (43.92-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.40 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.190 , 0.233 0.190 , 0.233	Depositor DCC
$R_{free}$ test set	2786 reflections (5.19%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtriage
Anisotropy	0.774	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 40.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	4 of 56507 reflections (0.007%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7119	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, 1Q6, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	5/3406 (0.1%)	0.69	0/4621
1	B	0.72	8/3438 (0.2%)	0.70	0/4661
All	All	0.70	13/6844 (0.2%)	0.70	0/9282

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	421	TRP	CD2-CE2	6.22	1.48	1.41
1	B	510	TRP	CD2-CE2	5.69	1.48	1.41
1	B	711	TRP	CD2-CE2	5.59	1.48	1.41
1	A	510	TRP	CD2-CE2	5.51	1.48	1.41
1	A	561	TRP	CD2-CE2	5.45	1.47	1.41
1	B	561	TRP	CD2-CE2	5.44	1.47	1.41
1	A	421	TRP	CD2-CE2	5.41	1.47	1.41
1	B	716	TRP	CD2-CE2	5.31	1.47	1.41
1	A	716	TRP	CD2-CE2	5.19	1.47	1.41
1	A	676	TRP	CD2-CE2	5.19	1.47	1.41
1	B	475	TRP	CD2-CE2	5.15	1.47	1.41
1	B	553	TRP	CD2-CE2	5.08	1.47	1.41
1	B	625	TRP	CD2-CE2	5.07	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens



added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3313	0	0	10	0
1	B	3345	0	0	4	0
2	A	43	0	0	2	0
2	B	43	0	0	2	0
3	A	17	0	0	0	0
3	B	17	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0
5	A	24	0	25	4	0
5	B	24	0	25	2	0
6	A	1	0	0	0	0
7	A	106	0	0	4	0
7	B	178	0	0	1	0
All	All	7119	0	50	17	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (17) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:706:TYR:OH	2:B:801:HEM:O2D	2.09	0.70
5:A:804:1Q6:H4	7:A:917:HOH:O	1.98	0.63
1:A:403:TYR:CE1	1:A:407:HIS:CE1	2.92	0.58
1:A:307:GLU:CG	7:B:908:HOH:O	2.58	0.51
1:A:350:THR:N	1:A:353:GLN:NE2	2.59	0.50
2:B:801:HEM:CBA	5:B:804:1Q6:H18	2.43	0.49
1:A:567:VAL:CG2	5:A:804:1Q6:H17	2.44	0.48
2:A:801:HEM:O2A	5:A:804:1Q6:H8	2.14	0.48
1:B:388:ILE:O	1:B:392:SER:N	2.47	0.47
1:A:605:ASN:ND2	7:A:925:HOH:O	2.47	0.47
1:B:567:VAL:CG2	5:B:804:1Q6:H17	2.47	0.45
1:A:466:THR:N	7:A:959:HOH:O	2.52	0.43
1:A:706:TYR:OH	2:A:801:HEM:O2D	2.36	0.43
1:A:535:GLN:CG	7:A:922:HOH:O	2.66	0.43
1:A:350:THR:OG1	1:A:353:GLN:NE2	2.53	0.42
1:B:445:HIS:CD2	1:B:445:HIS:C	2.95	0.40
1:A:592:GLU:OE1	5:A:804:1Q6:N01	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/422 (96%)	390 (97%)	13 (3%)	0	100	100
1	B	407/422 (96%)	399 (98%)	8 (2%)	0	100	100
All	All	810/844 (96%)	789 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/377 (96%)	353 (97%)	10 (3%)	56	59
1	B	366/377 (97%)	355 (97%)	11 (3%)	53	55
All	All	729/754 (97%)	708 (97%)	21 (3%)	55	57

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	371	ARG
1	A	380	ARG
1	A	489	ASP
1	A	507	GLN
1	A	523	LEU
1	A	527	ASN
1	A	547	ARG
1	A	569	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	645	LYS
1	A	715	VAL
1	B	320	SER
1	B	332	MET
1	B	336	MET
1	B	351	LYS
1	B	353	GLN
1	B	390	SER
1	B	392	SER
1	B	423	LYS
1	B	540	LEU
1	B	547	ARG
1	B	717	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	801	1	49,50,50	4.51	26 (53%)	46,82,82	2.48	17 (36%)
3	H4B	A	802	-	18,18,18	0.90	1 (5%)	24,26,26	1.75	8 (33%)
4	ACT	A	803	-	1,3,3	1.58	0	0,3,3	0.00	-
5	1Q6	A	804	-	25,25,25	0.59	0	33,33,33	1.70	6 (18%)
2	HEM	B	801	1	49,50,50	4.69	24 (48%)	46,82,82	2.75	16 (34%)
3	H4B	B	802	-	18,18,18	1.33	3 (16%)	24,26,26	1.66	2 (8%)
4	ACT	B	803	-	1,3,3	1.27	0	0,3,3	0.00	-
5	1Q6	B	804	-	25,25,25	0.66	0	33,33,33	2.09	9 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	801	1	-	0/14/114/114	0/0/8/8
3	H4B	A	802	-	-	0/8/17/17	0/0/2/2
4	ACT	A	803	-	-	0/0/0/0	0/0/0/0
5	1Q6	A	804	-	-	0/13/13/13	0/2/2/2
2	HEM	B	801	1	-	0/14/114/114	0/0/8/8
3	H4B	B	802	-	-	0/8/17/17	0/0/2/2
4	ACT	B	803	-	-	0/0/0/0	0/0/0/0
5	1Q6	B	804	-	-	0/13/13/13	0/2/2/2

All (54) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	HEM	C2B-C1B	17.22	1.48	1.44
2	B	801	HEM	C2D-C1D	16.81	1.48	1.44
2	A	801	HEM	C2B-C1B	15.89	1.48	1.44
2	A	801	HEM	C2D-C1D	13.84	1.48	1.44
2	B	801	HEM	C3D-C4D	12.32	1.47	1.44
2	A	801	HEM	C3D-C4D	9.61	1.47	1.44
2	A	801	HEM	C4A-C3A	8.24	1.50	1.40
2	A	801	HEM	CHA-C4D	6.98	1.45	1.35
2	B	801	HEM	CHA-C4D	6.05	1.44	1.35
2	A	801	HEM	CHD-C4C	5.82	1.47	1.36
2	B	801	HEM	C4A-C3A	5.72	1.47	1.40
2	B	801	HEM	CHD-C4C	5.37	1.46	1.36
2	A	801	HEM	CHB-C1B	5.34	1.43	1.35
2	A	801	HEM	C4C-NC	5.20	1.45	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	C1C-NC	5.17	1.45	1.38
2	A	801	HEM	CHC-C1C	5.07	1.45	1.36
2	A	801	HEM	C4A-NA	4.91	1.46	1.36
2	B	801	HEM	CHC-C1C	4.90	1.45	1.36
2	A	801	HEM	C1A-NA	4.84	1.46	1.36
2	B	801	HEM	C4A-NA	4.51	1.45	1.36
2	B	801	HEM	C4C-NC	4.30	1.44	1.38
2	B	801	HEM	CHB-C1B	4.08	1.41	1.35
2	B	801	HEM	C1C-NC	4.04	1.43	1.38
2	A	801	HEM	FE-NB	4.01	2.12	1.97
2	B	801	HEM	C3B-C4B	3.86	1.49	1.44
2	B	801	HEM	C1A-NA	3.80	1.44	1.36
2	B	801	HEM	FE-NB	3.73	2.11	1.97
2	A	801	HEM	C3B-C4B	3.64	1.48	1.44
2	A	801	HEM	FE-ND	3.43	2.10	1.97
2	A	801	HEM	CHC-C4B	3.38	1.47	1.39
2	A	801	HEM	C2A-C3A	3.35	1.47	1.37
2	B	801	HEM	C4D-ND	-3.31	1.32	1.39
2	A	801	HEM	CHD-C1D	3.22	1.46	1.39
2	A	801	HEM	C3C-C2C	3.19	1.49	1.43
2	B	801	HEM	FE-ND	3.11	2.09	1.97
2	B	801	HEM	C2A-C3A	3.05	1.46	1.37
2	B	801	HEM	CHD-C1D	2.99	1.46	1.39
2	A	801	HEM	C3C-CAC	2.94	1.49	1.40
3	B	802	H4B	C2-N1	2.91	1.37	1.33
2	A	801	HEM	C3B-CAB	2.87	1.49	1.40
3	B	802	H4B	C7-C6	2.78	1.55	1.52
2	B	801	HEM	CHC-C4B	2.77	1.45	1.39
2	B	801	HEM	C3C-CAC	2.69	1.48	1.40
2	B	801	HEM	C1A-C2A	2.67	1.48	1.43
2	A	801	HEM	C4D-ND	-2.55	1.34	1.39
3	A	802	H4B	C2-N2	2.55	1.36	1.32
2	A	801	HEM	C1A-C2A	2.42	1.47	1.43
2	A	801	HEM	C4A-CHB	2.39	1.46	1.39
2	B	801	HEM	C3B-CAB	2.37	1.47	1.40
2	B	801	HEM	C1B-NB	-2.31	1.34	1.39
3	B	802	H4B	C2-N2	2.30	1.36	1.32
2	B	801	HEM	C3C-C2C	2.13	1.47	1.43
2	A	801	HEM	C2C-C1C	2.13	1.49	1.43
2	A	801	HEM	C3D-C2D	2.06	1.47	1.43

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	HEM	CBA-CAA-C2A	-8.25	98.16	112.69
2	B	801	HEM	C3A-C4A-NA	7.18	114.83	109.41
2	A	801	HEM	CBA-CAA-C2A	-7.07	100.24	112.69
2	B	801	HEM	C1B-NB-C4B	-5.81	99.22	105.16
5	B	804	1Q6	C02-N01-C06	5.80	122.30	118.23
3	B	802	H4B	C4-C4A-C8A	5.78	119.92	114.56
2	A	801	HEM	C1B-NB-C4B	-5.39	99.65	105.16
5	B	804	1Q6	C22-N21-C26	5.00	121.75	118.23
2	A	801	HEM	C3A-C4A-NA	4.92	113.12	109.41
2	B	801	HEM	C4A-CHB-C1B	-4.84	121.10	127.47
2	B	801	HEM	CHD-C1D-ND	-4.72	120.66	124.58
2	B	801	HEM	C4A-C3A-C2A	-4.72	103.71	107.00
5	A	804	1Q6	C22-N21-C26	4.65	121.50	118.23
5	A	804	1Q6	O09-C10-C11	4.53	118.32	109.26
5	B	804	1Q6	C08-O09-C10	-4.46	102.12	112.81
2	B	801	HEM	C4A-NA-C1A	-4.16	101.28	106.76
3	A	802	H4B	C4-C4A-C8A	4.13	118.38	114.56
2	A	801	HEM	C4A-NA-C1A	-3.91	101.62	106.76
2	A	801	HEM	C4A-CHB-C1B	-3.89	122.35	127.47
5	A	804	1Q6	C02-N01-C06	3.88	120.95	118.23
2	A	801	HEM	C1A-CHA-C4D	-3.82	122.45	127.47
2	B	801	HEM	C2D-C1D-ND	3.81	117.44	112.93
2	A	801	HEM	CHD-C1D-ND	-3.73	121.48	124.58
2	A	801	HEM	C2D-C1D-ND	3.61	117.19	112.93
2	A	801	HEM	CBD-CAD-C3D	-3.37	107.03	114.37
2	A	801	HEM	C2A-C1A-NA	3.27	114.28	109.73
2	A	801	HEM	C4D-ND-C1D	-3.18	101.91	105.16
2	B	801	HEM	C4D-ND-C1D	-3.17	101.91	105.16
2	B	801	HEM	C3B-C4B-NB	3.13	116.24	114.00
5	B	804	1Q6	C25-C26-N21	-3.11	119.53	122.99
2	A	801	HEM	CAD-CBD-CGD	-3.10	103.80	113.48
5	A	804	1Q6	C05-C06-N01	-3.06	119.59	122.99
5	B	804	1Q6	O29-C28-C26	2.94	117.95	111.60
5	B	804	1Q6	O09-C10-C11	2.87	115.01	109.26
2	A	801	HEM	CHB-C1B-NB	-2.87	120.36	124.31
3	A	802	H4B	C7-C6-C9	-2.80	109.33	113.66
2	B	801	HEM	C3A-C4A-CHB	-2.80	120.70	126.00
2	B	801	HEM	CBD-CAD-C3D	-2.72	108.44	114.37
3	A	802	H4B	N2-C2-N3	2.71	120.84	117.86
2	A	801	HEM	C4A-C3A-C2A	-2.69	105.13	107.00
2	B	801	HEM	CMB-C2B-C3B	2.68	132.47	126.16
5	A	804	1Q6	C25-C26-N21	-2.66	120.03	122.99
3	A	802	H4B	C4-C4A-N5	2.64	122.94	119.10

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	802	H4B	N8-C8A-N1	2.60	119.63	115.82
5	B	804	1Q6	C28-O29-C11	-2.51	109.59	114.09
2	A	801	HEM	C2A-C1A-CHA	-2.47	121.32	126.00
2	B	801	HEM	C2A-C1A-NA	2.45	113.14	109.73
2	A	801	HEM	C3B-C4B-NB	2.43	115.73	114.00
3	A	802	H4B	C2-N1-C8A	2.34	120.93	117.61
5	B	804	1Q6	C05-C06-N01	-2.32	120.41	122.99
3	A	802	H4B	C6-C7-N8	-2.29	108.70	111.66
2	B	801	HEM	C1A-CHA-C4D	-2.28	124.47	127.47
3	A	802	H4B	C4A-C8A-N8	2.27	122.35	119.23
2	A	801	HEM	CMC-C2C-C3C	2.21	131.38	126.16
5	A	804	1Q6	C08-O09-C10	-2.15	107.66	112.81
2	B	801	HEM	O1D-CGD-CBD	-2.12	115.73	123.03
5	B	804	1Q6	C07-C04-C05	-2.05	117.63	120.93
3	A	802	H4B	C4A-N5-C6	-2.01	115.70	121.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	407/422 (96%)	0.45	34 (8%) 11 12	29, 50, 86, 112	0
1	B	411/422 (97%)	-0.00	11 (2%) 52 57	27, 40, 66, 88	0
All	All	818/844 (96%)	0.22	45 (5%) 24 26	27, 45, 80, 112	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	715	VAL	6.1
1	A	716	TRP	5.5
1	B	300	PHE	5.4
1	A	488	PRO	4.9
1	A	355	PHE	4.6
1	B	321	THR	3.9
1	B	619	ARG	3.6
1	A	352	ASP	3.6
1	A	507	GLN	3.4
1	B	348	VAL	3.3
1	A	486	LYS	3.3
1	A	712	ASN	3.1
1	B	350	THR	3.0
1	A	485	TYR	2.9
1	A	503	GLU	2.9
1	A	390	SER	2.9
1	A	300	PHE	2.8
1	A	321	THR	2.8
1	A	491	SER	2.8
1	A	677	VAL	2.7
1	B	718	GLY	2.7
1	A	678	TRP	2.6
1	A	714	HIS	2.6
1	A	713	THR	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	551	PHE	2.5
1	B	352	ASP	2.4
1	B	620	LYS	2.4
1	A	514	ARG	2.4
1	A	490	GLY	2.4
1	A	506	ILE	2.4
1	A	487	GLN	2.3
1	A	469	LYS	2.3
1	B	680	VAL	2.3
1	A	676	TRP	2.3
1	A	322	LEU	2.3
1	A	392	SER	2.2
1	A	489	ASP	2.2
1	A	351	LYS	2.2
1	A	386	LYS	2.2
1	A	680	VAL	2.1
1	B	353	GLN	2.1
1	A	591	THR	2.1
1	A	511	LYS	2.1
1	B	691	PHE	2.1
1	A	711	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	ACT	A	803	4/4	0.22	6.14	65,65,66,66	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	1Q6	A	804	24/24	0.22	1.33	23,55,84,87	0
4	ACT	B	803	4/4	0.11	1.12	48,48,49,49	0
2	HEM	B	801	43/43	0.15	0.91	26,31,41,48	0
5	1Q6	B	804	24/24	0.17	0.76	27,54,74,76	0
2	HEM	A	801	43/43	0.17	0.47	29,33,39,45	0
3	H4B	A	802	17/17	0.17	0.23	35,36,40,41	0
3	H4B	B	802	17/17	0.15	-0.29	33,35,40,40	0
6	ZN	A	805	1/1	0.08	-0.93	38,38,38,38	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.