



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 02:30 PM GMT

PDB ID : 1K6Z
Title : Crystal Structure of the Yersinia Secretion Chaperone SycE
Authors : Evdokimov, A.G.; Tropea, J.E.; Routzahn, K.M.; Waugh, D.S.
Deposited on : 2001-10-17
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	141	
1	B	141	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	IMD	B	2502	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2161 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type III secretion chaperone SycE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	122	Total	C	N	O	S	Se	0	1	0
			983	625	168	187	1	2			
1	B	121	Total	C	N	O	S	Se	0	1	0
			974	620	166	185	1	2			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	996	ASP	-	FLAG TAG	UNP P31491
A	997	TYR	-	FLAG TAG	UNP P31491
A	998	LYS	-	FLAG TAG	UNP P31491
A	999	ASP	-	FLAG TAG	UNP P31491
A	1000	GLU	-	FLAG TAG	UNP P31491
A	1001	MSE	MET	MODIFIED RESIDUE	UNP P31491
A	1048	MSE	MET	MODIFIED RESIDUE	UNP P31491
A	1109	MSE	MET	MODIFIED RESIDUE	UNP P31491
A	1131	HIS	-	EXPRESSION TAG	UNP P31491
A	1132	HIS	-	EXPRESSION TAG	UNP P31491
A	1133	HIS	-	EXPRESSION TAG	UNP P31491
A	1134	HIS	-	EXPRESSION TAG	UNP P31491
A	1135	HIS	-	EXPRESSION TAG	UNP P31491
A	1136	HIS	-	EXPRESSION TAG	UNP P31491
B	1996	ASP	-	FLAG TAG	UNP P31491
B	1997	TYR	-	FLAG TAG	UNP P31491
B	1998	LYS	-	FLAG TAG	UNP P31491
B	1999	ASP	-	FLAG TAG	UNP P31491
B	2000	GLU	-	FLAG TAG	UNP P31491
B	2001	MSE	MET	MODIFIED RESIDUE	UNP P31491
B	2048	MSE	MET	MODIFIED RESIDUE	UNP P31491
B	2109	MSE	MET	MODIFIED RESIDUE	UNP P31491
B	2131	HIS	-	EXPRESSION TAG	UNP P31491
B	2132	HIS	-	EXPRESSION TAG	UNP P31491
B	2133	HIS	-	EXPRESSION TAG	UNP P31491

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2134	HIS	-	EXPRESSION TAG	UNP P31491
B	2135	HIS	-	EXPRESSION TAG	UNP P31491
B	2136	HIS	-	EXPRESSION TAG	UNP P31491

- Molecule 2 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N 5 3 2	0	0
2	B	1	Total C N 5 3 2	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	100	Total O 100 100	0	0
3	B	94	Total O 94 94	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type III secretion chaperone SycE

Chain A: 



- Molecule 1: Type III secretion chaperone SycE

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.88Å 56.04Å 54.99Å 90.00° 114.04° 90.00°	Depositor
Resolution (Å)	35.00 – 2.00 34.96 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (35.00-2.00) 95.3 (34.96-1.99)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 2.00Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.194 , 0.248 0.198 , 0.234	Depositor DCC
R_{free} test set	1002 reflections (5.61%)	DCC
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 76.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 19757 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2161	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1007	0.98	6/1365 (0.4%)
1	B	0.39	0/998	0.97	3/1353 (0.2%)
All	All	0.39	0/2005	0.98	9/2718 (0.3%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1081	ASP	CB-CG-OD1	6.79	124.41	118.30
1	A	1092[A]	ARG	CD-NE-CZ	6.72	133.00	123.60
1	A	1092[B]	ARG	CD-NE-CZ	6.72	133.00	123.60
1	B	2127	ARG	NE-CZ-NH1	-6.40	117.10	120.30
1	A	1127	ARG	NE-CZ-NH1	-6.22	117.19	120.30
1	B	2092[A]	ARG	NE-CZ-NH2	5.65	123.12	120.30
1	B	2092[B]	ARG	NE-CZ-NH2	5.65	123.12	120.30
1	A	1092[A]	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	A	1092[B]	ARG	NE-CZ-NH1	5.58	123.09	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	983	0	961	15	0
1	B	974	0	953	23	0
2	A	5	0	5	0	0
2	B	5	0	5	1	0
3	A	100	0	0	1	0
3	B	94	0	0	3	0
All	All	2161	0	1924	37	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (37) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1067:ASN:HD21	1:A:1078:LEU:H	1.05	1.00
1:B:2009:THR:HA	1:B:2019:ILE:HD11	1.48	0.92
1:B:2067:ASN:HD21	1:B:2078:LEU:H	1.23	0.83
1:B:2046:ILE:HD11	1:B:2095:LEU:HD13	1.68	0.76
1:B:2010:GLN:O	1:B:2014:GLN:HG3	1.86	0.74
1:A:1067:ASN:ND2	1:A:1078:LEU:H	1.84	0.74
1:A:1010:GLN:O	1:A:1014:GLN:HG3	1.90	0.71
1:A:1046:ILE:HD11	1:A:1095:LEU:HD13	1.71	0.71
1:A:1051:LEU:HB3	3:A:3025:HOH:O	1.92	0.68
1:A:1008:ILE:HG21	1:A:1027:ILE:HD13	1.77	0.66
1:B:2035:ALA:HB1	1:B:2132:HIS:CE1	2.37	0.59
1:B:2031:VAL:CG1	1:B:2127:ARG:HD2	2.33	0.58
1:B:2067:ASN:ND2	1:B:2078:LEU:H	1.99	0.57
1:B:2083:VAL:O	2:B:2502:IMD:H2	2.04	0.56
1:A:1060:LYS:HG2	1:A:1061:GLU:N	2.22	0.55
1:A:1071:GLN:HG2	1:B:2041:HIS:CE1	2.43	0.53
1:A:1059:GLU:HA	1:A:1059:GLU:OE2	2.07	0.53
1:B:2020:PRO:HD3	1:B:2131:HIS:NE2	2.26	0.50
1:A:1024:GLU:OE1	1:A:1024:GLU:HA	2.12	0.50
1:B:2101:ASN:HD22	1:B:2101:ASN:N	2.09	0.50
1:B:2046:ILE:CD1	1:B:2095:LEU:HD13	2.40	0.49
1:B:2016:SER:OG	1:B:2016:SER:O	2.31	0.47
1:B:2019:ILE:O	1:B:2019:ILE:HG22	2.14	0.47
1:A:1011:LEU:O	1:A:1015:LEU:HG	2.15	0.46
1:B:2014:GLN:NE2	3:B:3075:HOH:O	2.50	0.45
1:B:2099:ASP:OD1	1:B:2102:SER:N	2.50	0.45
1:A:1023:ILE:HG22	1:A:1023:ILE:O	2.18	0.44
1:B:2086:HIS:HB2	1:B:2087:PRO:HD2	2.01	0.43
1:B:2007:ALA:HB1	1:B:2103:LEU:HD11	2.00	0.43
1:B:2071:GLN:HE21	1:B:2071:GLN:HB2	1.58	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:2066:HIS:HE1	3:B:3020:HOH:O	2.02	0.42
1:A:1059:GLU:OE1	1:A:1060:LYS:HD2	2.20	0.42
1:B:2062:THR:HG23	3:B:3158:HOH:O	2.21	0.41
1:A:1019:ILE:HA	1:A:1020:PRO:HD3	1.69	0.41
1:B:2029:VAL:CA	1:B:2132:HIS:HB3	2.52	0.40
1:B:2109:MSE:HE2	1:B:2109:MSE:HA	2.04	0.40
1:A:1024:GLU:HB3	1:A:1025:PRO:HD2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/141 (84%)	115 (97%)	4 (3%)	0	100	100
1	B	118/141 (84%)	117 (99%)	1 (1%)	0	100	100
All	All	237/282 (84%)	232 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/129 (88%)	106 (93%)	8 (7%)	21	14
1	B	113/129 (88%)	101 (89%)	12 (11%)	10	5
All	All	227/258 (88%)	207 (91%)	20 (9%)	14	8

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1021	ASP
1	A	1022	THR
1	A	1023	ILE
1	A	1024	GLU
1	A	1051	LEU
1	A	1059	GLU
1	A	1060	LYS
1	A	1097	SER
1	B	2016	SER
1	B	2018	SER
1	B	2019	ILE
1	B	2024	GLU
1	B	2051	LEU
1	B	2056	ASN
1	B	2057	ASN
1	B	2071	GLN
1	B	2099	ASP
1	B	2107	LEU
1	B	2130	SER
1	B	2131	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1013	GLN
1	A	1066	HIS
1	A	1067	ASN
1	A	1112	GLN
1	B	2014	GLN
1	B	2041	HIS
1	B	2066	HIS
1	B	2067	ASN
1	B	2071	GLN
1	B	2093	GLN
1	B	2101	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IMD	A	2501	-	5,5,5	1.19	1 (20%)	5,5,5	1.01	0
2	IMD	B	2502	-	5,5,5	1.13	1 (20%)	5,5,5	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMD	A	2501	-	-	0/0/0/0	0/1/1/1
2	IMD	B	2502	-	-	0/0/0/0	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2501	IMD	C5-N1	2.22	1.40	1.36
2	B	2502	IMD	C5-N1	2.19	1.40	1.36

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	122/141 (86%)	-0.38	1 (0%) 83 84	12, 24, 50, 88	0
1	B	121/141 (85%)	-0.40	2 (1%) 67 67	11, 24, 53, 88	0
All	All	243/282 (86%)	-0.39	3 (1%) 75 76	11, 24, 55, 88	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2132	HIS	2.7
1	B	2022	THR	2.4
1	A	1023	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	IMD	B	2502	5/5	0.22	6.12	75,75,77,77	0
2	IMD	A	2501	5/5	0.10	0.68	9,15,21,28	0

6.5 Other polymers ⓘ

There are no such residues in this entry.