



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 02:26 PM GMT

PDB ID : 3KDN
Title : Crystal structure of Type III Rubisco SP4 mutant complexed with 2-CABP
Authors : Nishitani, Y.; Fujihashi, M.; Doi, T.; Yoshida, S.; Atomi, H.; Imanaka, T.; Miki, K.
Deposited on : 2009-10-23
Resolution : 2.09 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	444	
1	B	444	
1	C	444	
1	D	444	
1	E	444	
1	F	444	
1	G	444	
1	H	444	
1	I	444	
1	J	444	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 36999 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribulose biphosphate carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	0	0
			3349	2145	577	617	10			
1	B	437	Total	C	N	O	S	0	0	0
			3430	2201	588	631	10			
1	C	437	Total	C	N	O	S	0	0	0
			3417	2195	585	627	10			
1	D	437	Total	C	N	O	S	0	0	0
			3421	2196	586	629	10			
1	E	438	Total	C	N	O	S	0	0	0
			3428	2197	586	635	10			
1	F	437	Total	C	N	O	S	0	0	0
			3404	2185	586	623	10			
1	G	437	Total	C	N	O	S	0	0	0
			3433	2204	588	631	10			
1	H	438	Total	C	N	O	S	0	0	0
			3396	2182	584	620	10			
1	I	436	Total	C	N	O	S	0	0	0
			3421	2195	586	630	10			
1	J	437	Total	C	N	O	S	0	0	0
			3428	2199	586	633	10			

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	326	GLU	GLY	ENGINEERED	UNP O93627
A	327	ARG	LYS	ENGINEERED	UNP O93627
A	328	ASP	TRP	ENGINEERED	UNP O93627
A	329	ILE	ASP	ENGINEERED	UNP O93627
A	330	THR	VAL	ENGINEERED	UNP O93627
B	326	GLU	GLY	ENGINEERED	UNP O93627
B	327	ARG	LYS	ENGINEERED	UNP O93627
B	328	ASP	TRP	ENGINEERED	UNP O93627
B	329	ILE	ASP	ENGINEERED	UNP O93627

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	330	THR	VAL	ENGINEERED	UNP O93627
C	326	GLU	GLY	ENGINEERED	UNP O93627
C	327	ARG	LYS	ENGINEERED	UNP O93627
C	328	ASP	TRP	ENGINEERED	UNP O93627
C	329	ILE	ASP	ENGINEERED	UNP O93627
C	330	THR	VAL	ENGINEERED	UNP O93627
D	326	GLU	GLY	ENGINEERED	UNP O93627
D	327	ARG	LYS	ENGINEERED	UNP O93627
D	328	ASP	TRP	ENGINEERED	UNP O93627
D	329	ILE	ASP	ENGINEERED	UNP O93627
D	330	THR	VAL	ENGINEERED	UNP O93627
E	326	GLU	GLY	ENGINEERED	UNP O93627
E	327	ARG	LYS	ENGINEERED	UNP O93627
E	328	ASP	TRP	ENGINEERED	UNP O93627
E	329	ILE	ASP	ENGINEERED	UNP O93627
E	330	THR	VAL	ENGINEERED	UNP O93627
F	326	GLU	GLY	ENGINEERED	UNP O93627
F	327	ARG	LYS	ENGINEERED	UNP O93627
F	328	ASP	TRP	ENGINEERED	UNP O93627
F	329	ILE	ASP	ENGINEERED	UNP O93627
F	330	THR	VAL	ENGINEERED	UNP O93627
G	326	GLU	GLY	ENGINEERED	UNP O93627
G	327	ARG	LYS	ENGINEERED	UNP O93627
G	328	ASP	TRP	ENGINEERED	UNP O93627
G	329	ILE	ASP	ENGINEERED	UNP O93627
G	330	THR	VAL	ENGINEERED	UNP O93627
H	326	GLU	GLY	ENGINEERED	UNP O93627
H	327	ARG	LYS	ENGINEERED	UNP O93627
H	328	ASP	TRP	ENGINEERED	UNP O93627
H	329	ILE	ASP	ENGINEERED	UNP O93627
H	330	THR	VAL	ENGINEERED	UNP O93627
I	326	GLU	GLY	ENGINEERED	UNP O93627
I	327	ARG	LYS	ENGINEERED	UNP O93627
I	328	ASP	TRP	ENGINEERED	UNP O93627
I	329	ILE	ASP	ENGINEERED	UNP O93627
I	330	THR	VAL	ENGINEERED	UNP O93627
J	326	GLU	GLY	ENGINEERED	UNP O93627
J	327	ARG	LYS	ENGINEERED	UNP O93627
J	328	ASP	TRP	ENGINEERED	UNP O93627
J	329	ILE	ASP	ENGINEERED	UNP O93627
J	330	THR	VAL	ENGINEERED	UNP O93627

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Mg	0	0
			1	1		
2	J	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		
2	E	1	Total	Mg	0	0
			1	1		
2	H	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		
2	I	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	F	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-CARBOXYARABINITOL-1,5-DIPHOSPHATE (three-letter code: CAP) (formula: C₆H₁₄O₁₃P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			21	6	13	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	O	P	0	0
			21	6	13	2		
3	C	1	Total	C	O	P	0	0
			21	6	13	2		
3	D	1	Total	C	O	P	0	0
			21	6	13	2		
3	E	1	Total	C	O	P	0	0
			21	6	13	2		
3	F	1	Total	C	O	P	0	0
			21	6	13	2		
3	G	1	Total	C	O	P	0	0
			21	6	13	2		
3	H	1	Total	C	O	P	0	0
			21	6	13	2		
3	I	1	Total	C	O	P	0	0
			21	6	13	2		
3	J	1	Total	C	O	P	0	0
			21	6	13	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	173	Total	O	0	0
			173	173		
4	B	290	Total	O	0	0
			290	290		
4	C	311	Total	O	0	0
			311	311		
4	D	308	Total	O	0	0
			308	308		
4	E	258	Total	O	0	0
			258	258		
4	F	268	Total	O	0	0
			268	268		
4	G	267	Total	O	0	0
			267	267		
4	H	185	Total	O	0	0
			185	185		
4	I	291	Total	O	0	0
			291	291		
4	J	301	Total	O	0	0
			301	301		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ribulose biphosphate carboxylase




• Molecule 1: Ribulose biphosphate carboxylase




• Molecule 1: Ribulose biphosphate carboxylase






• Molecule 1: Ribulose biphosphate carboxylase

Chain D:




• Molecule 1: Ribulose biphosphate carboxylase

Chain E:



• Molecule 1: Ribulose biphosphate carboxylase

Chain F:



• Molecule 1: Ribulose biphosphate carboxylase


Chain G:






• Molecule 1: Ribulose biphosphate carboxylase

Chain H:




• Molecule 1: Ribulose biphosphate carboxylase

Chain I:



• Molecule 1: Ribulose biphosphate carboxylase

Chain J:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	97.47Å 246.24Å 133.08Å 90.00° 104.10° 90.00°	Depositor
Resolution (Å)	42.86 – 2.09 42.86 – 2.09	Depositor EDS
% Data completeness (in resolution range)	96.5 (42.86-2.09) 96.5 (42.86-2.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.213 , 0.253 0.202 , 0.236	Depositor DCC
R_{free} test set	17388 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.556	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 345090 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	36999	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CAP, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/3421	0.40	0/4652
1	B	0.26	0/3505	0.42	0/4753
1	C	0.26	0/3492	0.42	0/4740
1	D	0.27	0/3496	0.44	0/4744
1	E	0.25	0/3503	0.42	0/4753
1	F	0.26	0/3478	0.43	0/4719
1	G	0.25	0/3508	0.43	0/4757
1	H	0.24	0/3471	0.41	0/4713
1	I	0.25	0/3496	0.42	0/4742
1	J	0.26	0/3503	0.43	0/4753
All	All	0.25	0/34873	0.42	0/47326

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3349	0	3204	204	0
1	B	3430	0	3346	68	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3417	0	3325	81	0
1	D	3421	0	3329	75	0
1	E	3428	0	3319	78	0
1	F	3404	0	3312	70	0
1	G	3433	0	3355	60	0
1	H	3396	0	3282	118	0
1	I	3421	0	3333	56	0
1	J	3428	0	3335	82	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
3	A	21	0	7	3	0
3	B	21	0	7	0	0
3	C	21	0	8	1	0
3	D	21	0	8	0	0
3	E	21	0	7	0	0
3	F	21	0	8	0	0
3	G	21	0	8	0	0
3	H	21	0	8	1	0
3	I	21	0	7	0	0
3	J	21	0	7	0	0
4	A	173	0	0	9	0
4	B	290	0	0	5	0
4	C	311	0	0	7	0
4	D	308	0	0	6	0
4	E	258	0	0	9	0
4	F	268	0	0	10	0
4	G	267	0	0	4	0
4	H	185	0	0	9	0
4	I	291	0	0	8	0
4	J	301	0	0	9	0
All	All	36999	0	33215	856	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 856 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:149:MET:HE3	1:A:250:LYS:CD	1.57	1.31
1:A:38:THR:HG23	1:A:41:GLN:OE1	1.53	1.07
1:A:149:MET:HE3	1:A:250:LYS:HD2	1.07	1.04
1:A:438:TRP:O	1:A:441:VAL:HG12	1.58	1.03
1:A:424:TYR:CE2	1:A:428:HIS:CE1	2.46	1.03

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/444 (98%)	394 (91%)	37 (8%)	3 (1%)	30	23
1	B	434/444 (98%)	418 (96%)	16 (4%)	0	100	100
1	C	434/444 (98%)	419 (96%)	15 (4%)	0	100	100
1	D	434/444 (98%)	422 (97%)	12 (3%)	0	100	100
1	E	435/444 (98%)	418 (96%)	16 (4%)	1 (0%)	56	57
1	F	434/444 (98%)	419 (96%)	14 (3%)	1 (0%)	56	57
1	G	434/444 (98%)	419 (96%)	15 (4%)	0	100	100
1	H	435/444 (98%)	399 (92%)	32 (7%)	4 (1%)	25	17
1	I	433/444 (98%)	418 (96%)	15 (4%)	0	100	100
1	J	434/444 (98%)	420 (97%)	13 (3%)	1 (0%)	56	57
All	All	4341/4440 (98%)	4146 (96%)	185 (4%)	10 (0%)	56	57

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	426	LYS
1	E	284	MET
1	J	284	MET
1	A	117	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	325	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	323/356 (91%)	316 (98%)	7 (2%)	64	68
1	B	341/356 (96%)	336 (98%)	5 (2%)	76	81
1	C	338/356 (95%)	332 (98%)	6 (2%)	71	75
1	D	339/356 (95%)	333 (98%)	6 (2%)	71	75
1	E	339/356 (95%)	334 (98%)	5 (2%)	76	81
1	F	335/356 (94%)	328 (98%)	7 (2%)	66	70
1	G	342/356 (96%)	336 (98%)	6 (2%)	71	75
1	H	330/356 (93%)	323 (98%)	7 (2%)	66	70
1	I	340/356 (96%)	338 (99%)	2 (1%)	92	95
1	J	341/356 (96%)	331 (97%)	10 (3%)	55	57
All	All	3368/3560 (95%)	3307 (98%)	61 (2%)	71	75

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	328	ASP
1	F	332	GLN
1	J	329	ILE
1	E	363	PHE
1	F	87	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	354	GLN
1	E	354	GLN
1	I	376	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	440	HIS
1	E	312	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	A	189	1,2	11,11,12	5.39	2 (18%)	10,12,14	1.76	4 (40%)
1	KCX	B	189	1,2	11,11,12	5.33	2 (18%)	10,12,14	1.53	3 (30%)
1	KCX	C	189	1,2	11,11,12	5.32	2 (18%)	10,12,14	1.56	4 (40%)
1	KCX	D	189	1,2	11,11,12	5.31	2 (18%)	10,12,14	1.82	3 (30%)
1	KCX	E	189	1,2	11,11,12	5.38	2 (18%)	10,12,14	1.65	3 (30%)
1	KCX	F	189	1,2	11,11,12	5.36	2 (18%)	10,12,14	1.69	3 (30%)
1	KCX	G	189	1,2	11,11,12	5.33	2 (18%)	10,12,14	1.60	2 (20%)
1	KCX	H	189	1,2	11,11,12	5.37	2 (18%)	10,12,14	1.55	2 (20%)
1	KCX	I	189	1,2	11,11,12	5.51	2 (18%)	10,12,14	1.71	3 (30%)
1	KCX	J	189	1,2	11,11,12	5.30	2 (18%)	10,12,14	1.58	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	B	189	1,2	-	0/8/10/12	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	C	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	D	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	E	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	F	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	G	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	H	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	I	189	1,2	-	0/8/10/12	0/0/0/0
1	KCX	J	189	1,2	-	0/8/10/12	0/0/0/0

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	189	KCX	O-C	17.89	1.23	1.11
1	A	189	KCX	O-C	17.55	1.23	1.11
1	E	189	KCX	O-C	17.53	1.23	1.11
1	H	189	KCX	O-C	17.46	1.23	1.11
1	F	189	KCX	O-C	17.40	1.23	1.11

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	189	KCX	OQ2-CX-NZ	3.30	120.43	116.33
1	D	189	KCX	OQ1-CX-NZ	-3.17	119.67	124.94
1	A	189	KCX	C-CA-N	-3.12	110.71	113.83
1	I	189	KCX	C-CA-N	-3.12	110.72	113.83
1	G	189	KCX	OQ2-CX-NZ	3.04	120.11	116.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 20 ligands modelled in this entry, 10 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CAP	A	600	2	20,20,20	0.77	0	31,31,31	0.91	1 (3%)
3	CAP	B	600	2	20,20,20	0.72	0	31,31,31	0.90	1 (3%)
3	CAP	C	600	2	20,20,20	0.75	0	31,31,31	0.83	1 (3%)
3	CAP	D	600	2	20,20,20	0.72	0	31,31,31	0.92	1 (3%)
3	CAP	E	600	2	20,20,20	0.72	0	31,31,31	0.89	1 (3%)
3	CAP	F	600	2	20,20,20	0.71	0	31,31,31	0.86	1 (3%)
3	CAP	G	600	2	20,20,20	0.70	0	31,31,31	0.89	1 (3%)
3	CAP	H	600	2	20,20,20	0.71	0	31,31,31	0.93	1 (3%)
3	CAP	I	600	2	20,20,20	0.72	0	31,31,31	0.84	1 (3%)
3	CAP	J	600	2	20,20,20	0.77	0	31,31,31	0.82	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CAP	A	600	2	-	0/29/29/29	0/0/0/0
3	CAP	B	600	2	-	0/29/29/29	0/0/0/0
3	CAP	C	600	2	-	0/29/29/29	0/0/0/0
3	CAP	D	600	2	-	0/29/29/29	0/0/0/0
3	CAP	E	600	2	-	0/29/29/29	0/0/0/0
3	CAP	F	600	2	-	0/29/29/29	0/0/0/0
3	CAP	G	600	2	-	0/29/29/29	0/0/0/0
3	CAP	H	600	2	-	0/29/29/29	0/0/0/0
3	CAP	I	600	2	-	0/29/29/29	0/0/0/0
3	CAP	J	600	2	-	0/29/29/29	0/0/0/0

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	600	CAP	O7-C-C2	3.02	120.17	113.58
3	A	600	CAP	O7-C-C2	2.76	119.59	113.58
3	B	600	CAP	O7-C-C2	2.76	119.59	113.58
3	G	600	CAP	O7-C-C2	2.75	119.56	113.58
3	D	600	CAP	O7-C-C2	2.74	119.55	113.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	437/444 (98%)	0.81	50 (11%) 6 6	22, 36, 52, 58	0
1	B	437/444 (98%)	-0.20	4 (0%) 81 85	13, 22, 34, 37	0
1	C	437/444 (98%)	-0.10	2 (0%) 88 91	12, 21, 40, 45	0
1	D	437/444 (98%)	-0.17	4 (0%) 81 85	11, 20, 32, 39	0
1	E	438/444 (98%)	-0.08	9 (2%) 60 65	15, 24, 47, 51	0
1	F	437/444 (98%)	0.04	9 (2%) 60 65	12, 24, 38, 46	0
1	G	437/444 (98%)	-0.13	3 (0%) 84 88	13, 24, 36, 41	0
1	H	438/444 (98%)	0.50	31 (7%) 16 17	22, 32, 57, 61	0
1	I	436/444 (98%)	-0.19	1 (0%) 93 95	15, 23, 34, 40	0
1	J	437/444 (98%)	-0.15	6 (1%) 72 76	11, 20, 41, 47	0
All	All	4371/4440 (98%)	0.03	119 (2%) 50 57	11, 24, 42, 61	0

The worst 5 of 119 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	59	TYR	5.0
1	E	7	THR	4.6
1	A	59	TYR	4.4
1	H	7	THR	4.2
1	A	414	ALA	4.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	KCX	F	189	12/13	0.17	1.19	17,18,20,20	0
1	KCX	A	189	12/13	0.18	0.76	29,30,31,31	0
1	KCX	B	189	12/13	0.12	0.48	18,19,19,20	0
1	KCX	D	189	12/13	0.13	0.45	11,14,15,15	0
1	KCX	J	189	12/13	0.13	0.45	18,18,19,20	0
1	KCX	G	189	12/13	0.13	0.41	16,17,17,17	0
1	KCX	I	189	12/13	0.11	0.09	17,18,18,18	0
1	KCX	H	189	12/13	0.13	-0.12	30,30,31,31	0
1	KCX	E	189	12/13	0.11	-0.25	21,22,23,23	0
1	KCX	C	189	12/13	0.11	-0.27	14,16,17,17	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CAP	H	600	21/21	0.17	0.06	31,34,36,37	0
3	CAP	J	600	21/21	0.12	0.03	17,21,22,22	0
3	CAP	I	600	21/21	0.12	-0.22	16,18,18,19	0
3	CAP	E	600	21/21	0.11	-0.24	21,24,26,26	0
3	CAP	A	600	21/21	0.16	-0.31	27,33,34,35	0
3	CAP	G	600	21/21	0.11	-0.34	17,17,18,18	0
3	CAP	C	600	21/21	0.12	-0.51	19,21,22,23	0
3	CAP	F	600	21/21	0.12	-0.63	15,17,17,18	0
3	CAP	B	600	21/21	0.09	-0.63	17,20,21,22	0
3	CAP	D	600	21/21	0.10	-0.68	13,14,15,16	0
2	MG	H	500	1/1	0.11	-1.55	29,29,29,29	0
2	MG	J	500	1/1	0.10	-1.80	16,16,16,16	0
2	MG	B	500	1/1	0.07	-2.09	18,18,18,18	0
2	MG	E	500	1/1	0.09	-2.37	21,21,21,21	0
2	MG	F	500	1/1	0.09	-2.49	19,19,19,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	I	500	1/1	0.08	-3.05	13,13,13,13	0
2	MG	C	500	1/1	0.07	-3.56	16,16,16,16	0
2	MG	G	500	1/1	0.05	-3.77	16,16,16,16	0
2	MG	A	500	1/1	0.11	-5.60	29,29,29,29	0
2	MG	D	500	1/1	0.06	-6.25	11,11,11,11	0

6.5 Other polymers ⓘ

There are no such residues in this entry.