



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 05:38 AM GMT

PDB ID : 1KF9
Title : PHAGE DISPLAY DERIVED VARIANT OF HUMAN GROWTH HORMONE COMPLEXED WITH TWO COPIES OF THE EXTRACELLULAR DOMAIN OF ITS RECEPTOR
Authors : Schiffer, C.A.; Ultsch, M.; Walsh, S.; Somers, W.; De Vos, A.M.; Kossiakoff, A.A.
Deposited on : 2001-11-19
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	191	
1	D	191	
2	B	238	
2	C	238	
2	E	238	
2	F	238	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8678 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHAGE DISPLAY DERIVED VARIANT HUMAN GROWTH HORMONE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	160	Total	C	N	O	S	0	0	0
			1284	825	211	242	6			
1	D	158	Total	C	N	O	S	0	0	0
			1264	811	208	239	6			

- Molecule 2 is a protein called EXTRACELLULAR DOMAIN HUMAN GROWTH HORMONE RECEPTOR (1-238).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	193	Total	C	N	O	S	0	0	0
			1577	1011	258	298	10			
2	C	175	Total	C	N	O	S	0	0	0
			1419	916	229	265	9			
2	E	193	Total	C	N	O	S	0	0	0
			1576	1011	258	298	9			
2	F	184	Total	C	N	O	S	0	0	0
			1484	952	239	284	9			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	11	Total	O	0	0
			11	11		
3	B	13	Total	O	0	0
			13	13		
3	C	28	Total	O	0	0
			28	28		
3	D	3	Total	O	0	0
			3	3		
3	E	12	Total	O	0	0
			12	12		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	7	Total	O	0	0
			7	7		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: PHAGE DISPLAY DERIVED VARIANT HUMAN GROWTH HORMONE




• Molecule 1: PHAGE DISPLAY DERIVED VARIANT HUMAN GROWTH HORMONE




• Molecule 2: EXTRACELLULAR DOMAIN HUMAN GROWTH HORMONE RECEPTOR (1-238)






• Molecule 2: EXTRACELLULAR DOMAIN HUMAN GROWTH HORMONE RECEPTOR (1-238)

Chain C:




• Molecule 2: EXTRACELLULAR DOMAIN HUMAN GROWTH HORMONE RECEPTOR (1-238)

Chain E:



• Molecule 2: EXTRACELLULAR DOMAIN HUMAN GROWTH HORMONE RECEPTOR (1-238)

Chain F:



P1698	V1699	Y1700	S1701	L1702	K1703	V1704		R1711	V1712	R1713	S1714	K1715	Q1716	R1717	N1718	S1719		Y1722	G1723	E1724	F1725		V1728	L1729		T1732	L1733	P1734	GLN	MET	SER	GLN
-------	-------	-------	-------	-------	-------	-------	--	-------	-------	-------	-------	-------	-------	-------	-------	-------	--	-------	-------	-------	-------	--	-------	-------	--	-------	-------	-------	-----	-----	-----	-----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.29Å 111.94Å 95.29Å 90.00° 90.06° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60	Depositor
% Data completeness (in resolution range)	90.5 (20.00-2.60)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.234 , 0.326	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8678	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/1312	0.84	2/1781 (0.1%)
1	D	0.71	0/1289	0.84	2/1747 (0.1%)
2	B	0.70	0/1622	0.87	0/2208
2	C	0.67	0/1459	0.86	1/1987 (0.1%)
2	E	0.69	0/1621	0.86	1/2208 (0.0%)
2	F	0.65	0/1525	0.84	0/2078
All	All	0.68	0/8828	0.85	6/12009 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	THR	N-CA-C	-9.03	86.61	111.00
1	D	1107	ASP	CB-CG-OD1	6.76	124.39	118.30
2	C	668	GLY	N-CA-C	-6.00	98.09	113.10
2	E	1324	SER	N-CA-C	-5.23	96.87	111.00
1	A	1	PHE	C-N-CD	5.20	139.32	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1284	0	1257	102	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1264	0	1241	88	0
2	B	1577	0	1506	91	0
2	C	1419	0	1338	99	0
2	E	1576	0	1508	104	0
2	F	1484	0	1395	115	0
3	A	11	0	0	2	0
3	B	13	0	0	1	0
3	C	28	0	0	2	0
3	D	3	0	0	0	0
3	E	12	0	0	1	0
3	F	7	0	0	0	0
All	All	8678	0	8245	550	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 33.

The worst 5 of 550 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:104:GLY:CA	2:C:666:GLN:HB3	1.65	1.26
2:E:1346:LEU:HB2	2:F:1701:SER:HB3	1.24	1.17
1:A:104:GLY:HA2	2:C:666:GLN:HB3	1.25	1.14
2:F:1581:LYS:HZ3	2:F:1581:LYS:HB2	1.08	1.09
1:A:104:GLY:HA3	2:C:666:GLN:HB3	1.41	1.02

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	154/191 (81%)	129 (84%)	19 (12%)	6 (4%)	5 6
1	D	150/191 (78%)	129 (86%)	17 (11%)	4 (3%)	8 13
2	B	189/238 (79%)	168 (89%)	20 (11%)	1 (0%)	38 67
2	C	167/238 (70%)	144 (86%)	21 (13%)	2 (1%)	19 39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	189/238 (79%)	171 (90%)	16 (8%)	2 (1%)	21	42
2	F	178/238 (75%)	155 (87%)	20 (11%)	3 (2%)	14	26
All	All	1027/1334 (77%)	896 (87%)	113 (11%)	18 (2%)	13	25

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	SER
1	A	107	ASP
2	B	274	GLN
1	D	1002	PRO
2	E	1274	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	143/175 (82%)	112 (78%)	31 (22%)	1	2
1	D	142/175 (81%)	112 (79%)	30 (21%)	1	2
2	B	177/218 (81%)	130 (73%)	47 (27%)	1	1
2	C	157/218 (72%)	113 (72%)	44 (28%)	0	1
2	E	177/218 (81%)	134 (76%)	43 (24%)	1	2
2	F	165/218 (76%)	121 (73%)	44 (27%)	1	1
All	All	961/1222 (79%)	722 (75%)	239 (25%)	1	2

5 of 239 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	672	LEU
1	D	1055	SER
2	F	1658	GLU
2	C	692	ILE
2	C	729	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such

sidechains are listed below:

Mol	Chain	Res	Type
2	C	716	GLN
1	D	1122	GLN
2	F	1643	ASN
1	D	1068	GLN
1	D	1099	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.