



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 12:57 AM GMT

PDB ID : 3KF2
Title : The HCV NS3/NS4A protease apo structure
Authors : Lindberg, J.D.; Nystrom, S.; Cummings, M.D.
Deposited on : 2009-10-27
Resolution : 2.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	200	
1	B	200	
2	C	23	
2	D	23	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2782 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	154	Total	C	N	O	S	0	0	0
			1142	713	207	214	8			
1	B	181	Total	C	N	O	S	0	0	0
			1341	836	244	252	9			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	EXPRESSION TAG	UNP B1PBR5
A	-9	ALA	-	EXPRESSION TAG	UNP B1PBR5
A	-8	SER	-	EXPRESSION TAG	UNP B1PBR5
A	-7	MET	-	EXPRESSION TAG	UNP B1PBR5
A	-6	THR	-	EXPRESSION TAG	UNP B1PBR5
A	-5	GLY	-	EXPRESSION TAG	UNP B1PBR5
A	-4	GLY	-	EXPRESSION TAG	UNP B1PBR5
A	-3	GLN	-	EXPRESSION TAG	UNP B1PBR5
A	-2	GLN	-	EXPRESSION TAG	UNP B1PBR5
A	-1	MET	-	EXPRESSION TAG	UNP B1PBR5
A	0	GLY	-	EXPRESSION TAG	UNP B1PBR5
A	176	GLU	GLY	ENGINEERED	UNP B1PBR5
A	182	GLY	-	EXPRESSION TAG	UNP B1PBR5
A	183	SER	-	EXPRESSION TAG	UNP B1PBR5
A	184	HIS	-	EXPRESSION TAG	UNP B1PBR5
A	185	HIS	-	EXPRESSION TAG	UNP B1PBR5
A	186	HIS	-	EXPRESSION TAG	UNP B1PBR5
A	187	HIS	-	EXPRESSION TAG	UNP B1PBR5
A	188	HIS	-	EXPRESSION TAG	UNP B1PBR5
A	189	HIS	-	EXPRESSION TAG	UNP B1PBR5
B	-10	MET	-	EXPRESSION TAG	UNP B1PBR5
B	-9	ALA	-	EXPRESSION TAG	UNP B1PBR5
B	-8	SER	-	EXPRESSION TAG	UNP B1PBR5
B	-7	MET	-	EXPRESSION TAG	UNP B1PBR5
B	-6	THR	-	EXPRESSION TAG	UNP B1PBR5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLY	-	EXPRESSION TAG	UNP B1PBR5
B	-4	GLY	-	EXPRESSION TAG	UNP B1PBR5
B	-3	GLN	-	EXPRESSION TAG	UNP B1PBR5
B	-2	GLN	-	EXPRESSION TAG	UNP B1PBR5
B	-1	MET	-	EXPRESSION TAG	UNP B1PBR5
B	0	GLY	-	EXPRESSION TAG	UNP B1PBR5
B	176	GLU	GLY	ENGINEERED	UNP B1PBR5
B	182	GLY	-	EXPRESSION TAG	UNP B1PBR5
B	183	SER	-	EXPRESSION TAG	UNP B1PBR5
B	184	HIS	-	EXPRESSION TAG	UNP B1PBR5
B	185	HIS	-	EXPRESSION TAG	UNP B1PBR5
B	186	HIS	-	EXPRESSION TAG	UNP B1PBR5
B	187	HIS	-	EXPRESSION TAG	UNP B1PBR5
B	188	HIS	-	EXPRESSION TAG	UNP B1PBR5
B	189	HIS	-	EXPRESSION TAG	UNP B1PBR5

- Molecule 2 is a protein called 19-mer peptide from Genome polyprotein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	16	Total	C	N	O	0	0	0
			108	70	20	18			
2	D	21	Total	C	N	O	0	0	0
			149	99	27	23			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	LYS	-	EXPRESSION TAG	UNP Q6GYR8
C	20	LYS	-	EXPRESSION TAG	UNP Q6GYR8
C	40	LYS	-	EXPRESSION TAG	UNP Q6GYR8
C	41	LYS	-	EXPRESSION TAG	UNP Q6GYR8
D	19	LYS	-	EXPRESSION TAG	UNP Q6GYR8
D	20	LYS	-	EXPRESSION TAG	UNP Q6GYR8
D	40	LYS	-	EXPRESSION TAG	UNP Q6GYR8
D	41	LYS	-	EXPRESSION TAG	UNP Q6GYR8

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total 11	O 11	0	0
4	C	5	Total 5	O 5	0	0
4	B	20	Total 20	O 20	0	0
4	D	4	Total 4	O 4	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	224.92Å 224.92Å 75.55Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	112.46 – 2.50 112.46 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (112.46-2.50) 99.5 (112.46-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.52Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.205 , 0.235 0.205 , 0.234	Depositor DCC
R_{free} test set	1285 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 17.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 25152 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2782	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/1166	0.76	0/1591
1	B	0.62	0/1367	0.76	0/1864
2	C	0.63	0/108	0.56	0/145
2	D	0.68	0/150	0.72	0/201
All	All	0.61	0/2791	0.75	0/3801

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1142	0	1145	14	0
1	B	1341	0	1357	17	0
2	C	108	0	125	4	0
2	D	149	0	180	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	11	0	0	0	0
4	B	20	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	5	0	0	0	0
4	D	4	0	0	0	0
All	All	2782	0	2807	30	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (30) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:130:ARG:HD3	4:B:198:HOH:O	1.64	0.97
1:A:99:CYS:HA	1:B:127:LEU:HD12	1.67	0.75
1:B:159:CYS:HB3	1:B:164:ALA:HA	1.69	0.73
1:A:94:LEU:HD13	1:A:150:ALA:HB2	1.74	0.70
1:A:63:THR:HG1	2:C:22:SER:HG	1.42	0.65
1:B:82:LEU:HD11	1:B:175:LEU:HD12	1.79	0.63
1:B:95:THR:OG1	1:B:149:HIS:HD2	1.87	0.57
1:A:28:GLN:HE22	1:A:90:GLY:HA2	1.72	0.55
1:A:158:VAL:HB	1:A:166:ALA:HB3	1.89	0.55
1:B:42:THR:OG1	1:B:109:ARG:NH2	2.40	0.54
1:B:95:THR:OG1	1:B:149:HIS:CD2	2.61	0.54
1:B:104:LEU:HB2	1:B:116:VAL:HG22	1.90	0.54
1:B:77:ASN:ND2	1:B:80:GLN:HG3	2.22	0.53
1:B:11:ARG:HH12	1:B:25:ASP:CG	2.13	0.52
1:B:142:PRO:HB2	1:B:144:LEU:HD13	1.91	0.52
1:B:102:SER:HB2	1:B:118:ARG:HB3	1.91	0.52
1:B:121:ASP:O	1:B:171:PRO:HG3	2.12	0.50
1:A:36:VAL:HG13	2:C:23:VAL:CG1	2.42	0.49
1:A:149:HIS:HE1	4:B:193:HOH:O	1.96	0.49
1:A:33:VAL:HB	2:C:29:ILE:HB	1.97	0.47
1:A:28:GLN:NE2	1:A:90:GLY:HA2	2.32	0.44
1:B:77:ASN:ND2	1:B:80:GLN:CG	2.80	0.44
1:A:114:ILE:HA	1:A:115:PRO:HD3	1.81	0.44
1:B:33:VAL:HB	2:D:29:ILE:HB	2.00	0.44
1:A:47:CYS:HA	1:A:51:VAL:O	2.20	0.42
1:B:139:SER:HB2	4:B:194:HOH:O	2.20	0.42
1:B:44:LEU:HD21	1:B:109:ARG:HB3	2.02	0.41
1:A:102:SER:CB	1:A:118:ARG:HB3	2.51	0.41
1:A:36:VAL:HG13	2:C:23:VAL:HG13	2.03	0.41
1:A:118:ARG:HH21	1:A:121:ASP:HA	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	152/200 (76%)	149 (98%)	3 (2%)	0	100	100
1	B	179/200 (90%)	175 (98%)	4 (2%)	0	100	100
2	C	14/23 (61%)	14 (100%)	0	0	100	100
2	D	19/23 (83%)	19 (100%)	0	0	100	100
All	All	364/446 (82%)	357 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/161 (78%)	107 (85%)	19 (15%)	4	7
1	B	147/161 (91%)	133 (90%)	14 (10%)	12	22
2	C	12/19 (63%)	12 (100%)	0	100	100
2	D	17/19 (90%)	17 (100%)	0	100	100
All	All	302/360 (84%)	269 (89%)	33 (11%)	9	16

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	VAL
1	A	34	GLN
1	A	61	THR
1	A	72	ILE
1	A	82	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	95	THR
1	A	109	ARG
1	A	113	VAL
1	A	114	ILE
1	A	116	VAL
1	A	117	ARG
1	A	139	SER
1	A	143	LEU
1	A	144	LEU
1	A	153	LEU
1	A	165	LYS
1	A	168	ASP
1	A	176	GLU
1	A	181	SER
1	B	13	LEU
1	B	36	VAL
1	B	37	SER
1	B	42	THR
1	B	61	THR
1	B	62	ARG
1	B	92	ARG
1	B	99	CYS
1	B	109	ARG
1	B	114	ILE
1	B	117	ARG
1	B	136	LYS
1	B	139	SER
1	B	144	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	49	ASN
1	A	149	HIS
1	B	27	ASN
1	B	149	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	154/200 (77%)	0.41	4 (2%) 53 55	22, 34, 50, 68	0
1	B	181/200 (90%)	0.29	1 (0%) 86 88	19, 31, 45, 53	0
2	C	16/23 (69%)	0.08	1 (6%) 19 19	20, 27, 38, 44	0
2	D	21/23 (91%)	0.20	0 100 100	21, 27, 38, 47	0
All	All	372/446 (83%)	0.33	6 (1%) 68 71	19, 32, 47, 68	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	180	ARG	3.1
1	B	121	ASP	2.7
2	C	36	ALA	2.6
1	A	28	GLN	2.4
1	A	79	ASP	2.1
1	A	176	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ZN	A	901	1/1	0.18	1.43	30,30,30,30	0
3	ZN	B	902	1/1	0.16	-0.19	38,38,38,38	0

6.5 Other polymers

There are no such residues in this entry.