



Full wwPDB X-ray Structure Validation Report

Oct 20, 2014 – 06:51 PM EDT

PDB ID : 4KG5
Title : Crystal Structure of AmpC beta-lactamase N152G Mutant in Complex with Cefotaxime
Authors : Docter, B.E.; Baggett, V.L.; Powers, R.A.; Wallar, B.J.
Deposited on : 2013-04-28
Resolution : 2.11 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable24103
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable24103

1 Overall quality at a glance

The reported resolution of this entry is 2.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3409 (2.14-2.10)
Clashscore	79885	4090 (2.14-2.10)
Ramachandran outliers	78287	4048 (2.14-2.10)
Sidechain outliers	78261	4049 (2.14-2.10)
RSRZ outliers	66119	3410 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	358	
1	B	358	
1	C	358	
1	D	358	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	PO4	A	401	-	X
2	PO4	A	402	-	X
2	PO4	A	403	-	X
2	PO4	B	401	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
2	PO4	D	401	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11765 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	358	Total	C	N	O	S	0	1	0
			2755	1776	468	505	6			
1	A	358	Total	C	N	O	S	0	5	0
			2803	1801	479	517	6			
1	C	358	Total	C	N	O	S	0	3	0
			2786	1793	472	515	6			
1	D	358	Total	C	N	O	S	0	3	0
			2790	1793	475	516	6			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	152	GLY	ASN	ENGINEERED MUTATION	UNP P00811
A	152	GLY	ASN	ENGINEERED MUTATION	UNP P00811
C	152	GLY	ASN	ENGINEERED MUTATION	UNP P00811
D	152	GLY	ASN	ENGINEERED MUTATION	UNP P00811

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is (2R)-2-[(1R)-1-{[(2Z)-2-(2-AMINO-1,3-THIAZOL-4-YL)-2-(METHOXYIMINO)ACETYL]AMINO}-2-OXOETHYL]-5-METHYLIDENE-5,6-DIHYDRO-2H-1,3-THIAZINE-4-CARBOXYLICACID (three-letter code: PCZ) (formula: C₁₄H₁₅N₅O₅S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			26	14	5	5	2		
3	A	1	Total	C	N	O	S	0	0
			26	14	5	5	2		
3	C	1	Total	C	N	O	S	0	0
			26	14	5	5	2		
3	D	1	Total	C	N	O	S	0	0
			26	14	5	5	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	113	Total	O	0	1
			114	114		
4	A	131	Total	O	0	1
			132	132		
4	C	120	Total	O	0	3
			123	123		
4	D	132	Total	O	0	1
			133	133		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-lactamase




- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	163.37Å 80.35Å 129.79Å 90.00° 103.99° 90.00°	Depositor
Resolution (Å)	28.23 – 2.11 28.21 – 2.11	Depositor EDS
% Data completeness (in resolution range)	96.8 (28.23-2.11) 96.9 (28.21-2.11)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.12Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.179 , 0.232 0.186 , 0.234	Depositor DCC
R_{free} test set	4580 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 91499 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11765	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 50.21 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.6803e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PCZ, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	0/2883	0.88	0/3947
1	B	0.85	1/2835 (0.0%)	0.81	0/3883
1	C	0.83	0/2866	0.87	1/3923 (0.0%)
1	D	0.88	0/2870	0.89	1/3929 (0.0%)
All	All	0.87	1/11454 (0.0%)	0.86	2/15682 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	324	SER	CB-OG	-5.85	1.34	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	47	ASP	CB-CG-OD1	6.95	124.56	118.30
1	D	275	ASP	CB-CG-OD1	6.11	123.79	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2803	0	2725	15	0
1	B	2755	0	2680	13	0
1	C	2786	0	2712	25	0
1	D	2790	0	2710	10	1
2	A	15	0	0	0	0
2	B	5	0	0	0	0
2	D	5	0	0	0	0
3	A	26	0	13	2	0
3	B	26	0	13	0	0
3	C	26	0	13	2	0
3	D	26	0	13	2	0
4	A	132	0	0	1	0
4	B	114	0	0	0	0
4	C	123	0	0	5	1
4	D	133	0	0	1	0
All	All	11765	0	10879	64	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (64) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:253[B]:GLN:NE2	4:C:592:HOH:O	1.92	1.02
1:C:355[B]:GLN:NE2	4:C:597:HOH:O	1.99	0.94
1:C:296:ARG:HD3	4:C:513:HOH:O	1.92	0.68
1:C:95:GLU:OE2	1:C:95:GLU:N	2.32	0.63
1:D:238:LEU:HD23	1:D:330:PRO:HA	1.83	0.59
1:A:304:PRO:HG2	1:D:303:PRO:HB3	1.85	0.59
1:C:243:ILE:HD12	1:C:252:ILE:HD12	1.86	0.58
1:B:26:PRO:HB3	1:B:48:ILE:HD11	1.86	0.58
3:D:402:PCZ:H15	3:D:402:PCZ:O91	2.04	0.58
1:D:344:TYR:CZ	1:D:349:ARG:HG2	2.38	0.58
1:D:243:ILE:CD1	1:D:252:ILE:HD12	2.34	0.57
3:C:401:PCZ:H15	3:C:401:PCZ:O91	2.04	0.57
1:C:5:PRO:HB2	1:C:7:GLN:NE2	2.20	0.57
1:A:172:GLN:O	1:A:176:THR:HG23	2.03	0.56
3:A:404:PCZ:H522	3:A:404:PCZ:O12	2.07	0.54
1:C:33:ILE:HD12	1:C:33:ILE:N	2.22	0.53
1:A:262:THR:HG21	1:A:296[A]:ARG:CB	2.39	0.53
1:C:344:TYR:CE2	1:C:349:ARG:HG2	2.44	0.52
1:C:265:MET:CE	1:C:272:GLU:HB3	2.41	0.51
1:A:164:LYS:HB2	1:A:165:PRO:HD3	1.93	0.51
1:C:197:LYS:HE2	4:C:543:HOH:O	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:164:LYS:HB2	1:B:165:PRO:HD3	1.93	0.50
1:B:304:PRO:HD2	1:C:304:PRO:HD2	1.94	0.50
1:A:296[A]:ARG:NH2	4:A:595:HOH:O	2.44	0.50
1:D:172:GLN:NE2	4:D:608:HOH:O	2.41	0.50
1:B:304:PRO:HG2	1:C:303:PRO:HB3	1.94	0.49
1:C:239:LYS:HD2	1:C:333:GLU:OE2	2.13	0.48
1:C:265:MET:HE3	1:C:272:GLU:HB3	1.96	0.48
1:C:244:ASN:HD22	1:C:244:ASN:N	2.12	0.47
1:A:180:GLN:HB2	1:A:181:PRO:HD3	1.96	0.47
3:D:402:PCZ:O91	3:D:402:PCZ:C15	2.62	0.47
1:B:304:PRO:HG2	1:C:303:PRO:CB	2.44	0.47
1:B:17:THR:HB	1:B:18:PRO:HD3	1.98	0.46
1:A:95:GLU:N	1:A:95:GLU:OE1	2.39	0.45
1:C:5:PRO:CB	1:C:7:GLN:HE22	2.29	0.45
1:B:33:ILE:HD12	1:B:33:ILE:N	2.32	0.44
3:C:401:PCZ:C15	3:C:401:PCZ:O91	2.61	0.43
1:B:78:ILE:HD11	1:B:85:LEU:HG	2.01	0.43
1:D:33:ILE:HD12	1:D:33:ILE:N	2.34	0.43
1:A:262:THR:HG21	1:A:296[B]:ARG:CB	2.49	0.43
1:A:304:PRO:HD2	1:D:304:PRO:HD2	2.00	0.43
1:A:346:ASN:HB2	1:A:347:PRO:HD3	2.00	0.43
1:D:61:GLU:HB2	1:D:322:PHE:CD1	2.54	0.43
1:B:260:TRP:CE3	1:B:300:ALA:HA	2.55	0.42
1:C:182:LEU:O	1:C:232:ARG:HD3	2.18	0.42
1:A:241:LEU:HD21	1:A:253[B]:GLN:NE2	2.35	0.42
1:A:262:THR:HG21	1:A:296[A]:ARG:HB2	2.00	0.42
1:C:98:ALA:HB1	1:C:100:GLN:OE1	2.19	0.42
1:C:118:PRO:HD2	1:C:153:SER:OG	2.20	0.42
1:D:238:LEU:CD2	1:D:330:PRO:HA	2.48	0.42
3:A:404:PCZ:O91	3:A:404:PCZ:H15	2.19	0.41
1:C:243:ILE:CD1	1:C:252:ILE:HD12	2.50	0.41
1:B:216:LEU:HA	1:B:219:GLU:OE1	2.21	0.41
1:C:75:GLY:HA2	1:C:78:ILE:HD12	2.03	0.41
1:B:241:LEU:HD23	1:B:249:GLN:NE2	2.36	0.41
1:C:17:THR:HB	1:C:18:PRO:HD3	2.04	0.41
1:B:283:ILE:HD13	1:B:283:ILE:HG21	1.79	0.40
1:A:224:LYS:HD2	1:A:224:LYS:N	2.37	0.40
1:B:336:ILE:HG12	1:B:337:VAL:N	2.36	0.40
1:A:96:LEU:HD12	1:A:96:LEU:HA	1.89	0.40
1:C:250:GLN:HG2	4:C:592:HOH:O	2.22	0.40
1:A:5:PRO:HD2	1:A:8:ILE:HD12	2.03	0.40
1:C:5:PRO:HB3	1:C:7:GLN:HE22	1.87	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:243:ILE:HD12	1:D:252:ILE:HD12	2.01	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:57[B]:GLN:NE2	4:C:616:HOH:O[3_455]	1.97	0.23

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/358 (101%)	351 (97%)	10 (3%)	0	100	100
1	B	357/358 (100%)	349 (98%)	6 (2%)	2 (1%)	33	27
1	C	359/358 (100%)	351 (98%)	8 (2%)	0	100	100
1	D	359/358 (100%)	348 (97%)	10 (3%)	1 (0%)	50	48
All	All	1436/1432 (100%)	1399 (97%)	34 (2%)	3 (0%)	56	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	331	GLU
1	D	36	GLY
1	B	36	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/291 (98%)	283 (99%)	3 (1%)	85	90
1	B	279/291 (96%)	273 (98%)	6 (2%)	64	68
1	C	285/291 (98%)	279 (98%)	6 (2%)	66	69
1	D	285/291 (98%)	281 (99%)	4 (1%)	78	83
All	All	1135/1164 (98%)	1116 (98%)	19 (2%)	73	77

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	13	HIS
1	B	199	TYR
1	B	236	SER
1	B	262	THR
1	B	303	PRO
1	B	314	HIS
1	A	314	HIS
1	A	324	SER
1	A	361	GLN
1	C	21	GLU
1	C	128	SER
1	C	197	LYS
1	C	250	GLN
1	C	265	MET
1	C	314	HIS
1	D	129	SER
1	D	232	ARG
1	D	236	SER
1	D	314	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	249	GLN
1	C	57	GLN
1	C	198	ASN
1	C	244	ASN
1	D	190	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	401	-	4,4,4	0.30	0	6,6,6	0.33	0
2	PO4	A	402	-	4,4,4	0.33	0	6,6,6	0.30	0
2	PO4	A	403	-	4,4,4	0.15	0	6,6,6	0.33	0
3	PCZ	A	404	1	27,27,27	1.93	6 (22%)	32,37,37	4.97	10 (31%)
2	PO4	B	401	-	4,4,4	0.28	0	6,6,6	0.37	0
3	PCZ	B	402	1	27,27,27	1.92	6 (22%)	32,37,37	4.89	13 (40%)
3	PCZ	C	401	1	27,27,27	1.66	3 (11%)	32,37,37	4.26	10 (31%)
2	PO4	D	401	-	4,4,4	0.26	0	6,6,6	0.32	0
3	PCZ	D	402	1	27,27,27	2.04	6 (22%)	32,37,37	4.45	10 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	401	-	-	0/0/0/0	0/0/0/0
2	PO4	A	402	-	-	0/0/0/0	0/0/0/0
2	PO4	A	403	-	-	0/0/0/0	0/0/0/0
3	PCZ	A	404	1	-	0/15/38/38	0/1/2/2
2	PO4	B	401	-	-	0/0/0/0	0/0/0/0
3	PCZ	B	402	1	-	0/15/38/38	0/1/2/2
3	PCZ	C	401	1	-	0/15/38/38	0/1/2/2
2	PO4	D	401	-	-	0/0/0/0	0/0/0/0
3	PCZ	D	402	1	-	0/15/38/38	0/1/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	PCZ	O12-N11	-7.16	1.26	1.40
3	D	402	PCZ	O12-N11	-6.95	1.26	1.40
3	C	401	PCZ	O12-N11	-6.83	1.26	1.40
3	A	404	PCZ	O12-N11	-5.98	1.28	1.40
3	A	404	PCZ	O62-C61	4.11	1.34	1.22
3	D	402	PCZ	C2-N8	-3.85	1.42	1.46
3	D	402	PCZ	O62-C61	3.84	1.34	1.22
3	A	404	PCZ	C2-N8	-3.37	1.42	1.46
3	B	402	PCZ	C2-C1	3.27	1.53	1.49
3	A	404	PCZ	C2-C1	3.02	1.52	1.49
3	D	402	PCZ	O63-C61	-2.96	1.20	1.30
3	C	401	PCZ	C2-C1	2.82	1.52	1.49
3	D	402	PCZ	C52-S1	-2.64	1.76	1.82
3	B	402	PCZ	C14-C10	-2.53	1.43	1.48
3	B	402	PCZ	C10-C9	-2.49	1.46	1.50
3	D	402	PCZ	C3-S1	-2.41	1.76	1.82
3	A	404	PCZ	C52-S1	-2.39	1.76	1.82
3	B	402	PCZ	C52-S1	-2.39	1.76	1.82
3	A	404	PCZ	C6-C61	-2.39	1.45	1.48
3	C	401	PCZ	C3-S1	-2.23	1.77	1.82
3	B	402	PCZ	C2-N8	-2.03	1.44	1.46

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	404	PCZ	C13-O12-N11	16.25	126.02	108.50
3	B	402	PCZ	C3-N7-C6	16.15	130.38	116.18
3	B	402	PCZ	O12-N11-C10	14.88	130.09	111.33
3	C	401	PCZ	C3-N7-C6	14.15	128.62	116.18
3	A	404	PCZ	C3-N7-C6	13.69	128.22	116.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	402	PCZ	O12-N11-C10	13.55	128.42	111.33
3	D	402	PCZ	C3-N7-C6	13.16	127.75	116.18
3	C	401	PCZ	O12-N11-C10	11.60	125.96	111.33
3	D	402	PCZ	C13-O12-N11	10.31	119.62	108.50
3	A	404	PCZ	O12-N11-C10	10.23	124.23	111.33
3	C	401	PCZ	C13-O12-N11	9.56	118.81	108.50
3	A	404	PCZ	C2-C3-N7	-9.36	101.53	111.19
3	B	402	PCZ	C13-O12-N11	9.16	118.38	108.50
3	B	402	PCZ	C52-S1-C3	8.71	110.96	94.37
3	D	402	PCZ	C52-S1-C3	8.06	109.72	94.37
3	C	401	PCZ	C52-S1-C3	7.55	108.74	94.37
3	A	404	PCZ	C52-S1-C3	7.51	108.67	94.37
3	A	404	PCZ	C61-C6-N7	-5.40	108.78	115.86
3	C	401	PCZ	C2-C3-N7	-5.33	105.69	111.19
3	D	402	PCZ	C15-C14-N18	5.14	116.98	109.64
3	D	402	PCZ	C2-C3-N7	-4.90	106.14	111.19
3	B	402	PCZ	C15-C14-N18	4.77	116.44	109.64
3	B	402	PCZ	C2-C3-N7	-4.50	106.55	111.19
3	A	404	PCZ	C5-C52-S1	-4.32	102.38	111.97
3	C	401	PCZ	C10-C9-N8	3.95	120.82	114.27
3	D	402	PCZ	C14-C15-S16	-3.90	107.41	111.97
3	B	402	PCZ	C14-C15-S16	-3.70	107.66	111.97
3	C	401	PCZ	C2-C3-S1	-3.32	106.72	112.76
3	B	402	PCZ	C10-C14-N18	-3.24	116.64	123.86
3	C	401	PCZ	C15-C14-N18	3.23	114.25	109.64
3	A	404	PCZ	C15-C14-N18	3.09	114.05	109.64
3	C	401	PCZ	C61-C6-N7	-3.04	111.88	115.86
3	B	402	PCZ	O63-C61-O62	-2.85	116.89	123.61
3	B	402	PCZ	C52-C5-C51	-2.84	115.10	121.40
3	A	404	PCZ	O91-C9-C10	2.73	123.62	120.52
3	B	402	PCZ	C61-C6-N7	-2.72	112.29	115.86
3	B	402	PCZ	C3-C2-N8	-2.71	104.65	109.93
3	A	404	PCZ	O62-C61-C6	-2.56	115.59	120.97
3	C	401	PCZ	O91-C9-C10	-2.52	117.67	120.52
3	D	402	PCZ	C52-C5-C51	-2.43	116.02	121.40
3	D	402	PCZ	O63-C61-O62	-2.30	118.19	123.61
3	D	402	PCZ	C61-C6-N7	-2.19	112.99	115.86
3	B	402	PCZ	C2-C3-S1	-2.06	109.02	112.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	358/358 (100%)	-0.26	2 (0%) 86 90	22, 33, 51, 68	0
1	B	358/358 (100%)	-0.23	6 (1%) 67 71	25, 37, 55, 69	0
1	C	358/358 (100%)	-0.22	7 (1%) 62 66	24, 35, 54, 77	0
1	D	358/358 (100%)	-0.22	3 (0%) 83 87	23, 33, 52, 70	0
All	All	1432/1432 (100%)	-0.23	18 (1%) 74 78	22, 34, 53, 77	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	4	ALA	4.1
1	D	241	LEU	3.1
1	D	205	GLU	2.7
1	C	49	ALA	2.6
1	C	244	ASN	2.4
1	C	326	VAL	2.4
1	D	337	VAL	2.4
1	B	331	GLU	2.4
1	B	241	LEU	2.2
1	C	339	LEU	2.2
1	A	242	ASP	2.2
1	B	36	GLY	2.2
1	B	337	VAL	2.2
1	A	339	LEU	2.1
1	C	336	ILE	2.1
1	B	326	VAL	2.1
1	B	336	ILE	2.1
1	C	331	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	PO4	A	402	5/5	0.53	9.63	122,137,144,153	0
2	PO4	B	401	5/5	0.23	4.28	58,67,87,88	0
2	PO4	A	403	5/5	0.20	3.75	70,74,74,76	0
2	PO4	D	401	5/5	0.17	3.40	64,71,85,86	0
2	PO4	A	401	5/5	0.23	2.29	69,72,77,85	0
3	PCZ	A	404	26/26	0.18	1.76	38,58,70,93	0
3	PCZ	B	402	26/26	0.15	1.47	42,58,63,79	0
3	PCZ	C	401	26/26	0.14	0.81	35,58,66,70	0
3	PCZ	D	402	26/26	0.14	0.81	38,56,69,74	0

6.5 Other polymers ⓘ

There are no such residues in this entry.