



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 05:42 AM GMT

PDB ID : 4KML
Title : Probing the N-terminal beta-sheet conversion in the crystal structure of the full-length human prion protein bound to a Nanobody
Authors : Abskharon, R.N.N; Giachin, G.; Wohlkonig, A.; Soror, S.H.; Pardon, E.; Legname, G.; Steyaert, J.
Deposited on : 2013-05-08
Resolution : 1.50 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1513 (1.50-1.50)
Clashscore	79885	1768 (1.50-1.50)
Ramachandran outliers	78287	1720 (1.50-1.50)
Sidechain outliers	78261	1718 (1.50-1.50)
RSRZ outliers	66119	1514 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	241	
2	B	130	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1920 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major prion protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	1	0
			894	554	155	175	10			

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	EXPRESSION TAG	UNP P04156
A	-8	GLY	-	EXPRESSION TAG	UNP P04156
A	-7	SER	-	EXPRESSION TAG	UNP P04156
A	-6	SER	-	EXPRESSION TAG	UNP P04156
A	-5	HIS	-	EXPRESSION TAG	UNP P04156
A	-4	HIS	-	EXPRESSION TAG	UNP P04156
A	-3	HIS	-	EXPRESSION TAG	UNP P04156
A	-2	HIS	-	EXPRESSION TAG	UNP P04156
A	-1	HIS	-	EXPRESSION TAG	UNP P04156
A	0	HIS	-	EXPRESSION TAG	UNP P04156
A	1	SER	-	EXPRESSION TAG	UNP P04156
A	2	SER	-	EXPRESSION TAG	UNP P04156
A	3	GLY	-	EXPRESSION TAG	UNP P04156
A	4	LEU	-	EXPRESSION TAG	UNP P04156
A	5	VAL	-	EXPRESSION TAG	UNP P04156
A	6	PRO	-	EXPRESSION TAG	UNP P04156
A	7	ARG	-	EXPRESSION TAG	UNP P04156
A	8	GLY	-	EXPRESSION TAG	UNP P04156
A	9	SER	-	EXPRESSION TAG	UNP P04156
A	10	HIS	-	EXPRESSION TAG	UNP P04156
A	11	MET	-	EXPRESSION TAG	UNP P04156
A	12	ALA	-	EXPRESSION TAG	UNP P04156
A	13	SER	-	EXPRESSION TAG	UNP P04156
A	14	MET	-	EXPRESSION TAG	UNP P04156
A	15	THR	-	EXPRESSION TAG	UNP P04156
A	16	GLY	-	EXPRESSION TAG	UNP P04156
A	17	GLY	-	EXPRESSION TAG	UNP P04156

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	GLN	-	EXPRESSION TAG	UNP P04156
A	19	GLN	-	EXPRESSION TAG	UNP P04156
A	20	MET	-	EXPRESSION TAG	UNP P04156
A	21	GLY	-	EXPRESSION TAG	UNP P04156
A	22	ARG	-	EXPRESSION TAG	UNP P04156
A	23	GLY	-	EXPRESSION TAG	UNP P04156

- Molecule 2 is a protein called Nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	123	Total	C	N	O	S	0	0	0
			937	579	171	182	5			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	43	Total	O	0	0
			43	43		
3	B	46	Total	O	0	0
			46	46		

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	131.44Å 45.92Å 44.96Å 90.00° 96.09° 90.00°	Depositor
Resolution (Å)	31.61 – 1.50 31.61 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.5 (31.61-1.50) 98.5 (31.61-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.38 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.175 , 0.192 0.186 , 0.202	Depositor DCC
R_{free} test set	2112 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	14.1	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 34.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 42231 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1920	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	2/913 (0.2%)	1.35	9/1232 (0.7%)
2	B	1.26	1/955 (0.1%)	1.29	10/1286 (0.8%)
All	All	1.30	3/1868 (0.2%)	1.32	19/2518 (0.8%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	168	GLU	CD-OE2	6.10	1.32	1.25
2	B	60	TYR	CE1-CZ	-6.08	1.30	1.38
1	A	152	GLU	CD-OE2	-5.07	1.20	1.25

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	151	ARG	NE-CZ-NH2	-9.78	115.41	120.30
1	A	151	ARG	NE-CZ-NH1	8.09	124.35	120.30
2	B	67	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	A	164	ARG	NE-CZ-NH2	-7.52	116.54	120.30
2	B	108	ARG	NE-CZ-NH2	7.39	124.00	120.30
1	A	167	ASP	CB-CG-OD1	7.10	124.69	118.30
2	B	112	ASP	CB-CG-OD1	6.53	124.18	118.30
2	B	60	TYR	CB-CG-CD1	-6.36	117.18	121.00
2	B	56	ASP	CB-CG-OD1	6.36	124.02	118.30
2	B	68	PHE	CB-CG-CD1	5.96	124.97	120.80
1	A	136	ARG	NE-CZ-NH2	5.68	123.14	120.30
2	B	80	TYR	CB-CG-CD1	-5.57	117.66	121.00
1	A	145	TYR	CB-CG-CD2	5.46	124.28	121.00
2	B	90	ASP	CB-CG-OD1	5.31	123.08	118.30
2	B	18	LEU	CA-CB-CG	5.26	127.40	115.30
1	A	156	ARG	NE-CZ-NH1	5.21	122.90	120.30
1	A	175	PHE	CG-CD1-CE1	5.10	126.41	120.80
2	B	108	ARG	CA-CB-CG	5.07	124.55	113.40
1	A	164	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	894	0	833	7	0
2	B	937	0	904	3	0
3	A	43	0	0	2	0
3	B	46	0	0	0	0
All	All	1920	0	1737	9	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (9) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:192:THR:HG23	3:A:343:HOH:O	2.03	0.57
1:A:122:VAL:HG13	1:A:186:GLN:HE21	1.70	0.57
2:B:12:VAL:HG11	2:B:86:LEU:HD13	1.97	0.46
2:B:22:CYS:SG	2:B:96:CYS:CB	3.05	0.44
1:A:117:ALA:HB3	1:A:129[A]:MET:CE	2.49	0.43
1:A:188:THR:O	1:A:192:THR:HG23	2.18	0.43
1:A:174:ASN:ND2	2:B:106:ARG:HH11	2.17	0.42
1:A:192:THR:CG2	3:A:343:HOH:O	2.65	0.42
1:A:117:ALA:HB3	1:A:129[A]:MET:HE2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	108/241 (45%)	108 (100%)	0	0	100	100
2	B	121/130 (93%)	119 (98%)	2 (2%)	0	100	100
All	All	229/371 (62%)	227 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/183 (53%)	95 (98%)	2 (2%)	66	30
2	B	97/104 (93%)	96 (99%)	1 (1%)	85	65
All	All	194/287 (68%)	191 (98%)	3 (2%)	75	47

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166	MET
1	A	204	LYS
2	B	123	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	109/241 (45%)	0.16	9 (8%) 11 13	8, 16, 40, 63	0
2	B	123/130 (94%)	0.07	2 (1%) 68 75	8, 16, 27, 45	0
All	All	232/371 (62%)	0.11	11 (4%) 30 34	8, 16, 37, 63	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	225	TYR	7.3
1	A	193	THR	5.0
1	A	195	GLY	4.4
1	A	194	LYS	3.6
1	A	192	THR	3.4
1	A	224	ALA	3.1
2	B	1	ALA	2.9
2	B	27	ARG	2.8
1	A	190	THR	2.5
1	A	223	GLN	2.4
1	A	142	GLY	2.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.