



Full wwPDB X-ray Structure Validation Report (i)

Sep 16, 2014 – 02:06 AM EDT

PDB ID : 4KMO
Title : Crystal Structure of the Vps33-Vps16 HOPS subcomplex from *Chaetomium thermophilum*
Authors : Baker, R.W.; Jeffrey, P.D.; Hughson, F.M.
Deposited on : 2013-05-08
Resolution : 2.60 Å (reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23489
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23489

1 Overall quality at a glance (i)


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6810 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Small conjugating protein ligase-like protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	574	Total	C 4536	N 2870	O 803	S 852	Se 4	0	0	0

- Molecule 2 is a protein called Putative vacuolar protein sorting-associated protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	265	Total	C 2122	N 1347	O 374	S 392	Se 2	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O 5	S 4	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	118	Total O 118 118	0	0
4	B	29	Total O 29 29	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Small conjugating protein ligase-like protein

Chain A:



- Molecule 2: Putative vacuolar protein sorting-associated protein



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	100.27Å 100.27Å 176.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.66 – 2.60 48.66 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.1 (48.66-2.60) 98.1 (48.66-2.60)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.65 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R , R_{free}	0.225 , 0.254 0.222 , 0.253	Depositor DCC
R_{free} test set	1599 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	56.8	Xtriage
Anisotropy	0.197	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 35.1	EDS
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
L-test for twinning	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	0 of 31609 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6810	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section:
SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.21	0/4597	0.38	0/6185
2	B	0.21	0/2153	0.39	0/2894
All	All	0.21	0/6750	0.38	0/9079

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4536	0	4597	44	0
2	B	2122	0	2138	33	0
3	B	5	0	0	0	0
4	A	118	0	0	1	0
4	B	29	0	0	1	0
All	All	6810	0	6735	74	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (74) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:452:GLN:NE2	1:A:654:ALA:O	2.28	0.67
1:A:312:LEU:HD23	1:A:323:LEU:HD23	1.81	0.62
1:A:539:LEU:O	1:A:543:LYS:N	2.30	0.59
2:B:762:ASN:N	2:B:762:ASN:OD1	2.35	0.58
1:A:600:ASP:OD1	1:A:600:ASP:N	2.36	0.58
1:A:358:GLN:NE2	4:A:814:HOH:O	2.37	0.57
2:B:636:THR:HB	2:B:671:LEU:HD13	1.86	0.57
2:B:697:LEU:HB3	2:B:706:ALA:HB2	1.87	0.56
1:A:607:VAL:HG22	1:A:636:CYS:HB2	1.87	0.56
2:B:776:LEU:HD11	2:B:784:MSE:HE1	1.87	0.55
1:A:447:GLN:HB2	2:B:690:VAL:HB	1.89	0.55
2:B:536:VAL:HG13	2:B:537:PRO:HD3	1.90	0.54
2:B:581:VAL:HG13	2:B:588:SER:HA	1.89	0.52
1:A:73:ARG:HG3	1:A:105:HIS:HA	1.92	0.51
1:A:499:LEU:HD13	1:A:568:ILE:HG21	1.91	0.51
2:B:721:TRP:HZ3	2:B:740:ILE:HD13	1.76	0.50
1:A:627:GLU:OE2	1:A:630:ARG:NH1	2.44	0.50
1:A:174:ARG:HA	1:A:206:MSE:HE1	1.93	0.50
1:A:301:GLN:NE2	1:A:303:ASP:OD2	2.34	0.50
1:A:78:ILE:HG12	1:A:110:PHE:HB2	1.94	0.49
1:A:496:ARG:HA	1:A:501:LEU:HB2	1.94	0.49
1:A:137:LEU:HD22	1:A:139:LEU:HG	1.94	0.48
2:B:546:GLU:HG2	2:B:580:VAL:HG13	1.95	0.48
1:A:23:LEU:HD12	1:A:137:LEU:HG	1.95	0.48
2:B:652:GLY:O	2:B:654:GLN:N	2.46	0.47
1:A:443:ARG:O	1:A:447:GLN:HG2	2.15	0.46
2:B:722:TRP:CH2	2:B:752:PRO:HB3	2.50	0.46
1:A:66:LYS:NZ	1:A:89:ALA:O	2.48	0.46
2:B:740:ILE:O	2:B:743:GLN:HG2	2.15	0.46
1:A:494:TYR:HD2	1:A:495:LEU:HD13	1.80	0.46
1:A:115:ARG:HG3	1:A:134:ILE:HG21	1.96	0.46
1:A:96:ARG:HG3	1:A:100:GLU:OE1	2.16	0.45
1:A:442:ARG:HG2	1:A:457:LEU:HD13	1.99	0.45
2:B:574:LEU:HD13	2:B:574:LEU:HA	1.81	0.45
1:A:170:PHE:CE2	1:A:174:ARG:HD2	2.52	0.45
2:B:738:GLU:HG2	2:B:742:ARG:HH21	1.81	0.45
2:B:640:SER:HB2	2:B:668:LYS:HD2	1.97	0.45
2:B:545:ASP:O	2:B:547:LEU:N	2.48	0.45
2:B:628:ARG:NH1	4:B:1018:HOH:O	2.50	0.45
2:B:724:ARG:HE	2:B:740:ILE:HD11	1.82	0.45
1:A:237:ASP:HB3	1:A:240:VAL:HG23	1.99	0.45
1:A:58:ASP:N	1:A:58:ASP:OD1	2.50	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:770:ILE:HD11	2:B:781:THR:HG23	1.98	0.45
1:A:385:LYS:HB3	1:A:409:LEU:HD21	2.00	0.44
1:A:443:ARG:NH2	2:B:666:GLU:OE1	2.51	0.44
1:A:438:LEU:HD22	1:A:468:SER:HB2	2.00	0.44
1:A:67:ASN:ND2	1:A:67:ASN:O	2.51	0.44
2:B:776:LEU:HA	2:B:776:LEU:HD12	1.71	0.44
1:A:494:TYR:CE2	1:A:498:GLN:HG3	2.53	0.43
1:A:452:GLN:HB2	1:A:452:GLN:HE21	1.62	0.43
1:A:75:ILE:HB	1:A:107:PHE:HD1	1.83	0.43
1:A:382:ILE:HD13	1:A:414:THR:HG22	2.00	0.43
2:B:697:LEU:HD12	2:B:697:LEU:HA	1.86	0.43
1:A:654:ALA:HB3	2:B:631:LEU:HD13	1.99	0.42
2:B:595:ALA:O	2:B:605:ASP:HB2	2.19	0.42
1:A:317:PHE:HA	1:A:320:VAL:HG23	2.01	0.42
2:B:609:LEU:HA	2:B:609:LEU:HD12	1.84	0.42
1:A:655:THR:O	1:A:655:THR:OG1	2.33	0.42
2:B:540:LEU:HA	2:B:540:LEU:HD12	1.90	0.42
2:B:770:ILE:HD13	2:B:784:MSE:SE	2.70	0.42
1:A:409:LEU:O	1:A:414:THR:HG23	2.19	0.42
2:B:655:LYS:HD3	2:B:655:LYS:H	1.85	0.41
1:A:398:ASP:HA	1:A:399:PRO:HD3	1.91	0.41
1:A:32:ASN:HD21	1:A:71:SER:HB3	1.85	0.41
1:A:76:VAL:HG22	1:A:108:HIS:HB2	2.03	0.41
1:A:192:LYS:HE2	1:A:572:ALA:HB3	2.02	0.41
2:B:734:TRP:HB3	2:B:765:LEU:HD22	2.03	0.41
1:A:386:LEU:HD23	1:A:422:LEU:HD11	2.03	0.41
1:A:331:LEU:HD12	1:A:364:LEU:HD22	2.03	0.41
2:B:531:ARG:O	2:B:533:GLY:N	2.49	0.40
2:B:770:ILE:HD12	2:B:771:PRO:HA	2.03	0.40
2:B:782:ILE:HG13	2:B:782:ILE:H	1.56	0.40
2:B:534:ARG:HA	2:B:534:ARG:HD3	1.67	0.40
1:A:499:LEU:HA	1:A:499:LEU:HD12	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	562/669 (84%)	528 (94%)	27 (5%)	7 (1%)	19 39
2	B	261/333 (78%)	239 (92%)	15 (6%)	7 (3%)	8 13
All	All	823/1002 (82%)	767 (93%)	42 (5%)	14 (2%)	14 26

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	506	VAL
2	B	530	PRO
2	B	774	THR
1	A	71	SER
1	A	97	ILE
1	A	505	GLU
2	B	781	THR
1	A	105	HIS
1	A	558	GLY
2	B	533	GLY
2	B	653	ASN
2	B	703	HIS
1	A	485	GLY
2	B	771	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	494/556 (89%)	465 (94%)	29 (6%)	28 52
2	B	226/272 (83%)	212 (94%)	14 (6%)	26 49
All	All	720/828 (87%)	677 (94%)	43 (6%)	27 51

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	27	VAL
1	A	31	LYS
1	A	39	LEU
1	A	58	ASP
1	A	63	LEU
1	A	85	ARG
1	A	96	ARG
1	A	114	ARG
1	A	115	ARG
1	A	117	LEU
1	A	124	GLU
1	A	137	LEU
1	A	162	LEU
1	A	176	LEU
1	A	297	LYS
1	A	298	ARG
1	A	323	LEU
1	A	324	LEU
1	A	426	TYR
1	A	452	GLN
1	A	495	LEU
1	A	499	LEU
1	A	509	GLN
1	A	524	LEU
1	A	533	LEU
1	A	603	THR
1	A	620	LEU
1	A	630	ARG
2	B	525	LEU
2	B	528	HIS
2	B	536	VAL
2	B	547	LEU
2	B	574	LEU
2	B	581	VAL
2	B	584	ARG
2	B	609	LEU
2	B	655	LYS
2	B	669	MSE
2	B	682	THR
2	B	697	LEU
2	B	758	LEU
2	B	762	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	452	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	901	-	4,4,4	0.24	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	B	901	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	574/669 (85%)	0.28	33 (5%) 23 19	26, 48, 115, 157	0
2	B	265/333 (79%)	0.43	33 (12%) 5 3	39, 64, 115, 140	0
All	All	839/1002 (83%)	0.33	66 (7%) 13 10	26, 54, 115, 157	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	558	GLY	20.3
1	A	357	TYR	9.0
1	A	546	GLY	6.6
1	A	559	ALA	6.1
2	B	780	GLN	5.3
1	A	557	GLY	5.1
2	B	656	GLU	4.6
1	A	545	SER	4.6
1	A	544	GLY	4.5
2	B	790	MSE	4.1
1	A	104	SER	4.0
2	B	528	HIS	3.9
1	A	99	ARG	3.8
2	B	532	ALA	3.7
1	A	296	ARG	3.7
1	A	543	LYS	3.7
1	A	556	GLY	3.6
1	A	100	GLU	3.5
1	A	73	ARG	3.4
1	A	60	PHE	3.4
2	B	679	ARG	3.4
2	B	686	VAL	3.3
1	A	600	ASP	3.3
1	A	268	ASP	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	96	ARG	3.1
1	A	480	SER	3.0
2	B	721	TRP	3.0
2	B	778	PRO	3.0
2	B	671	LEU	2.9
2	B	538	LEU	2.9
1	A	76	VAL	2.9
2	B	760	ALA	2.9
1	A	131	ASP	2.8
2	B	758	LEU	2.8
2	B	789	GLY	2.8
1	A	483	SER	2.8
2	B	779	GLY	2.8
2	B	523	THR	2.8
2	B	545	ASP	2.7
2	B	653	ASN	2.7
2	B	529	GLU	2.7
1	A	61	PHE	2.6
2	B	685	PHE	2.6
2	B	521	LEU	2.5
2	B	534	ARG	2.5
2	B	788	CYS	2.5
1	A	436	LYS	2.5
2	B	717	GLU	2.5
1	A	132	ALA	2.5
1	A	361	GLN	2.5
2	B	687	GLY	2.4
2	B	773	CYS	2.4
2	B	784	MSE	2.4
2	B	771	PRO	2.4
2	B	783	THR	2.4
1	A	35	ILE	2.4
1	A	90	ILE	2.4
1	A	360	GLU	2.2
2	B	761	GLY	2.1
1	A	97	ILE	2.1
1	A	69	GLY	2.1
2	B	742	ARG	2.1
1	A	118	PHE	2.1
2	B	524	GLU	2.1
2	B	754	PHE	2.1
1	A	126	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	B	901	5/5	0.13	-0.81	79,80,81,81	0

6.5 Other polymers (i)

There are no such residues in this entry.