



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 08:46 AM GMT

PDB ID : 4KP3
Title : Crystal Structure of MyoVa-GTD in Complex with Two Cargos
Authors : Wei, Z.; Liu, X.; Yu, C.; Zhang, M.
Deposited on : 2013-05-13
Resolution : 2.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	404	
1	B	404	
2	C	103	
2	D	103	
3	E	43	
3	F	43	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7359 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-Va.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	361	Total	C	N	O	S	0	1	0
			2908	1850	501	536	21			
1	B	355	Total	C	N	O	S	0	0	0
			2847	1810	491	525	21			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1450	HIS	-	EXPRESSION TAG	UNP Q99104
A	1451	HIS	-	EXPRESSION TAG	UNP Q99104
A	1452	HIS	-	EXPRESSION TAG	UNP Q99104
A	1453	HIS	-	EXPRESSION TAG	UNP Q99104
A	1454	HIS	-	EXPRESSION TAG	UNP Q99104
A	1455	HIS	-	EXPRESSION TAG	UNP Q99104
A	1456	SER	-	EXPRESSION TAG	UNP Q99104
A	1457	SER	-	EXPRESSION TAG	UNP Q99104
A	1458	GLY	-	EXPRESSION TAG	UNP Q99104
A	1459	LEU	-	EXPRESSION TAG	UNP Q99104
A	1460	GLY	-	EXPRESSION TAG	UNP Q99104
A	1461	VAL	-	EXPRESSION TAG	UNP Q99104
A	1462	LEU	-	EXPRESSION TAG	UNP Q99104
A	1463	PHE	-	EXPRESSION TAG	UNP Q99104
A	1464	GLN	-	EXPRESSION TAG	UNP Q99104
A	1465	GLY	-	EXPRESSION TAG	UNP Q99104
A	1466	PRO	-	EXPRESSION TAG	UNP Q99104
A	1467	GLY	-	EXPRESSION TAG	UNP Q99104
A	1468	SER	-	EXPRESSION TAG	UNP Q99104
A	1674	SER	CYS	ENGINEERED MUTATION	UNP Q99104
B	1450	HIS	-	EXPRESSION TAG	UNP Q99104
B	1451	HIS	-	EXPRESSION TAG	UNP Q99104
B	1452	HIS	-	EXPRESSION TAG	UNP Q99104
B	1453	HIS	-	EXPRESSION TAG	UNP Q99104
B	1454	HIS	-	EXPRESSION TAG	UNP Q99104

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1455	HIS	-	EXPRESSION TAG	UNP Q99104
B	1456	SER	-	EXPRESSION TAG	UNP Q99104
B	1457	SER	-	EXPRESSION TAG	UNP Q99104
B	1458	GLY	-	EXPRESSION TAG	UNP Q99104
B	1459	LEU	-	EXPRESSION TAG	UNP Q99104
B	1460	GLY	-	EXPRESSION TAG	UNP Q99104
B	1461	VAL	-	EXPRESSION TAG	UNP Q99104
B	1462	LEU	-	EXPRESSION TAG	UNP Q99104
B	1463	PHE	-	EXPRESSION TAG	UNP Q99104
B	1464	GLN	-	EXPRESSION TAG	UNP Q99104
B	1465	GLY	-	EXPRESSION TAG	UNP Q99104
B	1466	PRO	-	EXPRESSION TAG	UNP Q99104
B	1467	GLY	-	EXPRESSION TAG	UNP Q99104
B	1468	SER	-	EXPRESSION TAG	UNP Q99104
B	1674	SER	CYS	ENGINEERED MUTATION	UNP Q99104

- Molecule 2 is a protein called RILP-like protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	81	Total	C	N	O	S	0	0	0
			642	398	110	130	4			
2	D	82	Total	C	N	O	S	0	1	0
			659	409	115	131	4			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	GLY	-	EXPRESSION TAG	UNP Q99LE1
C	-4	PRO	-	EXPRESSION TAG	UNP Q99LE1
C	-3	GLY	-	EXPRESSION TAG	UNP Q99LE1
C	-2	SER	-	EXPRESSION TAG	UNP Q99LE1
C	-1	GLU	-	EXPRESSION TAG	UNP Q99LE1
C	0	PHE	-	EXPRESSION TAG	UNP Q99LE1
D	-5	GLY	-	EXPRESSION TAG	UNP Q99LE1
D	-4	PRO	-	EXPRESSION TAG	UNP Q99LE1
D	-3	GLY	-	EXPRESSION TAG	UNP Q99LE1
D	-2	SER	-	EXPRESSION TAG	UNP Q99LE1
D	-1	GLU	-	EXPRESSION TAG	UNP Q99LE1
D	0	PHE	-	EXPRESSION TAG	UNP Q99LE1

- Molecule 3 is a protein called Melanophilin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	14	Total	C	N	O	0	0	0
			121	78	22	21			
3	F	14	Total	C	N	O	0	0	0
			125	81	23	21			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	166	GLY	-	EXPRESSION TAG	UNP Q91V27
E	167	PRO	-	EXPRESSION TAG	UNP Q91V27
E	168	GLY	-	EXPRESSION TAG	UNP Q91V27
E	169	SER	-	EXPRESSION TAG	UNP Q91V27
F	166	GLY	-	EXPRESSION TAG	UNP Q91V27
F	167	PRO	-	EXPRESSION TAG	UNP Q91V27
F	168	GLY	-	EXPRESSION TAG	UNP Q91V27
F	169	SER	-	EXPRESSION TAG	UNP Q91V27


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	33	Total	O	0	0
			33	33		
4	C	6	Total	O	0	0
			6	6		
4	E	1	Total	O	0	0
			1	1		
4	B	13	Total	O	0	0
			13	13		
4	D	3	Total	O	0	0
			3	3		
4	F	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Unconventional myosin-Va



- Molecule 1: Unconventional myosin-Va



- Molecule 2: RILP-like protein 2




- Molecule 2: RILP-like protein 2




- Molecule 3: Melanophilin





● Molecule 3: Melanophilin

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.85Å 107.93Å 83.51Å 90.00° 96.14° 90.00°	Depositor
Resolution (Å)	38.69 – 2.40 38.69 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.8 (38.69-2.40) 97.9 (38.69-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.192 , 0.241 0.193 , 0.243	Depositor DCC
R_{free} test set	2795 reflections (5.09%)	DCC
Wilson B-factor (Å ²)	48.2	Xtriage
Anisotropy	0.224	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 28.6	EDS
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 54945 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7359	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2960	0.44	0/3995
1	B	0.24	0/2894	0.42	0/3906
2	C	0.25	0/646	0.44	0/869
2	D	0.25	0/666	0.41	0/894
3	E	0.24	0/122	0.36	0/160
3	F	0.24	0/126	0.34	0/164
All	All	0.25	0/7414	0.42	0/9988

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2908	0	0	9	0
1	B	2847	0	0	10	0
2	C	642	0	0	0	0
2	D	659	0	0	0	0
3	E	121	0	0	0	0
3	F	125	0	0	1	0
4	A	33	0	0	0	0
4	B	13	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	6	0	0	0	0
4	D	3	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
All	All	7359	0	0	19	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (19) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1726:GLU:OE2	1:A:1729:ARG:NH1	2.28	0.66
1:B:1630:HIS:O	1:B:1716:GLN:NE2	2.30	0.64
1:B:1590:ASN:OD1	3:F:184:LYS:NZ	2.33	0.61
1:A:1630:HIS:O	1:A:1716:GLN:NE2	2.34	0.60
1:A:1763:GLU:OE1	1:A:1805:ARG:NH2	2.36	0.59
1:A:1667:ASN:OD1	1:A:1731:LYS:NZ	2.40	0.54
1:B:1610:GLN:OE1	1:B:1613:ARG:NH1	2.41	0.53
1:B:1548:ASP:OD2	1:B:1551:THR:OG1	2.28	0.52
1:B:1812:SER:O	1:B:1814:GLN:NE2	2.45	0.49
1:B:1754:VAL:O	1:B:1756:LYS:NZ	2.47	0.48
1:A:1713:LYS:NZ	1:A:1716:GLN:OE1	2.47	0.48
1:A:1656:THR:OG1	1:A:1657:TYR:N	2.48	0.45
1:B:1628:LEU:O	1:B:1630:HIS:N	2.50	0.45
1:B:1667:ASN:OD1	1:B:1731:LYS:NZ	2.49	0.45
1:A:1740:GLU:N	1:A:1740:GLU:OE1	2.49	0.45
1:A:1692:TYR:OH	1:A:1818:ASP:O	2.35	0.44
1:B:1726:GLU:OE1	1:B:1729:ARG:NH1	2.51	0.43
1:B:1692:TYR:OH	1:B:1818:ASP:O	2.37	0.43
1:A:1811:ASP:N	1:A:1811:ASP:OD1	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	358/404 (89%)	351 (98%)	7 (2%)	0	100	100
1	B	349/404 (86%)	341 (98%)	7 (2%)	1 (0%)	50	68
2	C	79/103 (77%)	79 (100%)	0	0	100	100
2	D	81/103 (79%)	81 (100%)	0	0	100	100
3	E	12/43 (28%)	12 (100%)	0	0	100	100
3	F	12/43 (28%)	11 (92%)	1 (8%)	0	100	100
All	All	891/1100 (81%)	875 (98%)	15 (2%)	1 (0%)	59	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1629	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/365 (90%)	323 (98%)	6 (2%)	71	88
1	B	321/365 (88%)	315 (98%)	6 (2%)	69	87
2	C	72/89 (81%)	71 (99%)	1 (1%)	78	92
2	D	74/89 (83%)	74 (100%)	0	100	100
3	E	13/40 (32%)	13 (100%)	0	100	100
3	F	14/40 (35%)	14 (100%)	0	100	100
All	All	823/988 (83%)	810 (98%)	13 (2%)	75	89

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1502	LEU
1	A	1658	THR
1	A	1701	ASN
1	A	1732	ASN
1	A	1760	ASP
1	A	1809	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	43	MET
1	B	1479	ARG
1	B	1502	LEU
1	B	1543	LYS
1	B	1660	ASP
1	B	1701	ASN
1	B	1734	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	361/404 (89%)	-0.23	3 (0%) 83 82	23, 44, 83, 118	0
1	B	355/404 (87%)	-0.09	2 (0%) 86 86	27, 55, 85, 119	1 (0%)
2	C	81/103 (78%)	-0.40	1 (1%) 75 75	28, 44, 73, 95	2 (2%)
2	D	82/103 (79%)	-0.24	0 100 100	27, 43, 71, 96	0
3	E	14/43 (32%)	-0.09	1 (7%) 16 14	41, 57, 84, 104	0
3	F	14/43 (32%)	0.71	2 (14%) 3 3	66, 86, 114, 124	0
All	All	907/1100 (82%)	-0.17	9 (0%) 79 79	23, 50, 87, 124	3 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	185	LYS	3.8
1	B	1656	THR	3.6
3	F	184	LYS	3.3
2	C	15	ASP	2.5
3	E	197	GLU	2.4
1	A	1656	THR	2.3
1	B	1573	GLU	2.2
1	A	1805	ARG	2.2
1	A	1790	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.