



wwPDB X-ray Structure Validation Summary Report i

Mar 1, 2014 – 12:33 AM GMT

PDB ID : 3KRB
Title : Structure of Aldose Reductase from Giardia Lamblia at 1.75A Resolution
Authors : Seattle Structural Genomics Center for Infectious Disease (Ssgcid)
Deposited on : 2009-11-18
Resolution : 1.75 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : **FAILED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1304 (1.76-1.76)
Ramachandran outliers	78287	1288 (1.76-1.76)
Sidechain outliers	78261	1288 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	334	
1	B	334	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5675 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	307	2455	1580	412	448	15	0	4	0
1	B	307	2453	1574	417	447	15	0	5	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP A8BGX6
A	-19	ALA	-	EXPRESSION TAG	UNP A8BGX6
A	-18	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-17	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-16	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-15	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-14	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-13	HIS	-	EXPRESSION TAG	UNP A8BGX6
A	-12	MET	-	EXPRESSION TAG	UNP A8BGX6
A	-11	GLY	-	EXPRESSION TAG	UNP A8BGX6
A	-10	THR	-	EXPRESSION TAG	UNP A8BGX6
A	-9	LEU	-	EXPRESSION TAG	UNP A8BGX6
A	-8	GLU	-	EXPRESSION TAG	UNP A8BGX6
A	-7	ALA	-	EXPRESSION TAG	UNP A8BGX6
A	-6	GLN	-	EXPRESSION TAG	UNP A8BGX6
A	-5	THR	-	EXPRESSION TAG	UNP A8BGX6
A	-4	GLN	-	EXPRESSION TAG	UNP A8BGX6
A	-3	GLY	-	EXPRESSION TAG	UNP A8BGX6
A	-2	PRO	-	EXPRESSION TAG	UNP A8BGX6
A	-1	GLY	-	EXPRESSION TAG	UNP A8BGX6
A	0	SER	-	EXPRESSION TAG	UNP A8BGX6
B	-20	MET	-	EXPRESSION TAG	UNP A8BGX6
B	-19	ALA	-	EXPRESSION TAG	UNP A8BGX6
B	-18	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-17	HIS	-	EXPRESSION TAG	UNP A8BGX6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-15	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-14	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-13	HIS	-	EXPRESSION TAG	UNP A8BGX6
B	-12	MET	-	EXPRESSION TAG	UNP A8BGX6
B	-11	GLY	-	EXPRESSION TAG	UNP A8BGX6
B	-10	THR	-	EXPRESSION TAG	UNP A8BGX6
B	-9	LEU	-	EXPRESSION TAG	UNP A8BGX6
B	-8	GLU	-	EXPRESSION TAG	UNP A8BGX6
B	-7	ALA	-	EXPRESSION TAG	UNP A8BGX6
B	-6	GLN	-	EXPRESSION TAG	UNP A8BGX6
B	-5	THR	-	EXPRESSION TAG	UNP A8BGX6
B	-4	GLN	-	EXPRESSION TAG	UNP A8BGX6
B	-3	GLY	-	EXPRESSION TAG	UNP A8BGX6
B	-2	PRO	-	EXPRESSION TAG	UNP A8BGX6
B	-1	GLY	-	EXPRESSION TAG	UNP A8BGX6
B	0	SER	-	EXPRESSION TAG	UNP A8BGX6

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total X 1 1	0	0
3	A	1	Total X 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	379	Total O 379 379	0	0
5	B	282	Total O 282 282	0	0

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: Aldose reductase

Chain A: 



- Molecule 1: Aldose reductase

Chain B: 



4 Data and refinement statistics (i)

EDS failed to run properly - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	196.77Å 66.09Å 56.29Å 90.00° 92.26° 90.00°	Depositor
Resolution (Å)	50.00 – 1.75	Depositor
% Data completeness (in resolution range)	99.9 (50.00-1.75)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.5.0104	Depositor
R , R_{free}	0.144 , 0.173	Depositor
Wilson B-factor (Å ²)	15.1	Xtrriage
Anisotropy	0.080	Xtrriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtrriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	0 of 72824 reflections	Xtrriage
Total number of atoms	5675	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, NAP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/2532	0.74	2/3446 (0.1%)
1	B	0.70	0/2533	0.70	1/3452 (0.0%)
All	All	0.73	0/5065	0.72	3/6898 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	LYS	CD-CE-NZ	5.20	123.65	111.70
1	A	6	ARG	NE-CZ-NH2	-5.09	117.75	120.30
1	B	6	ARG	NE-CZ-NH2	-5.03	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2455	0	2426	5	0
1	B	2453	0	2400	13	0
2	A	48	0	25	2	0
2	B	48	0	25	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	6	1	0
4	B	4	0	6	0	0
5	A	379	0	0	1	0
5	B	282	0	0	4	0
All	All	5675	0	4888	18	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:177:GLN:NE2	2:B:400:NAP:H2N	2.09	0.67
1:B:13:GLN:HE22	1:B:220:GLN:HE22	1.43	0.64
1:B:44[B]:GLU:OE2	5:B:445:HOH:O	2.15	0.61
1:B:138[B]:ARG:HA	1:B:138[B]:ARG:HE	1.65	0.61
1:B:156:ASN:HD21	2:B:400:NAP:H72N	1.50	0.60

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/334 (92%)	302 (98%)	5 (2%)	0	100	100
1	B	308/334 (92%)	303 (98%)	5 (2%)	0	100	100
All	All	615/668 (92%)	605 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/286 (93%)	263 (99%)	2 (1%)	89	80
1	B	262/286 (92%)	259 (99%)	3 (1%)	84	70
All	All	527/572 (92%)	522 (99%)	5 (1%)	85	75

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	SER
1	A	75	TYR
1	B	75	TYR
1	B	129	GLU
1	B	142	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	13	GLN
1	B	177	GLN
1	B	142	GLN
1	A	177	GLN
1	B	20	GLN

5.3.3 RNA [i](#)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are unknown - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAP	A	400	-	52,52,52	1.26	2 (3%)	80,80,80	1.43	11 (13%)
4	EDO	A	401	-	3,3,3	0.48	0	2,2,2	0.63	0
2	NAP	B	400	-	52,52,52	1.34	7 (13%)	80,80,80	1.64	14 (17%)
4	EDO	B	401	-	3,3,3	0.50	0	2,2,2	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	400	-	-	0/35/67/67	0/3/5/5
4	EDO	A	401	-	-	0/1/1/1	0/0/0/0
2	NAP	B	400	-	-	0/35/67/67	0/3/5/5
4	EDO	B	401	-	-	0/1/1/1	0/0/0/0

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	NAP	O7N-C7N	5.31	1.36	1.24
2	B	400	NAP	O7N-C7N	5.08	1.36	1.24
2	B	400	NAP	PA-O3	3.44	1.66	1.59
2	A	400	NAP	C2A-N3A	2.90	1.37	1.32
2	B	400	NAP	PN-O1N	2.52	1.55	1.48

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	NAP	N3A-C2A-N1A	-7.36	122.56	128.71
2	B	400	NAP	N3A-C2A-N1A	-6.56	123.22	128.71
2	B	400	NAP	N3A-C4A-N9A	4.57	133.68	125.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	400	NAP	O4B-C1B-N9A	-4.19	104.54	108.44
2	B	400	NAP	C3N-C7N-N7N	-3.92	113.31	117.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS failed to run properly - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section will therefore be empty.

6.3 Carbohydrates

EDS failed to run properly - this section will therefore be empty.

6.4 Ligands

EDS failed to run properly - this section will therefore be empty.

6.5 Other polymers

EDS failed to run properly - this section will therefore be empty.