



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 03:23 PM GMT

PDB ID : 1KTK
Title : Complex of Streptococcal pyrogenic enterotoxin C (SpeC) with a human T cell receptor beta chain (Vbeta2.1)
Authors : Sundberg, E.J.; Li, H.; Llera, A.S.; McCormick, J.K.; Tormo, J.; Karjalainen, K.; Schlievert, P.M.; Mariuzza, R.A.
Deposited on : 2002-01-16
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1216 (3.00-3.00)
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	208	
1	B	208	
1	C	208	
1	D	208	
2	E	247	
2	F	247	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9511 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exotoxin type C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	0	0
			1664	1062	275	323	4			
1	B	200	Total	C	N	O	S	0	0	0
			1624	1042	267	311	4			
1	C	208	Total	C	N	O	S	0	0	0
			1699	1083	279	333	4			
1	D	198	Total	C	N	O	S	0	0	0
			1601	1028	267	303	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	ASP	ASN	CONFLICT	UNP P13380
B	26	ASP	ASN	CONFLICT	UNP P13380
C	26	ASP	ASN	CONFLICT	UNP P13380
D	26	ASP	ASN	CONFLICT	UNP P13380

- Molecule 2 is a protein called T-cell receptor beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	247	Total	C	N	O	S	0	0	0
			1879	1178	324	370	7			
2	F	144	Total	C	N	O	S	0	0	0
			1044	656	180	203	5			


There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	10	ARG	TRP	CONFLICT	UNP P01850
E	13	ALA	CYS	CONFLICT	UNP P01850
E	50	ALA	ASN	CONFLICT	UNP P01850
E	95	LEU	-	INSERTION	UNP P01850

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
E	96	ALA	ARG	CONFLICT	UNP P01850
E	99	GLY	GLU	CONFLICT	UNP P01850
E	99	GLY	THR	CONFLICT	UNP P01850
E	101	SER	-	INSERTION	UNP P01850
E	102	THR	-	INSERTION	UNP P01850
E	?	-	PRO	DELETION	UNP P01850
E	?	-	LYS	DELETION	UNP P01850
E	?	-	ASN	DELETION	UNP P01850
E	105	THR	GLU	CONFLICT	UNP P01850
E	107	TYR	PHE	CONFLICT	UNP P01850
E	?	-	VAL	DELETION	UNP P01850
E	191	ALA	CYS	CONFLICT	UNP P01850
F	10	ARG	TRP	CONFLICT	UNP P01850
F	13	ALA	CYS	CONFLICT	UNP P01850
F	50	ALA	ASN	CONFLICT	UNP P01850
F	96	LEU	-	INSERTION	UNP P01850
F	97	ALA	ARG	CONFLICT	UNP P01850
F	98	GLY	GLU	CONFLICT	UNP P01850
F	100	GLY	THR	CONFLICT	UNP P01850
F	102	SER	-	INSERTION	UNP P01850
F	103	THR	-	INSERTION	UNP P01850
F	?	-	PRO	DELETION	UNP P01850
F	?	-	LYS	DELETION	UNP P01850
F	?	-	ASN	DELETION	UNP P01850
F	105	THR	GLU	CONFLICT	UNP P01850
F	107	TYR	PHE	CONFLICT	UNP P01850
F	?	-	VAL	DELETION	UNP P01850
F	191	ALA	CYS	CONFLICT	UNP P01850






• Molecule 1: Exotoxin type C

Chain D:




• Molecule 2: T-cell receptor beta chain

Chain E:



• Molecule 2: T-cell receptor beta chain

Chain F:





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.38Å 146.76Å 135.69Å 90.00° 98.61° 90.00°	Depositor
Resolution (Å)	6.00 – 3.00 64.38 – 2.99	Depositor EDS
% Data completeness (in resolution range)	94.1 (6.00-3.00) 80.5 (64.38-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.09 (at 3.01Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.326 , 0.334 0.342 , 0.346	Depositor DCC
R_{free} test set	1913 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	78.6	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 53.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 43707 reflections	Xtriage
F_o, F_c correlation	0.81	EDS
Total number of atoms	9511	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.65	0/1697	0.88	4/2282 (0.2%)
1	B	0.48	0/1659	0.80	3/2240 (0.1%)
1	C	0.64	1/1735 (0.1%)	0.85	3/2338 (0.1%)
1	D	0.49	0/1633	0.82	2/2197 (0.1%)
2	E	0.55	0/1924	1.02	11/2620 (0.4%)
2	F	0.48	0/1055	0.96	5/1421 (0.4%)
All	All	0.56	1/9703 (0.0%)	0.89	28/13098 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	F	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	27	CYS	CB-SG	-5.29	1.73	1.81

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	7	HIS	C-N-CD	-15.24	87.06	120.60
2	F	85	GLU	N-CA-C	-9.54	85.23	111.00
1	B	4	LYS	N-CA-C	8.53	134.03	111.00
1	D	80	SER	N-CA-C	-8.38	88.37	111.00
1	A	79	ASN	N-CA-C	7.51	131.29	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	F	130	PHE	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1664	0	1608	183	0
1	B	1624	0	1554	212	0
1	C	1699	0	1634	143	0
1	D	1601	0	1539	158	0
2	E	1879	0	1775	308	1
2	F	1044	0	965	144	0
All	All	9511	0	9075	1104	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 59.

The worst 5 of 1104 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:E:57:GLU:OE2	2:E:63:ASP:HB2	1.43	1.17
1:A:180:THR:HG22	2:E:54:ALA:HB2	1.30	1.10
2:E:15:SER:HA	2:E:82:ALA:O	1.52	1.09
2:E:79:VAL:HG13	2:E:80:THR:H	1.17	1.08
2:F:86:ASP:HB3	2:F:115:THR:HA	1.15	1.08

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:E:141:GLN:OE1	2:E:223:ASP:OD2[1_655]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/208 (95%)	145 (74%)	33 (17%)	19 (10%)	1	4
1	B	196/208 (94%)	146 (74%)	32 (16%)	18 (9%)	1	5
1	C	206/208 (99%)	152 (74%)	37 (18%)	17 (8%)	1	6
1	D	190/208 (91%)	147 (77%)	25 (13%)	18 (10%)	1	4
2	E	245/247 (99%)	143 (58%)	42 (17%)	60 (24%)	0	0
2	F	122/247 (49%)	59 (48%)	31 (25%)	32 (26%)	0	0
All	All	1156/1326 (87%)	792 (68%)	200 (17%)	164 (14%)	0	1

5 of 164 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	ILE
1	A	9	VAL
1	A	65	ARG
1	A	66	ASP
1	A	80	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/192 (95%)	153 (84%)	30 (16%)	3	16
1	B	175/192 (91%)	156 (89%)	19 (11%)	9	35
1	C	186/192 (97%)	166 (89%)	20 (11%)	9	35
1	D	171/192 (89%)	157 (92%)	14 (8%)	17	52
2	E	201/213 (94%)	162 (81%)	39 (19%)	2	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	108/213 (51%)	87 (81%)	21 (19%)	2	11
All	All	1024/1194 (86%)	881 (86%)	143 (14%)	5	23

5 of 143 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	191	ASN
1	D	157	ARG
2	F	77	LEU
1	C	197	LYS
1	D	55	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	94	GLN
1	D	94	GLN
2	E	215	GLN
1	C	191	ASN
1	D	117	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/208 (97%)	0.16	1 (0%) 88 36	29, 56, 87, 119	0
1	B	200/208 (96%)	0.25	1 (0%) 88 36	49, 81, 103, 124	0
1	C	208/208 (100%)	0.17	1 (0%) 88 36	34, 57, 84, 104	0
1	D	198/208 (95%)	0.26	4 (2%) 62 12	50, 74, 110, 130	0
2	E	247/247 (100%)	0.32	2 (0%) 83 26	42, 80, 119, 130	0
2	F	144/247 (58%)	0.66	6 (4%) 35 7	69, 112, 137, 149	0
All	All	1200/1326 (90%)	0.28	15 (1%) 74 19	29, 74, 123, 149	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	89	PHE	4.8
2	F	151	GLY	4.7
2	F	150	THR	3.8
1	D	29	VAL	3.8
2	F	147	CYS	3.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.