



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 05:13 PM GMT

PDB ID : 3KT3
Title : Crystal structure of *S. cerevisiae* tryptophanyl-tRNA synthetase in complex with TrpAMP
Authors : Zhou, M.; Dong, X.; Zhong, C.; Shen, N.; Ding, J.
Deposited on : 2009-11-24
Resolution : 2.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	438	
1	B	438	
1	C	438	
1	D	438	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	SO4	A	439	-	X
3	SO4	A	440	-	X
3	SO4	B	439	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13496 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophanyl-tRNA synthetase, cytoplasmic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	0	0
			3295	2124	546	612	13			
1	B	405	Total	C	N	O	S	0	0	0
			3265	2105	541	606	13			
1	C	403	Total	C	N	O	S	0	0	0
			3250	2096	539	602	13			
1	D	409	Total	C	N	O	S	0	0	0
			3295	2124	546	612	13			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	433	HIS	-	EXPRESSION TAG	UNP Q12109
A	434	HIS	-	EXPRESSION TAG	UNP Q12109
A	435	HIS	-	EXPRESSION TAG	UNP Q12109
A	436	HIS	-	EXPRESSION TAG	UNP Q12109
A	437	HIS	-	EXPRESSION TAG	UNP Q12109
A	438	HIS	-	EXPRESSION TAG	UNP Q12109
B	433	HIS	-	EXPRESSION TAG	UNP Q12109
B	434	HIS	-	EXPRESSION TAG	UNP Q12109
B	435	HIS	-	EXPRESSION TAG	UNP Q12109
B	436	HIS	-	EXPRESSION TAG	UNP Q12109
B	437	HIS	-	EXPRESSION TAG	UNP Q12109
B	438	HIS	-	EXPRESSION TAG	UNP Q12109
C	433	HIS	-	EXPRESSION TAG	UNP Q12109
C	434	HIS	-	EXPRESSION TAG	UNP Q12109
C	435	HIS	-	EXPRESSION TAG	UNP Q12109
C	436	HIS	-	EXPRESSION TAG	UNP Q12109
C	437	HIS	-	EXPRESSION TAG	UNP Q12109
C	438	HIS	-	EXPRESSION TAG	UNP Q12109
D	433	HIS	-	EXPRESSION TAG	UNP Q12109
D	434	HIS	-	EXPRESSION TAG	UNP Q12109
D	435	HIS	-	EXPRESSION TAG	UNP Q12109

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	436	HIS	-	EXPRESSION TAG	UNP Q12109
D	437	HIS	-	EXPRESSION TAG	UNP Q12109
D	438	HIS	-	EXPRESSION TAG	UNP Q12109

- Molecule 2 is TRYPTOPHANYL-5'AMP (three-letter code: TYM) (formula: $C_{21}H_{24}N_7O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			37	21	7	8	1		
2	B	1	Total	C	N	O	P	0	0
			37	21	7	8	1		
2	C	1	Total	C	N	O	P	0	0
			37	21	7	8	1		
2	D	1	Total	C	N	O	P	0	0
			37	21	7	8	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	63	Total 63	O 63	0	0
4	B	36	Total 36	O 36	0	0
4	C	54	Total 54	O 54	0	0
4	D	25	Total 25	O 25	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	251.76Å 251.76Å 111.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.94 – 2.60 48.94 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.1 (48.94-2.60) 96.1 (48.94-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.98 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.214 , 0.237 0.212 , 0.212	Depositor DCC
R_{free} test set	5914 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	49.8	Xtriage
Anisotropy	0.492	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 24.3	EDS
Estimated twinning fraction	0.010 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 118158 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13496	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TYM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/3377	0.50	0/4557
1	B	0.34	0/3347	0.50	0/4517
1	C	0.35	0/3332	0.48	0/4496
1	D	0.35	0/3377	0.50	1/4557 (0.0%)
All	All	0.35	0/13433	0.49	1/18127 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	154	LEU	CA-CB-CG	5.56	128.10	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3295	0	0	6	0
1	B	3265	0	0	4	0
1	C	3250	0	0	5	0
1	D	3295	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	37	0	0	0	0
2	B	37	0	0	0	0
2	C	37	0	0	0	0
2	D	37	0	0	0	0
3	A	20	0	0	0	0
3	B	15	0	0	0	0
3	C	15	0	0	2	0
3	D	15	0	0	1	0
4	A	63	0	0	0	0
4	B	36	0	0	0	0
4	C	54	0	0	0	0
4	D	25	0	0	0	0
All	All	13496	0	0	19	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (19) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:97:TYR:O	1:D:419:ARG:NH2	2.30	0.65
1:A:97:TYR:O	1:A:419:ARG:NH2	2.31	0.64
1:B:97:TYR:O	1:B:419:ARG:NH2	2.33	0.62
1:A:25:GLN:OE1	1:A:46:ASP:N	2.35	0.60
1:A:290:GLN:OE1	1:A:303:THR:CG2	2.50	0.59
1:B:25:GLN:OE1	1:B:46:ASP:N	2.40	0.55
1:D:164:ARG:NH1	1:D:164:ARG:CG	2.72	0.53
1:C:74:GLU:N	3:C:439:SO4:O4	2.44	0.51
1:D:74:GLU:N	3:D:440:SO4:O3	2.45	0.50
1:C:271:ASP:N	1:C:271:ASP:OD2	2.45	0.49
1:C:97:TYR:O	1:C:419:ARG:NH2	2.47	0.48
1:C:73:ARG:NE	3:C:439:SO4:O1	2.49	0.45
1:A:144:ASP:OD2	1:A:184:SER:OG	2.37	0.42
1:D:248:LYS:O	1:D:276:SER:OG	2.38	0.41
1:C:144:ASP:OD2	1:C:184:SER:OG	2.37	0.41
1:B:88:ARG:NH2	1:B:260:ASP:OD1	2.54	0.41
1:B:144:ASP:OD2	1:B:184:SER:OG	2.39	0.41
1:A:106:TYR:CZ	1:A:233:THR:OG1	2.74	0.41
1:A:300:ASP:CB	1:A:303:THR:CB	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/438 (93%)	399 (98%)	7 (2%)	1 (0%)	56	82
1	B	403/438 (92%)	392 (97%)	11 (3%)	0	100	100
1	C	401/438 (92%)	393 (98%)	8 (2%)	0	100	100
1	D	407/438 (93%)	397 (98%)	9 (2%)	1 (0%)	56	82
All	All	1618/1752 (92%)	1581 (98%)	35 (2%)	2 (0%)	59	85

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	299	SER
1	A	291	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/391 (93%)	346 (96%)	16 (4%)	39	68
1	B	358/391 (92%)	344 (96%)	14 (4%)	43	74
1	C	356/391 (91%)	336 (94%)	20 (6%)	30	55
1	D	362/391 (93%)	340 (94%)	22 (6%)	26	50
All	All	1438/1564 (92%)	1366 (95%)	72 (5%)	34	61

All (72) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	69	GLN
1	A	86	SER
1	A	149	LEU
1	A	151	LYS
1	A	179	ASN
1	A	233	THR
1	A	289	LEU
1	A	294	THR
1	A	297	SER
1	A	328	VAL
1	A	359	LEU
1	A	365	LYS
1	A	371	LEU
1	A	420	LEU
1	A	421	VAL
1	B	49	LYS
1	B	68	LYS
1	B	87	GLU
1	B	149	LEU
1	B	164	ARG
1	B	165	GLU
1	B	230	GLN
1	B	272	LYS
1	B	295	LYS
1	B	308	THR
1	B	359	LEU
1	B	371	LEU
1	B	421	VAL
1	B	424	LYS
1	C	23	LYS
1	C	60	ASN
1	C	66	ARG
1	C	68	LYS
1	C	86	SER
1	C	106	TYR
1	C	114	ASP
1	C	133	VAL
1	C	149	LEU
1	C	207	SER
1	C	271	ASP
1	C	297	SER
1	C	308	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	328	VAL
1	C	329	SER
1	C	359	LEU
1	C	371	LEU
1	C	417	LYS
1	C	420	LEU
1	C	421	VAL
1	D	20	THR
1	D	32	VAL
1	D	33	GLU
1	D	62	GLU
1	D	83	LEU
1	D	112	SER
1	D	147	LYS
1	D	149	LEU
1	D	154	LEU
1	D	164	ARG
1	D	216	ASN
1	D	233	THR
1	D	271	ASP
1	D	289	LEU
1	D	293	THR
1	D	302	THR
1	D	328	VAL
1	D	359	LEU
1	D	360	LYS
1	D	371	LEU
1	D	401	GLU
1	D	421	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	439	-	4,4,4	0.20	0	6,6,6	0.08	0
3	SO4	A	440	-	4,4,4	0.17	0	6,6,6	0.13	0
3	SO4	A	441	-	4,4,4	0.21	0	6,6,6	0.06	0
3	SO4	A	442	-	4,4,4	0.22	0	6,6,6	0.06	0
2	TYM	A	500	-	41,41,41	1.15	3 (7%)	61,61,61	1.82	8 (13%)
3	SO4	B	439	-	4,4,4	0.27	0	6,6,6	0.09	0
3	SO4	B	440	-	4,4,4	0.25	0	6,6,6	0.14	0
3	SO4	B	441	-	4,4,4	0.19	0	6,6,6	0.10	0
2	TYM	B	700	-	41,41,41	1.17	3 (7%)	61,61,61	1.93	8 (13%)
3	SO4	C	439	-	4,4,4	0.33	0	6,6,6	0.13	0
3	SO4	C	440	-	4,4,4	0.19	0	6,6,6	0.13	0
3	SO4	C	441	-	4,4,4	0.10	0	6,6,6	0.08	0
2	TYM	C	600	-	41,41,41	1.18	3 (7%)	61,61,61	1.76	7 (11%)
3	SO4	D	439	-	4,4,4	0.27	0	6,6,6	0.08	0
3	SO4	D	440	-	4,4,4	0.19	0	6,6,6	0.08	0
3	SO4	D	441	-	4,4,4	0.19	0	6,6,6	0.13	0
2	TYM	D	800	-	41,41,41	1.17	3 (7%)	61,61,61	1.69	8 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	439	-	-	0/0/0/0	0/0/0/0
3	SO4	A	440	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	441	-	-	0/0/0/0	0/0/0/0
3	SO4	A	442	-	-	0/0/0/0	0/0/0/0
2	TYM	A	500	-	-	0/21/39/39	0/1/5/5
3	SO4	B	439	-	-	0/0/0/0	0/0/0/0
3	SO4	B	440	-	-	0/0/0/0	0/0/0/0
3	SO4	B	441	-	-	0/0/0/0	0/0/0/0
2	TYM	B	700	-	-	0/21/39/39	0/1/5/5
3	SO4	C	439	-	-	0/0/0/0	0/0/0/0
3	SO4	C	440	-	-	0/0/0/0	0/0/0/0
3	SO4	C	441	-	-	0/0/0/0	0/0/0/0
2	TYM	C	600	-	-	0/21/39/39	0/1/5/5
3	SO4	D	439	-	-	0/0/0/0	0/0/0/0
3	SO4	D	440	-	-	0/0/0/0	0/0/0/0
3	SO4	D	441	-	-	0/0/0/0	0/0/0/0
2	TYM	D	800	-	-	0/21/39/39	0/1/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	600	TYM	O4'-C1'	3.34	1.46	1.41
2	D	800	TYM	O4'-C1'	3.16	1.46	1.41
2	B	700	TYM	O4'-C1'	3.15	1.46	1.41
2	A	500	TYM	C4-N9	-3.13	1.33	1.37
2	C	600	TYM	C4-N9	-3.11	1.33	1.37
2	B	700	TYM	C4-N9	-3.09	1.33	1.37
2	A	500	TYM	O4'-C1'	2.90	1.45	1.41
2	D	800	TYM	C4-N9	-2.85	1.33	1.37
2	D	800	TYM	CD1-CG	-2.61	1.33	1.37
2	A	500	TYM	CD1-CG	-2.57	1.33	1.37
2	B	700	TYM	CD1-CG	-2.51	1.33	1.37
2	C	600	TYM	CD1-CG	-2.36	1.34	1.37

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	600	TYM	N3-C2-N1	-9.01	121.18	128.71
2	B	700	TYM	N3-C2-N1	-8.81	121.35	128.71
2	A	500	TYM	N3-C2-N1	-8.64	121.48	128.71
2	D	800	TYM	N3-C2-N1	-8.33	121.74	128.71
2	B	700	TYM	O4'-C1'-N9	-7.10	101.83	108.44
2	A	500	TYM	O4'-C1'-N9	-5.38	103.44	108.44
2	B	700	TYM	OPP-C-O	4.65	125.79	122.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	800	TYM	OPP-C-O	4.60	125.75	122.11
2	B	700	TYM	N3-C4-N9	4.30	133.20	125.43
2	D	800	TYM	N3-C4-N9	4.27	133.15	125.43
2	A	500	TYM	OPP-C-O	4.25	125.47	122.11
2	C	600	TYM	N3-C4-N9	4.21	133.03	125.43
2	A	500	TYM	N3-C4-N9	4.18	132.99	125.43
2	C	600	TYM	OPP-C-O	4.14	125.39	122.11
2	C	600	TYM	O4'-C1'-N9	-3.61	105.09	108.44
2	D	800	TYM	O4'-C1'-N9	-3.22	105.44	108.44
2	D	800	TYM	C5-C4-N3	-2.83	119.53	125.70
2	B	700	TYM	C5-C4-N3	-2.71	119.80	125.70
2	C	600	TYM	C5-C4-N3	-2.69	119.85	125.70
2	A	500	TYM	C5-C4-N3	-2.61	120.02	125.70
2	D	800	TYM	C4-C5-N7	-2.57	107.33	109.52
2	A	500	TYM	C4-C5-N7	-2.44	107.43	109.52
2	C	600	TYM	C2-N3-C4	2.32	120.60	114.01
2	C	600	TYM	C4-C5-N7	-2.26	107.58	109.52
2	A	500	TYM	C8-N9-C4	2.25	108.61	106.90
2	B	700	TYM	C2-N3-C4	2.23	120.36	114.01
2	D	800	TYM	C2-N3-C4	2.21	120.30	114.01
2	A	500	TYM	C2-N3-C4	2.19	120.24	114.01
2	B	700	TYM	C4-C5-N7	-2.12	107.71	109.52
2	B	700	TYM	C2'-C1'-N9	2.07	118.58	113.27
2	D	800	TYM	O1P-P-OPP	2.00	111.19	104.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	409/438 (93%)	-0.30	8 (1%) 62 60	31, 44, 78, 124	0
1	B	405/438 (92%)	-0.11	9 (2%) 59 56	30, 54, 87, 132	0
1	C	403/438 (92%)	-0.18	10 (2%) 54 52	32, 47, 75, 135	0
1	D	409/438 (93%)	-0.14	11 (2%) 52 49	38, 55, 87, 138	0
All	All	1626/1752 (92%)	-0.18	38 (2%) 57 54	30, 50, 84, 138	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	21	ASP	5.0
1	A	301	ASP	4.9
1	C	38	GLU	4.0
1	A	302	THR	3.8
1	C	39	GLN	3.8
1	B	22	VAL	3.7
1	D	302	THR	3.5
1	D	425	PRO	3.5
1	C	40	GLY	3.4
1	C	302	THR	3.2
1	C	41	ARG	3.2
1	D	424	LYS	3.1
1	D	413	VAL	2.9
1	D	18	LYS	2.8
1	C	36	VAL	2.8
1	A	300	ASP	2.7
1	A	17	LEU	2.7
1	B	300	ASP	2.7
1	B	39	GLN	2.6
1	D	420	LEU	2.6
1	B	37	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	18	LYS	2.5
1	D	72	GLY	2.5
1	C	300	ASP	2.5
1	C	425	PRO	2.4
1	A	33	GLU	2.4
1	A	22	VAL	2.4
1	A	21	ASP	2.4
1	B	40	GLY	2.4
1	D	68	LYS	2.3
1	D	69	GLN	2.2
1	D	21	ASP	2.2
1	C	312	LYS	2.2
1	C	301	ASP	2.2
1	B	69	GLN	2.2
1	B	36	VAL	2.1
1	D	102	PRO	2.1
1	B	38	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	A	440	5/5	0.23	7.51	47,57,57,57	5
3	SO4	A	439	5/5	0.26	5.22	83,90,91,91	5
3	SO4	B	439	5/5	0.24	3.74	64,67,67,67	5
3	SO4	D	439	5/5	0.24	1.97	75,80,81,81	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	SO4	C	439	5/5	0.16	1.41	47,48,49,49	5
3	SO4	B	440	5/5	0.15	0.04	45,55,56,56	5
3	SO4	C	440	5/5	0.16	-0.26	44,54,54,54	5
3	SO4	C	441	5/5	0.13	-0.42	33,33,34,37	5
3	SO4	B	441	5/5	0.11	-0.59	48,55,56,56	5
3	SO4	A	442	5/5	0.14	-0.92	48,55,55,56	5
3	SO4	D	441	5/5	0.13	-0.93	48,54,55,55	5
2	TYM	C	600	37/37	0.12	-1.21	29,31,31,31	0
3	SO4	D	440	5/5	0.18	-1.30	46,55,55,55	5
2	TYM	B	700	37/37	0.11	-1.53	32,36,37,37	0
2	TYM	D	800	37/37	0.11	-1.55	37,39,40,40	0
2	TYM	A	500	37/37	0.11	-1.64	27,29,30,31	0
3	SO4	A	441	5/5	0.16	-2.97	47,54,54,55	5

6.5 Other polymers ⓘ

There are no such residues in this entry.