



Full wwPDB X-ray Structure Validation Report (i)

Sep 29, 2014 – 06:11 PM EDT

PDB ID : 4KVG

Title : Crystal structure of RIAM RA-PH domains in complex with GTP bound Rap1

Authors : Zhang, H.; Chang, Y.E.; Brennan, M.L.; Wu, J.

Deposited on : 2013-05-22

Resolution : 1.65 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.16 November 2013

Xtriage (Phenix) : dev-1439

EDS : stable23828

Percentile statistics : 21963

Refmac : 5.8.0049

CCP4 : 6.3.0 (Settle)

Ideal geometry (proteins) : Engh & Huber (2001)


Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)

Validation Pipeline (wwPDB-VP) : stable23828

1 Overall quality at a glance (i)


The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1404 (1.68-1.64)
Clashscore	79885	1001 (1.66-1.66)
Ramachandran outliers	78287	1581 (1.68-1.64)
Sidechain outliers	78261	1580 (1.68-1.64)
RSRZ outliers	66119	1404 (1.68-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7731 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ras-related protein Rap-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	165	1378	868	237	263	10	0	10	0
1	C	160	1282	804	219	250	9	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	EXPRESSION TAG	UNP P62834
A	12	VAL	GLY	ENGINEERED MUTATION	UNP P62834
A	63	GLU	GLN	ENGINEERED MUTATION	UNP P62834
C	0	HIS	-	EXPRESSION TAG	UNP P62834
C	12	VAL	GLY	ENGINEERED MUTATION	UNP P62834
C	63	GLU	GLN	ENGINEERED MUTATION	UNP P62834


- Molecule 2 is a protein called Amyloid beta A4 precursor protein-binding family B member 1-interacting protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	246	2138	1384	349	388	17	0	17	0
2	D	247	2125	1373	348	387	17	0	12	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	178	MET	-	EXPRESSION TAG	UNP Q8R5A3
D	178	MET	-	EXPRESSION TAG	UNP Q8R5A3

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	32	10	5	14	3	0	0
3	C	1	32	10	5	14	3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	113	Total O 113 113	0	0
6	B	290	Total O 296 296	0	6
6	C	33	Total O 33 33	0	0
6	D	289	Total O 292 292	0	3

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ras-related protein Rap-1A

Chain A: 




- Molecule 1: Ras-related protein Rap-1A

Chain C: 




- Molecule 2: Amyloid beta A4 precursor protein-binding family B member 1-interacting protein

Chain B: 



- Molecule 2: Amyloid beta A4 precursor protein-binding family B member 1-interacting protein

Chain D: 



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	82.47 Å 85.85 Å 160.85 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.82 – 1.65 47.82 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.82-1.65) 99.3 (47.82-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.67 (at 1.65 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R , R_{free}	0.197 , 0.219 0.196 , 0.219	Depositor DCC
R_{free} test set	6891 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	19.0	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.9	EDS
Estimated twinning fraction	0.002 for k,h,-l	Xtriage
L-test for twinning	$< L > = 0.49$, $< L^2 > = 0.33$	Xtriage
Outliers	5 of 137213 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7731	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.35	1/1418 (0.1%)	0.50	0/1904
1	C	0.33	0/1299	0.47	0/1748
2	B	0.45	2/2233 (0.1%)	0.58	0/3005
2	D	0.44	1/2209 (0.0%)	0.58	0/2971
All	All	0.41	4/7159 (0.1%)	0.55	0/9628

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	250	TRP	CD2-CE2	5.24	1.47	1.41
1	A	138	TRP	CD2-CE2	5.10	1.47	1.41
2	D	330	TRP	CD2-CE2	5.08	1.47	1.41
2	B	330	TRP	CD2-CE2	5.07	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1378	0	1407	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1282	0	1282	9	0
2	B	2138	0	2185	24	0
2	D	2125	0	2155	21	0
3	A	32	0	12	0	0
3	C	32	0	12	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	B	4	0	6	0	0
5	D	4	0	6	0	0
6	A	113	0	0	0	0
6	B	296	0	0	3	0
6	C	33	0	0	0	0
6	D	292	0	0	1	0
All	All	7731	0	7065	60	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (60) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:311[B]:ILE:HD11	2:B:424[B]:LYS:CG	1.66	1.23
2:B:311[B]:ILE:HD11	2:B:424[B]:LYS:HG2	1.19	1.11
2:B:311[B]:ILE:CD1	2:B:424[B]:LYS:HG3	1.79	1.10
2:D:342[B]:ILE:HG22	2:D:360:ILE:HG22	1.34	1.10
2:B:311[B]:ILE:CD1	2:B:424[B]:LYS:CG	2.33	1.06
2:B:360[B]:ILE:HD11	2:B:400[B]:LYS:HG3	1.50	0.93
2:B:311[B]:ILE:HD13	2:B:424[B]:LYS:HG3	1.49	0.93
2:B:352:THR:HG21	2:B:355:ASP:HB3	1.59	0.84
2:B:311[B]:ILE:CD1	2:B:424[B]:LYS:HG2	2.07	0.72
2:B:227[B]:ILE:HD11	2:B:257:LYS:NZ	2.07	0.69
2:D:360:ILE:HD12	2:D:400[B]:LYS:HD2	1.77	0.65
1:A:150[A]:SER:O	1:A:151[A]:LYS:HB2	1.96	0.65
2:D:342[B]:ILE:CG2	2:D:360:ILE:HG22	2.20	0.65
2:B:311[B]:ILE:HD11	2:B:424[B]:LYS:HG3	1.47	0.59
1:A:9[A]:LEU:HD21	1:A:71:TYR:CE1	2.39	0.58
2:B:360[B]:ILE:HD11	2:B:400[B]:LYS:CG	2.30	0.57
1:A:24:VAL:HA	1:A:42[A]:LYS:HD2	1.87	0.57
2:D:359:PHE:CZ	2:D:400[B]:LYS:HD3	2.41	0.56
1:A:100:ILE:O	1:A:104:LYS:HG2	2.05	0.56
2:D:336:LEU:HD13	2:D:345:VAL:HG21	1.88	0.55
2:D:227[B]:ILE:HD11	2:D:257:LYS:HE3	1.89	0.55
1:C:80:LEU:HB3	1:C:93:LEU:HD23	1.88	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:313[B]:VAL:HG23	6:B:801:HOH:O	2.07	0.54
2:D:227[B]:ILE:HD11	2:D:257:LYS:CE	2.38	0.54
2:D:278:LEU:O	2:D:279:ASP:HB3	2.09	0.52
2:D:360:ILE:HD11	2:D:365:VAL:HG11	1.91	0.52
1:C:89:THR:O	1:C:93:LEU:HD12	2.10	0.51
2:B:227[B]:ILE:HD11	2:B:257:LYS:HZ3	1.75	0.51
1:A:97[B]:ARG:HD2	1:A:138:TRP:CD2	2.46	0.50
2:B:352:THR:CB	2:B:353:SER:HB3	2.41	0.50
1:C:113:LEU:HD23	1:C:138:TRP:CZ3	2.46	0.50
2:B:345:VAL:HG22	6:B:712:HOH:O	2.12	0.49
2:B:227[B]:ILE:HD11	2:B:257:LYS:HZ1	1.74	0.49
2:B:352:THR:HB	2:B:353:SER:C	2.32	0.49
1:A:24:VAL:HA	1:A:42[B]:LYS:HD2	1.95	0.48
2:B:352:THR:HG21	2:B:355:ASP:CB	2.39	0.48
1:C:82:TYR:HE1	1:C:113:LEU:CD1	2.27	0.48
2:D:336:LEU:HD13	2:D:345:VAL:CG2	2.43	0.47
1:C:113:LEU:HD12	1:C:143:PHE:CD2	2.49	0.47
2:D:227[B]:ILE:HG22	2:D:234:GLU:HB3	1.98	0.45
2:D:349:LYS:HE3	2:D:352:THR:HG21	1.98	0.45
2:B:352:THR:HB	2:B:353:SER:HB3	1.98	0.45
1:C:89:THR:O	1:C:93:LEU:CD1	2.65	0.45
1:C:82:TYR:HE1	1:C:113:LEU:HD11	1.82	0.45
2:D:321:TYR:HB2	2:D:403:CYS:HB3	1.99	0.45
2:D:360:ILE:HG21	2:D:387:LEU:HD13	1.99	0.44
2:B:189:ASP:O	2:B:190:SER:HB2	2.19	0.43
2:B:376:LYS:NZ	6:B:808:HOH:O	2.37	0.43
2:B:321:TYR:HB2	2:B:403:CYS:HB3	2.00	0.43
2:D:395:GLU:HG2	6:D:847:HOH:O	2.18	0.43
2:B:360[A]:ILE:HG21	2:B:387:LEU:HD13	2.02	0.42
2:D:311:ILE:HD13	2:D:423:GLY:HA2	2.01	0.42
2:B:232:GLN:OE1	2:B:376:LYS:HD3	2.20	0.42
1:C:113:LEU:HD12	1:C:143:PHE:HD2	1.85	0.41
2:D:342[A]:ILE:HG13	2:D:362:PHE:CZ	2.55	0.41
1:C:117:LYS:HG2	3:C:201:GTP:C6	2.56	0.40
2:D:209:ASN:OD1	2:D:213:LYS:HE2	2.20	0.40
2:D:261:LEU:HD13	2:D:263:LYS:HG2	2.03	0.40
2:D:225:TYR:CZ	2:D:263:LYS:HE3	2.57	0.40
2:D:395:GLU:O	2:D:396:SER:HB3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	171/168 (102%)	168 (98%)	3 (2%)	0	100 100
1	C	157/168 (94%)	154 (98%)	3 (2%)	0	100 100
2	B	259/260 (100%)	255 (98%)	4 (2%)	0	100 100
2	D	256/260 (98%)	250 (98%)	6 (2%)	0	100 100
All	All	843/856 (98%)	827 (98%)	16 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	157/150 (105%)	157 (100%)	0	100 100
1	C	144/150 (96%)	143 (99%)	1 (1%)	91 80
2	B	243/239 (102%)	242 (100%)	1 (0%)	95 91
2	D	240/239 (100%)	239 (100%)	1 (0%)	95 91
All	All	784/778 (101%)	781 (100%)	3 (0%)	95 91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	351	LYS
1	C	69	ASP
2	D	261	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GTP	A	201	4	34,34,34	1.29	5 (14%)	52,54,54	2.70	8 (15%)
5	EDO	B	501	-	3,3,3	0.50	0	2,2,2	0.23	0
3	GTP	C	201	4	34,34,34	1.26	5 (14%)	52,54,54	2.99	9 (17%)
5	EDO	D	501	-	3,3,3	0.49	0	2,2,2	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTP	A	201	4	-	0/22/38/38	0/3/3/3
5	EDO	B	501	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTP	C	201	4	-	0/22/38/38	0/3/3/3
5	EDO	D	501	-	-	0/1/1/1	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	201	GTP	C4-N9	-3.33	1.32	1.37
3	C	201	GTP	C5-C4	3.26	1.47	1.40
3	A	201	GTP	C5-C4	3.06	1.47	1.40
3	A	201	GTP	C2-N2	2.73	1.36	1.32
3	C	201	GTP	C4-N9	-2.66	1.33	1.37
3	A	201	GTP	C5-N7	-2.55	1.35	1.38
3	C	201	GTP	C2-N2	2.44	1.36	1.32
3	A	201	GTP	C2-N3	2.35	1.36	1.33
3	C	201	GTP	C5-N7	-2.35	1.35	1.38
3	C	201	GTP	C2-N3	2.34	1.36	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	201	GTP	C6-C5-N7	17.50	136.50	134.14
3	A	201	GTP	C6-C5-N7	15.45	136.22	134.14
3	C	201	GTP	C5-C4-N3	-6.16	118.97	126.07
3	C	201	GTP	N3-C4-N9	5.32	134.72	126.91
3	A	201	GTP	C5-C4-N3	-5.29	119.97	126.07
3	A	201	GTP	C6-N1-C2	4.77	122.89	120.20
3	A	201	GTP	N3-C4-N9	4.75	133.88	126.91
3	C	201	GTP	C2-N3-C4	4.71	120.95	115.30
3	C	201	GTP	C6-N1-C2	4.12	122.53	120.20
3	A	201	GTP	C2-N3-C4	3.97	120.06	115.30
3	C	201	GTP	C4-C5-N7	-2.91	106.60	109.41
3	C	201	GTP	PA-O3A-PB	-2.61	124.69	131.93
3	A	201	GTP	C4-C5-N7	-2.53	106.97	109.41
3	A	201	GTP	C8-N9-C4	2.43	108.94	106.96
3	A	201	GTP	PA-O3A-PB	-2.31	125.52	131.93
3	C	201	GTP	PB-O3B-PG	-2.14	126.01	131.93
3	C	201	GTP	C3'-C2'-C1'	2.05	104.13	100.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/168 (98%)	0.04	8 (4%) 29 27	15, 25, 45, 64	0
1	C	160/168 (95%)	1.15	43 (26%) 1 1	25, 54, 98, 113	0
2	B	246/260 (94%)	0.46	15 (6%) 21 18	11, 18, 41, 78	0
2	D	247/260 (95%)	0.45	19 (7%) 13 12	10, 19, 39, 71	0
All	All	818/856 (95%)	0.51	85 (10%) 7 5	10, 22, 75, 113	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	348	GLY	9.9
2	D	353	SER	9.4
2	B	350	THR	9.1
2	B	353	SER	8.0
1	C	108	ASP	7.7
1	A	66	ALA	6.9
2	D	354	ARG	6.1
1	C	138	TRP	6.1
1	C	139	CYS	6.0
2	B	354	ARG	6.0
2	D	398	TYR	5.9
2	B	349	LYS	5.9
1	C	29	VAL	5.5
2	B	347	LYS	5.4
1	C	101	LEU	5.2
1	A	67	MET	5.1
1	C	70	LEU	5.1
1	C	106	THR	4.9
1	C	109	VAL	4.8
2	B	352	THR	4.8
2	D	347	LYS	4.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	352	THR	4.7
2	B	178	MET	4.6
1	C	104	LYS	4.5
1	C	105	ASP	4.4
1	C	100	ILE	4.4
2	B	355	ASP	4.2
1	C	31	LYS	4.2
1	A	61	THR	4.1
2	B	346	PRO	4.0
2	D	279	ASP	4.0
2	B	351	LYS	4.0
1	C	122	ASP	3.9
1	C	61	THR	3.7
2	B	345	VAL	3.6
1	C	141	CYS	3.5
1	C	103	VAL	3.5
1	C	134	LEU	3.5
2	D	178	MET	3.5
2	D	351	LYS	3.4
2	B	252	ARG	3.4
1	C	140	ASN	3.3
1	C	137	GLN	3.3
2	D	344[A]	TYR	3.2
1	C	93	LEU	3.1
1	C	121	GLU	3.1
2	D	392[A]	ILE	3.0
2	D	355	ASP	2.9
1	C	90	PHE	2.9
1	C	120	LEU	2.8
1	C	95	ASP	2.8
1	C	9	LEU	2.7
1	C	88	SER	2.7
2	D	349	LYS	2.7
1	A	65	THR	2.6
1	C	129	GLU	2.6
2	D	346	PRO	2.6
1	C	85	THR	2.6
1	C	133	ASN	2.6
1	C	71	TYR	2.5
1	C	125	VAL	2.5
2	D	278	LEU	2.5
1	C	107	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	136	ARG	2.4
1	C	92	ASP	2.4
1	C	113	LEU	2.4
2	B	315	GLU	2.3
1	C	2	ARG	2.3
1	C	98	GLU	2.3
1	C	84	ILE	2.3
1	C	118	CYS	2.3
2	D	397	GLN	2.3
2	D	348	GLY	2.3
1	A	108	ASP	2.3
1	A	95	ASP	2.3
1	C	102	ARG	2.2
2	D	395	GLU	2.2
1	A	167	ARG	2.2
2	D	253	ASP	2.2
2	D	350	THR	2.2
1	C	99	GLN	2.1
2	B	278	LEU	2.1
1	C	73	LYS	2.1
1	A	71	TYR	2.0
1	C	86	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	EDO	B	501	4/4	0.09	1.62	18,19,19,20	0
5	EDO	D	501	4/4	0.12	0.96	19,19,20,22	0
3	GTP	C	201	32/32	0.12	-0.13	29,38,39,39	0
3	GTP	A	201	32/32	0.07	-0.42	15,19,20,21	0
4	MG	A	202	1/1	0.07	-0.71	15,15,15,15	0
4	MG	C	202	1/1	0.07	-0.80	24,24,24,24	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.