



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 07:54 AM GMT

PDB ID : 1L1D  
Title : Crystal structure of the C-terminal methionine sulfoxide reductase domain (MsrB) of *N. gonorrhoeae* pilB  
Authors : Lowther, W.T.; Weissbach, H.; Etienne, F.; Brot, N.; Matthews, B.W.  
Deposited on : 2002-02-15  
Resolution : 1.85 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1269 (1.86-1.86)
Clashscore	79885	1470 (1.86-1.86)
Ramachandran outliers	78287	1451 (1.86-1.86)
Sidechain outliers	78261	1451 (1.86-1.86)
RSRZ outliers	66119	1269 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	152	
1	B	152	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	CAC	A	601	X	-
2	CAC	B	602	X	-
2	CAC	B	603	X	X

## 2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 2454 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called peptide methionine sulfoxide reductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	144	Total	C	N	O	S	Se	0	2	0
			1149	721	197	226	3	2			
1	B	147	Total	C	N	O	S	Se	0	2	0
			1166	731	200	230	3	2			

- Molecule 2 is CACODYLATE ION (three-letter code: CAC) (formula: C<sub>2</sub>H<sub>6</sub>AsO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	As	C	O	0	0
			5	1	2	2		
2	B	1	Total	As	C	O	0	0
			5	1	2	2		
2	B	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	67	Total 67	O 67	0	0
3	B	57	Total 57	O 57	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: peptide methionine sulfoxide reductase

Chain A: 



- Molecule 1: peptide methionine sulfoxide reductase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.35Å 68.08Å 62.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.20 – 1.85 27.20 – 1.85	Depositor EDS
% Data completeness (in resolution range)	(Not available) (27.20-1.85) 87.9 (27.20-1.85)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.12 (at 1.85Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.207 , 0.237 0.215 , 0.208	Depositor DCC
$R_{free}$ test set	2256 reflections (11.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.2	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.45 , 43.5	EDS
Estimated twinning fraction	0.029 for k,h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 22213 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2454	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/1177	0.58	0/1587
1	B	0.34	0/1194	0.57	0/1611
All	All	0.34	0/2371	0.57	0/3198

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1149	0	1079	18	0
1	B	1166	0	1096	12	0
2	A	5	0	0	1	0
2	B	10	0	0	0	0
3	A	67	0	0	0	0
3	B	57	0	0	0	0
All	All	2454	0	2175	29	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (29) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:502:LYS:HE3	1:A:502:LYS:HA	1.58	0.84
1:B:396:GLN:NE2	1:B:400:ASN:HD22	1.87	0.72
1:B:396:GLN:HE22	1:B:400:ASN:HD22	1.39	0.70
1:A:382:SER:OG	1:A:385:GLU:HG3	1.95	0.67
1:B:450:ASP:HB3	1:B:453:SER:OG	1.96	0.65
1:A:493:ARG:NH1	1:A:495[B]:CYS:SG	2.74	0.60
1:B:440:CYS:HG	1:B:495[B]:CYS:HG	1.30	0.58
1:A:508:GLN:HG2	1:A:508:GLN:O	2.04	0.57
1:A:443:PRO:HD2	1:A:496:ILE:O	2.06	0.55
1:A:404:GLU:HG2	1:A:497:ASN:OD1	2.06	0.55
1:B:443:PRO:HD2	1:B:496:ILE:O	2.08	0.53
1:A:378:TYR:O	1:A:473:ALA:HB1	2.09	0.53
1:A:502:LYS:HA	1:A:502:LYS:CE	2.36	0.52
1:A:432:SER:OG	1:A:521:VAL:HG13	2.11	0.50
1:B:464:MSE:HG3	1:B:466:ARG:CD	2.42	0.50
1:B:444:SER:HA	1:B:494:TYR:O	2.13	0.48
1:B:493:ARG:NH1	1:B:495[B]:CYS:SG	2.85	0.48
1:A:380:LYS:HD2	1:A:474:ALA:O	2.14	0.46
1:A:486:PRO:HG3	1:B:437:ASP:OD1	2.17	0.45
1:B:516:ALA:C	1:B:517:LEU:HD12	2.39	0.43
1:A:487:ARG:HA	1:A:487:ARG:HE	1.82	0.43
1:A:421:VAL:HG12	1:A:428:PRO:HA	2.01	0.43
1:A:507:GLU:CD	1:A:507:GLU:H	2.23	0.42
1:B:464:MSE:HG3	1:B:466:ARG:HD2	2.02	0.41
1:A:480:HIS:ND1	2:A:601:CAC:O1	2.47	0.41
1:B:425:SER:HB3	1:B:474:ALA:HB2	2.02	0.41
1:A:450:ASP:HB3	1:A:453:SER:OG	2.21	0.41
1:A:416:LYS:HE2	1:A:416:LYS:HB3	1.77	0.41
1:A:463:ASN:O	1:A:464:MSE:HB3	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/152 (95%)	139 (96%)	4 (3%)	1 (1%)	30	12
1	B	147/152 (97%)	143 (97%)	4 (3%)	0	100	100
All	All	291/304 (96%)	282 (97%)	8 (3%)	1 (0%)	50	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	462	PHE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/124 (99%)	122 (99%)	1 (1%)	89	85
1	B	124/124 (100%)	123 (99%)	1 (1%)	89	85
All	All	247/248 (100%)	245 (99%)	2 (1%)	89	85

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	487	ARG
1	B	466	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	508	GLN
1	B	396	GLN
1	B	463	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CAC	A	601	-	4,4,4	6.48	4 (100%)	6,6,6	11.68	4 (66%)
2	CAC	B	602	-	4,4,4	6.06	4 (100%)	6,6,6	12.71	4 (66%)
2	CAC	B	603	-	4,4,4	6.11	4 (100%)	6,6,6	12.16	4 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CAC	A	601	-	-	0/0/0/0	0/0/0/0
2	CAC	B	602	-	-	0/0/0/0	0/0/0/0
2	CAC	B	603	-	-	0/0/0/0	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	CAC	O2-AS	10.56	1.80	1.68
2	B	602	CAC	O2-AS	9.27	1.78	1.68
2	B	603	CAC	O2-AS	9.03	1.78	1.68
2	B	603	CAC	AS-C1	5.43	1.99	1.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	602	CAC	AS-C1	4.88	1.98	1.90
2	B	602	CAC	O1-AS	4.82	1.82	1.66
2	B	603	CAC	O1-AS	4.73	1.82	1.66
2	A	601	CAC	O1-AS	4.56	1.81	1.66
2	A	601	CAC	AS-C1	4.38	1.98	1.90
2	A	601	CAC	AS-C2	4.05	1.97	1.90
2	B	603	CAC	AS-C2	3.96	1.97	1.90
2	B	602	CAC	AS-C2	3.72	1.96	1.90

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	602	CAC	O2-AS-O1	30.39	119.09	112.54
2	B	603	CAC	O2-AS-O1	28.97	118.78	112.54
2	A	601	CAC	O2-AS-O1	27.59	118.49	112.54
2	B	602	CAC	O2-AS-C1	4.97	113.22	109.15
2	B	603	CAC	O2-AS-C1	4.89	113.16	109.15
2	A	601	CAC	O2-AS-C2	4.76	113.05	109.15
2	A	601	CAC	O2-AS-C1	4.43	112.78	109.15
2	B	603	CAC	C2-AS-C1	-3.76	100.84	109.43
2	B	602	CAC	C2-AS-C1	-3.66	101.06	109.43
2	A	601	CAC	C2-AS-C1	-3.31	101.88	109.43
2	B	603	CAC	O2-AS-C2	2.62	111.30	109.15
2	B	602	CAC	O2-AS-C2	2.05	110.83	109.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	144/152 (94%)	0.31	10 (6%) 17 15	10, 17, 35, 56	0
1	B	147/152 (96%)	0.60	17 (11%) 5 5	10, 18, 37, 50	0
All	All	291/304 (95%)	0.45	27 (9%) 9 9	10, 17, 39, 56	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	462	PHE	8.9
1	B	460	PHE	5.6
1	B	458	ASP	4.9
1	A	462	PHE	4.6
1	B	376	ALA	3.7
1	B	465	ARG	3.6
1	B	457	HIS	3.5
1	A	463	ASN	3.4
1	A	378	TYR	3.2
1	B	451	ALA	2.9
1	B	463	ASN	2.9
1	A	379	LYS	2.8
1	A	464	MSE	2.8
1	B	466	ARG	2.8
1	B	461	SER	2.8
1	A	389	THR	2.6
1	A	465	ARG	2.5
1	B	495[A]	CYS	2.5
1	A	488	ASP	2.5
1	B	464	MSE	2.5
1	B	379	LYS	2.4
1	B	496	ILE	2.3
1	B	375	ALA	2.3
1	A	487	ARG	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	521	VAL	2.1
1	B	521	VAL	2.1
1	B	442	TRP	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CAC	B	603	5/5	0.24	6.04	63,63,64,64	0
2	CAC	B	602	5/5	0.13	1.63	40,40,41,42	0
2	CAC	A	601	5/5	0.13	-0.25	34,39,40,40	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.