



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 03:23 AM GMT

PDB ID : 3L6X  
Title : Crystal structure of p120 catenin in complex with E-cadherin  
Authors : Ishiyama, N.; Lee, S.-H.; Liu, S.; Li, G.-Y.; Smith, M.J.; Reichardt, L.F.; Ikura, M.  
Deposited on : 2009-12-27  
Resolution : 2.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	584	
2	B	18	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3664 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catenin delta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	434	3386	2130	612	633	11	0	3	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	319	GLY	-	EXPRESSION TAG	UNP O60716
A	320	SER	-	EXPRESSION TAG	UNP O60716
A	321	PRO	-	EXPRESSION TAG	UNP O60716
A	322	GLU	-	EXPRESSION TAG	UNP O60716
A	323	PHE	-	EXPRESSION TAG	UNP O60716
A	?	-	PRO	DELETION	UNP O60716
A	?	-	HIS	DELETION	UNP O60716
A	?	-	ALA	DELETION	UNP O60716
A	?	-	ALA	DELETION	UNP O60716
A	?	-	SER	DELETION	UNP O60716
A	?	-	CYS	DELETION	UNP O60716
A	?	-	PHE	DELETION	UNP O60716
A	?	-	GLY	DELETION	UNP O60716
A	?	-	ALA	DELETION	UNP O60716
A	?	-	LYS	DELETION	UNP O60716
A	?	-	LYS	DELETION	UNP O60716
A	?	-	GLY	DELETION	UNP O60716
A	?	-	LYS	DELETION	UNP O60716
A	?	-	ASP	SEE REMARK 999	UNP O60716
A	?	-	GLU	SEE REMARK 999	UNP O60716
A	?	-	TRP	SEE REMARK 999	UNP O60716
A	?	-	PHE	SEE REMARK 999	UNP O60716
A	?	-	SER	SEE REMARK 999	UNP O60716
A	?	-	ARG	SEE REMARK 999	UNP O60716
A	?	-	GLY	DELETION	UNP O60716
A	?	-	LYS	DELETION	UNP O60716
A	?	-	LYS	DELETION	UNP O60716

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	PRO	DELETION	UNP O60716
A	?	-	ILE	DELETION	UNP O60716
A	?	-	GLU	DELETION	UNP O60716
A	?	-	ASP	DELETION	UNP O60716
A	?	-	PRO	DELETION	UNP O60716
A	?	-	ALA	DELETION	UNP O60716
A	?	-	ASN	DELETION	UNP O60716
A	?	-	ASP	DELETION	UNP O60716
A	?	-	THR	DELETION	UNP O60716
A	?	-	VAL	DELETION	UNP O60716
A	?	-	ASP	DELETION	UNP O60716
A	?	-	PHE	DELETION	UNP O60716
A	?	-	PRO	DELETION	UNP O60716
A	?	-	LYS	DELETION	UNP O60716
A	?	-	ARG	DELETION	UNP O60716
A	932	LYS	-	SEE REMARK 999	UNP O60716
A	933	ILE	-	SEE REMARK 999	UNP O60716

- Molecule 2 is a protein called E-cadherin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	18	Total	C	N	O	0	0	0
			141	82	22	37			

- Molecule 3 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	111	Total	O	0	0
			111	111		
4	B	6	Total	O	0	0
			6	6		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Catenin delta-1

Chain A: 



#### • Molecule 2: E-cadherin

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.40Å 92.40Å 171.89Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.00 – 2.40 32.80 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.6 (32.00-2.40) 99.6 (32.80-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.86 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.4.0073	Depositor
R, $R_{free}$	0.201 , 0.234 0.201 , 0.233	Depositor DCC
$R_{free}$ test set	1707 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.7	Xtriage
Anisotropy	0.476	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.1	EDS
Estimated twinning fraction	0.015 for -h,-k,l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 33753 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3664	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/3448	0.70	6/4674 (0.1%)
2	B	0.63	0/142	0.68	0/189
All	All	0.58	0/3590	0.70	6/4863 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	545	ARG	NE-CZ-NH2	-8.96	115.82	120.30
1	A	546	ARG	NE-CZ-NH1	-8.82	115.89	120.30
1	A	549	ARG	NE-CZ-NH2	-7.72	116.44	120.30
1	A	549	ARG	NE-CZ-NH1	6.90	123.75	120.30
1	A	545	ARG	NE-CZ-NH1	5.89	123.25	120.30
1	A	701	ARG	NE-CZ-NH2	-5.55	117.52	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3386	0	3399	63	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	141	0	107	2	0
3	A	20	0	0	1	0
4	A	111	0	0	1	0
4	B	6	0	0	0	0
All	All	3664	0	3506	65	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (65) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:377:PHE:O	1:A:383:LYS:HE2	1.55	1.06
1:A:520[A]:HIS:CD2	1:A:521:ILE:H	1.77	1.02
1:A:482:HIS:HD2	1:A:484:SER:H	1.24	0.85
1:A:520[A]:HIS:HD2	1:A:521:ILE:H	1.27	0.82
1:A:666:LEU:HD22	1:A:712:LEU:HD11	1.65	0.77
1:A:789:LYS:HD2	1:A:792:GLU:OE1	1.85	0.76
1:A:791:ARG:HD3	1:A:829:LEU:HD23	1.67	0.75
1:A:736:ARG:NH2	4:A:103:HOH:O	2.16	0.74
1:A:787:ALA:HB1	1:A:829:LEU:CD1	2.17	0.74
1:A:508:TRP:O	1:A:509:GLU:HG3	1.88	0.73
1:A:546:ARG:HH11	1:A:546:ARG:HG2	1.54	0.72
1:A:701:ARG:NH2	1:A:734:ASP:OD1	2.22	0.72
1:A:565:GLU:OE1	1:A:570:ASP:OD2	2.08	0.72
1:A:814:ARG:HD2	1:A:846:ASN:O	1.89	0.72
1:A:715:ASN:HD22	1:A:717:HIS:H	1.37	0.71
2:B:767:GLN:HE21	2:B:768:ASP:H	1.39	0.70
1:A:789:LYS:CD	1:A:792:GLU:OE1	2.39	0.70
1:A:791:ARG:HB2	1:A:829:LEU:HD21	1.73	0.69
1:A:358:PRO:O	1:A:360:PRO:HD3	1.93	0.69
1:A:546:ARG:NH1	1:A:593:GLU:O	2.28	0.67
1:A:716:GLU:O	1:A:716:GLU:HG2	1.97	0.64
1:A:570:ASP:OD1	1:A:570:ASP:C	2.33	0.64
1:A:737:ASN:HB3	1:A:740:LEU:HB2	1.78	0.64
1:A:699:ALA:O	1:A:702:GLN:HG2	1.97	0.64
1:A:715:ASN:ND2	1:A:717:HIS:H	1.96	0.63
1:A:375:LEU:O	1:A:383:LYS:HG2	1.98	0.63
1:A:737:ASN:O	1:A:740:LEU:N	2.31	0.63
1:A:540:GLU:O	1:A:541:ARG:HB2	1.99	0.62
1:A:482:HIS:CD2	1:A:484:SER:H	2.12	0.61
1:A:814:ARG:CD	1:A:846:ASN:O	2.50	0.59
1:A:563[A]:GLN:CG	1:A:661:ILE:HD11	2.33	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:791:ARG:HD3	1:A:829:LEU:CD2	2.36	0.55
1:A:520[A]:HIS:CD2	1:A:521:ILE:N	2.61	0.54
1:A:482:HIS:HD2	1:A:484:SER:N	2.00	0.53
1:A:755:GLY:HA2	1:A:762:TRP:CZ2	2.44	0.53
1:A:545:ARG:NH2	3:A:2:SO4:O4	2.42	0.52
1:A:715:ASN:ND2	1:A:717:HIS:HB2	2.25	0.51
1:A:790:LEU:HD23	1:A:823:ILE:HD11	1.92	0.51
1:A:790:LEU:CD2	1:A:823:ILE:HD11	2.41	0.51
1:A:735:ALA:C	1:A:736:ARG:HG2	2.30	0.51
1:A:796:ILE:HG21	1:A:833:LEU:HD21	1.91	0.51
1:A:791:ARG:HB2	1:A:829:LEU:CD2	2.41	0.51
1:A:546:ARG:NH1	1:A:546:ARG:HG2	2.24	0.48
1:A:563[A]:GLN:NE2	1:A:657:GLU:OE2	2.47	0.47
2:B:767:GLN:NE2	2:B:768:ASP:H	2.10	0.47
1:A:466:MET:HE2	1:A:523:TRP:CD1	2.49	0.47
1:A:546:ARG:HH11	1:A:546:ARG:CG	2.18	0.46
1:A:502:ILE:O	1:A:506:SER:HB3	2.16	0.45
1:A:422:LYS:O	1:A:426:LEU:HB3	2.16	0.45
1:A:563[A]:GLN:HG2	1:A:661:ILE:HD11	1.96	0.45
1:A:808:ARG:HD3	1:A:812:GLU:OE2	2.17	0.45
1:A:814:ARG:HG3	1:A:845:VAL:HG13	1.98	0.45
1:A:831:LYS:HB2	1:A:832:PRO:HD3	1.99	0.45
1:A:453:VAL:HB	1:A:454:PRO:HD3	1.99	0.43
1:A:811:LYS:HE2	1:A:811:LYS:HB3	1.59	0.43
1:A:762:TRP:HH2	1:A:801:LEU:HD11	1.85	0.42
1:A:520[A]:HIS:CG	1:A:521:ILE:H	2.29	0.42
1:A:735:ALA:O	1:A:736:ARG:HG2	2.20	0.41
1:A:787:ALA:HB1	1:A:829:LEU:HD11	2.00	0.41
1:A:568:GLN:HA	1:A:568:GLN:OE1	2.19	0.41
1:A:794:GLN:HA	1:A:797:GLU:OE1	2.19	0.41
1:A:549:ARG:HB3	1:A:650:TYR:HB3	2.01	0.41
1:A:441:GLN:OE1	1:A:444:LYS:HD2	2.21	0.40
1:A:584[A]:ARG:HG3	1:A:681:ALA:N	2.35	0.40
1:A:830:ARG:O	1:A:834:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/584 (74%)	414 (96%)	16 (4%)	0	100	100
2	B	16/18 (89%)	16 (100%)	0	0	100	100
All	All	446/602 (74%)	430 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/503 (72%)	356 (99%)	5 (1%)	78	92
2	B	15/15 (100%)	15 (100%)	0	100	100
All	All	376/518 (73%)	371 (99%)	5 (1%)	80	93

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	540	GLU
1	A	556	ASP
1	A	570	ASP
1	A	690	TRP
1	A	736	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	391	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	482	HIS
1	A	683	GLN
1	A	684	ASN
1	A	715	ASN
1	A	730	ASN
1	A	821	GLN
2	B	767	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	1	-	4,4,4	0.21	0	6,6,6	0.43	0
3	SO4	A	2	-	4,4,4	0.23	0	6,6,6	0.51	0
3	SO4	A	3	-	4,4,4	0.11	0	6,6,6	0.14	0
3	SO4	A	4	-	4,4,4	0.15	0	6,6,6	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	1	-	-	0/0/0/0	0/0/0/0
3	SO4	A	2	-	-	0/0/0/0	0/0/0/0
3	SO4	A	3	-	-	0/0/0/0	0/0/0/0
3	SO4	A	4	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	434/584 (74%)	-0.14	16 (3%)	39 38	25, 38, 52, 65	0
2	B	18/18 (100%)	-0.49	0	100 100	30, 37, 46, 46	0
All	All	452/602 (75%)	-0.16	16 (3%)	42 40	25, 38, 52, 65	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	846	ASN	5.6
1	A	358	PRO	5.3
1	A	824	TRP	4.8
1	A	829	LEU	4.6
1	A	359	PRO	4.2
1	A	645	SER	4.0
1	A	845	VAL	4.0
1	A	826	TYR	3.3
1	A	520[A]	HIS	3.0
1	A	464	ARG	2.7
1	A	598	GLU	2.6
1	A	379	LEU	2.2
1	A	825	GLY	2.2
1	A	832	PRO	2.2
1	A	844	GLN	2.0
1	A	833	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	2	5/5	0.16	0.54	52,53,55,56	0
3	SO4	A	3	5/5	0.13	0.06	78,79,80,80	0
3	SO4	A	4	5/5	0.14	-0.71	85,86,86,87	0
3	SO4	A	1	5/5	0.11	-0.74	69,71,72,73	0

### 6.5 Other polymers ⓘ

There are no such residues in this entry.