



wwPDB X-ray Structure Validation Summary Report

Apr 7, 2014 – 03:54 PM EDT

PDB ID : 3LCM
Title : Crystal structure of Smu.1420 from Streptococcus mutans UA159
Authors : Wang, Z.X.; Su, X.-D.
Deposited on : 2010-01-11
Resolution : 1.80 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	FAILED
Xtriage (Phenix)	:	dev-1439
EDS	:	stable22978
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22978

1 Overall quality at a glance

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	196	
1	B	196	
1	C	196	
1	D	196	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NAP	C	198	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6676 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative oxidoreductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1559	1016	265	273	5			
1	B	188	Total	C	N	O	S	0	0	0
			1472	958	246	263	5			
1	C	196	Total	C	N	O	S	0	0	0
			1557	1011	263	278	5			
1	D	188	Total	C	N	O	S	0	0	0
			1456	949	240	262	5			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	189	LEU	-	EXPRESSION TAG	UNP Q8DTD1
A	190	GLU	-	EXPRESSION TAG	UNP Q8DTD1
A	191	HIS	-	EXPRESSION TAG	UNP Q8DTD1
A	192	HIS	-	EXPRESSION TAG	UNP Q8DTD1
A	193	HIS	-	EXPRESSION TAG	UNP Q8DTD1
A	194	HIS	-	EXPRESSION TAG	UNP Q8DTD1
A	195	HIS	-	EXPRESSION TAG	UNP Q8DTD1
A	196	HIS	-	EXPRESSION TAG	UNP Q8DTD1
B	189	LEU	-	EXPRESSION TAG	UNP Q8DTD1
B	190	GLU	-	EXPRESSION TAG	UNP Q8DTD1
B	191	HIS	-	EXPRESSION TAG	UNP Q8DTD1
B	192	HIS	-	EXPRESSION TAG	UNP Q8DTD1
B	193	HIS	-	EXPRESSION TAG	UNP Q8DTD1
B	194	HIS	-	EXPRESSION TAG	UNP Q8DTD1
B	195	HIS	-	EXPRESSION TAG	UNP Q8DTD1
B	196	HIS	-	EXPRESSION TAG	UNP Q8DTD1
C	189	LEU	-	EXPRESSION TAG	UNP Q8DTD1
C	190	GLU	-	EXPRESSION TAG	UNP Q8DTD1
C	191	HIS	-	EXPRESSION TAG	UNP Q8DTD1
C	192	HIS	-	EXPRESSION TAG	UNP Q8DTD1
C	193	HIS	-	EXPRESSION TAG	UNP Q8DTD1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	194	HIS	-	EXPRESSION TAG	UNP Q8DTD1
C	195	HIS	-	EXPRESSION TAG	UNP Q8DTD1
C	196	HIS	-	EXPRESSION TAG	UNP Q8DTD1
D	189	LEU	-	EXPRESSION TAG	UNP Q8DTD1
D	190	GLU	-	EXPRESSION TAG	UNP Q8DTD1
D	191	HIS	-	EXPRESSION TAG	UNP Q8DTD1
D	192	HIS	-	EXPRESSION TAG	UNP Q8DTD1
D	193	HIS	-	EXPRESSION TAG	UNP Q8DTD1
D	194	HIS	-	EXPRESSION TAG	UNP Q8DTD1
D	195	HIS	-	EXPRESSION TAG	UNP Q8DTD1
D	196	HIS	-	EXPRESSION TAG	UNP Q8DTD1

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 9	C 6	N 2	O 1		0	0
3	B	1	Total 9	C 6	N 2	O 1		0	0
3	C	1	Total 26	C 11	N 2	O 11	P 2	0	0
3	C	1	Total 9	C 6	N 2	O 1		0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	105	Total O 105 105	0	0
4	B	105	Total O 105 105	0	0
4	C	81	Total O 81 81	0	0
4	D	76	Total O 76 76	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative oxidoreductase

Chain A: 



- Molecule 1: Putative oxidoreductase

Chain B: 




- Molecule 1: Putative oxidoreductase

Chain C: 



- Molecule 1: Putative oxidoreductase

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.60Å 79.12Å 106.28Å 90.00° 90.01° 90.00°	Depositor
Resolution (Å)	29.02 – 1.80 29.02 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.8 (29.02-1.80) 94.8 (29.02-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.66 (at 1.80Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.160 , 0.188 0.162 , 0.186	Depositor DCC
R_{free} test set	1998 reflections (2.71%)	DCC
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 34.0	EDS
Estimated twinning fraction	0.425 for h,-k,-l 0.437 for h,-k,-l	Xtriage
Reported twinning fraction	0.425 for h,-k,-l	Depositor
L-test for twinning	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	2 of 73733 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6676	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	1/1605 (0.1%)	0.50	0/2188
1	B	0.47	0/1511	0.52	0/2063
1	C	0.37	0/1603	0.50	0/2187
1	D	0.47	0/1494	0.53	0/2042
All	All	0.44	1/6213 (0.0%)	0.51	0/8480

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	162	GLU	CD-OE1	-5.03	1.20	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1559	0	1511	20	0
1	B	1472	0	1434	23	0
1	C	1557	0	1491	20	0
1	D	1456	0	1403	22	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	53	0	29	1	0
2	B	53	0	28	1	0
2	C	53	0	31	2	0
2	D	53	0	29	3	0
3	A	9	0	6	2	0
3	B	9	0	6	0	0
3	C	35	0	19	5	0
4	A	105	0	0	1	0
4	B	105	0	0	1	0
4	C	81	0	0	0	0
4	D	76	0	0	7	0
All	All	6676	0	5987	88	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

The worst 5 of 88 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:191:HIS:HD2	1:A:192:HIS:HD2	1.15	0.94
1:A:191:HIS:CD2	1:A:192:HIS:HD2	1.95	0.84
1:A:143:LYS:HB2	1:B:139:GLN:HE22	1.47	0.80
1:A:191:HIS:HD2	1:A:192:HIS:CD2	2.01	0.78
1:A:143:LYS:HB2	1:B:139:GLN:NE2	1.98	0.77

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/196 (99%)	185 (95%)	9 (5%)	0	100	100
1	B	186/196 (95%)	178 (96%)	7 (4%)	1 (0%)	38	19
1	C	194/196 (99%)	180 (93%)	13 (7%)	1 (0%)	38	19
1	D	186/196 (95%)	175 (94%)	11 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	760/784 (97%)	718 (94%)	40 (5%)	2 (0%)	50 31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	140	ASP
1	B	50	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/178 (92%)	162 (99%)	1 (1%)	92	90
1	B	154/178 (86%)	152 (99%)	2 (1%)	80	71
1	C	163/178 (92%)	159 (98%)	4 (2%)	60	42
1	D	150/178 (84%)	146 (97%)	4 (3%)	57	39
All	All	630/712 (88%)	619 (98%)	11 (2%)	73	61

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	108	TYR
1	C	113	LEU
1	D	118	GLN
1	C	55	ARG
1	D	1	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	24	GLN
1	C	192	HIS
1	D	24	GLN
1	B	118	GLN
1	C	195	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

Mogul failed to run properly - this section will therefore be empty.

5.5 Carbohydrates ⓘ

Mogul failed to run properly - this section will therefore be empty.

5.6 Ligand geometry ⓘ

Mogul failed to run properly - this section will therefore be empty.

5.7 Other polymers ⓘ

Mogul failed to run properly - this section will therefore be empty.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/196 (100%)	0.17	2 (1%) 79 76	14, 19, 28, 43	0
1	B	188/196 (95%)	0.24	1 (0%) 88 87	14, 19, 30, 50	0
1	C	196/196 (100%)	0.36	3 (1%) 70 66	15, 20, 33, 50	0
1	D	188/196 (95%)	0.41	4 (2%) 60 54	16, 23, 33, 51	0
All	All	768/784 (97%)	0.29	10 (1%) 67 70	14, 20, 32, 51	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	111	VAL	6.2
1	D	111	VAL	5.8
1	D	137	PHE	4.4
1	A	108	TYR	3.4
1	C	113	LEU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAP	C	198	26/48	0.23	6.02	25,38,58,64	0
3	NAP	C	199	9/48	0.22	1.29	24,27,30,32	0
3	NAP	B	198	9/48	0.16	1.02	21,25,27,28	0
3	NAP	A	198	9/48	0.12	0.09	19,22,24,26	0
2	FAD	B	197	53/53	0.10	-0.10	14,18,22,24	0
2	FAD	A	197	53/53	0.10	-0.17	16,19,25,27	0
2	FAD	D	197	53/53	0.10	-0.59	19,21,27,31	0
2	FAD	C	197	53/53	0.10	-0.69	19,22,28,31	0

6.5 Other polymers ⓘ

There are no such residues in this entry.