



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 03:32 AM GMT

PDB ID : 1LDC
Title : X-RAY STRUCTURE OF TWO COMPLEXES OF THE Y143F FLAVO-CYTOCHROME B2 MUTANT CRYSTALLIZED IN THE PRESENCE OF LACTATE OR PHENYL-LACTATE
Authors : Tegoni, M.; Cambillau, C.
Deposited on : 1995-04-13
Resolution : 2.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	511	
1	B	511	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 8726 atoms, of which 1765 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called L-LACTATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	478	Total	C	H	N	O	S	0	0	0
			4530	2373	809	629	705	14			
1	B	382	Total	C	H	N	O	S	0	0	0
			3648	1893	666	507	571	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	143	PHE	TYR	ENGINEERED MUTATION	UNP P00175
B	143	PHE	TYR	ENGINEERED MUTATION	UNP P00175

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			35	17	4	4	9	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	H	N	O	P	0	0
			35	17	4	4	9	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Fe	H	N	O	0	0
			47	34	1	4	4	4		

- Molecule 4 is PYRUVIC ACID (three-letter code: PYR) (formula: $C_3H_4O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	130	Total	H	O	0	0
			386	256	130		
5	B	11	Total	H	O	0	0
			33	22	11		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: L-LACTATE DEHYDROGENASE

Chain A:



• Molecule 1: L-LACTATE DEHYDROGENASE

Chain B:





4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	164.50Å 164.50Å 114.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 – 2.90	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.90)	Depositor
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.195 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8726	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, HEM, PYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.08	10/3794 (0.3%)	1.95	102/5140 (2.0%)
1	B	1.00	0/3030	1.95	80/4094 (2.0%)
All	All	1.05	10/6824 (0.1%)	1.95	182/9234 (2.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	2
All	All	0	8

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	103	LYS	CD-CE	8.02	1.71	1.51
1	A	32	TYR	CE1-CZ	6.87	1.47	1.38
1	A	32	TYR	CG-CD2	6.32	1.47	1.39
1	A	32	TYR	CE2-CZ	6.13	1.46	1.38
1	A	382	ARG	CZ-NH1	5.96	1.40	1.33

The worst 5 of 182 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	463	ARG	NE-CZ-NH2	-18.93	110.83	120.30
1	B	289	ARG	NE-CZ-NH1	18.50	129.55	120.30
1	A	382	ARG	NE-CZ-NH2	-15.38	112.61	120.30
1	A	413	ARG	NE-CZ-NH2	-15.36	112.62	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	492	ASN	CA-C-N	-13.15	88.26	117.20

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	148	ALA	Peptide
1	A	32	TYR	Sidechain
1	A	34	TYR	Sidechain
1	A	433	ARG	Sidechain
1	A	492	ASN	Mainchain

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3721	809	2974	148	0
1	B	2982	666	2378	135	0
2	A	31	4	10	3	0
2	B	31	4	15	4	0
3	A	43	4	26	8	0
4	A	6	0	3	2	0
4	B	6	0	3	1	0
5	A	130	256	0	15	0
5	B	11	22	0	0	0
All	All	6961	1765	5409	279	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 20.

The worst 5 of 279 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:289:ARG:HH21	1:A:376:ARG:NH2	1.38	1.22
1:A:289:ARG:NH2	1:A:376:ARG:NH2	2.06	1.04
1:A:488:VAL:HG22	1:B:490:VAL:HG11	1.51	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:289:ARG:NH2	1:A:376:ARG:HH22	1.68	0.90
1:B:148:ALA:C	1:B:149:ASN:HD22	1.77	0.86

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/511 (93%)	425 (90%)	36 (8%)	13 (3%)	8	30
1	B	378/511 (74%)	331 (88%)	38 (10%)	9 (2%)	9	35
All	All	852/1022 (83%)	756 (89%)	74 (9%)	22 (3%)	8	32

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	PRO
1	A	104	GLU
1	A	114	SER
1	B	146	SER
1	B	149	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/440 (93%)	344 (84%)	67 (16%)	3	10
1	B	330/440 (75%)	277 (84%)	53 (16%)	3	10
All	All	741/880 (84%)	621 (84%)	120 (16%)	3	10

5 of 120 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	400	LEU
1	B	103	LYS
1	B	429	VAL
1	A	448	GLU
1	A	469	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	377	GLN
1	A	492	ASN
1	B	218	GLN
1	A	256	ASN
1	B	247	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	A	560	1	49,50,50	2.70	13 (26%)	46,82,82	2.14	9 (19%)
2	FMN	A	570	-	33,33,33	2.61	9 (27%)	46,50,50	4.10	23 (50%)
4	PYR	A	580	-	5,5,5	2.58	2 (40%)	6,6,6	1.63	1 (16%)
2	FMN	B	570	-	33,33,33	2.59	6 (18%)	46,50,50	3.13	19 (41%)
4	PYR	B	580	-	5,5,5	2.39	1 (20%)	6,6,6	1.86	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	560	1	-	0/14/114/114	0/0/8/8
2	FMN	A	570	-	2/2/4/4	0/18/18/18	0/0/3/3
4	PYR	A	580	-	-	0/4/4/4	0/0/0/0
2	FMN	B	570	-	2/2/4/4	0/18/18/18	0/0/3/3
4	PYR	B	580	-	-	0/4/4/4	0/0/0/0

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	570	FMN	C1'-N10	-11.52	1.35	1.48
2	A	570	FMN	C1'-N10	-10.32	1.36	1.48
3	A	560	HEM	CAA-C2A	9.12	1.67	1.52
3	A	560	HEM	C2D-C1D	-7.45	1.42	1.44
3	A	560	HEM	C3D-C4D	6.78	1.46	1.44

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	570	FMN	C2-N1-C10	16.02	131.12	114.98
2	A	570	FMN	C4A-C10-N1	-14.27	108.47	122.73
2	B	570	FMN	C2-N1-C10	13.59	128.68	114.98
2	B	570	FMN	C4A-C10-N1	-8.47	114.27	122.73
3	A	560	HEM	CAA-C2A-C1A	-7.79	101.98	125.50

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	570	FMN	C4'

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
2	B	570	FMN	C2'
2	A	570	FMN	C4'
2	A	570	FMN	C2'

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.