



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 12:28 AM GMT

PDB ID : 4LHD
Title : Crystal structure of Synechocystis sp. PCC 6803 glycine decarboxylase (P-protein), holo form with pyridoxal-5'-phosphate and glycine, closed flexible loop
Authors : Hasse, D.; Andersson, E.; Carlsson, G.; Masloboy, A.; Hagemann, M.; Bauwe, H.; Andersson, I.
Deposited on : 2013-07-01
Resolution : 1.80 Å (reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	983	
1	B	983	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	GLY	A	1001	-	X
2	GLY	A	1002	-	X
2	GLY	A	1003	-	X
2	GLY	A	1004	-	X
2	GLY	B	1001	-	X
3	BCT	A	1005	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
3	BCT	A	1006	-	X
3	BCT	B	1003	-	X
4	PO4	A	1008	-	X
4	PO4	A	1009	-	X
5	EDO	A	1010	-	X
5	EDO	A	1011	-	X
5	EDO	A	1012	-	X
5	EDO	A	1013	-	X
5	EDO	A	1014	-	X
5	EDO	A	1015	-	X
5	EDO	A	1016	-	X
5	EDO	A	1017	-	X
5	EDO	B	1007	-	X
5	EDO	B	1008	-	X
5	EDO	B	1009	-	X
5	EDO	B	1010	-	X

2 Entry composition i


There are 6 unique types of molecules in this entry. The entry contains 16216 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycine dehydrogenase [decarboxylating].

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	952	Total	C	N	O	P	S	0	22	0
			7424	4724	1253	1401	1	45			
1	B	940	Total	C	N	O	P	S	0	17	0
			7298	4645	1227	1380	1	45			

- Molecule 2 is GLYCINE (three-letter code: GLY) (formula: C₂H₅NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			5	2	1	2		
2	A	1	Total	C	N	O	0	0
			5	2	1	2		
2	A	1	Total	C	N	O	0	0
			5	2	1	2		
2	A	1	Total	C	N	O	0	0
			5	2	1	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			5	2	1	2		
2	B	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 3 is BICARBONATE ION (three-letter code: BCT) (formula: CHO_3).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	1	3		
3	A	1	Total	C	O	0	0
			4	1	3		
3	A	1	Total	C	O	0	0
			4	1	3		
3	B	1	Total	C	O	0	0
			4	1	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	697	Total O 698 698	0	1
6	B	675	Total O 678 678	0	3

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycine dehydrogenase [decarboxylating]

Chain A: 



- Molecule 1: Glycine dehydrogenase [decarboxylating]

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	159.99Å 159.99Å 159.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.76 – 1.80 49.76 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (49.76-1.80) 99.4 (49.76-1.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.163 , 0.191 0.159 , 0.188	Depositor DCC
R_{free} test set	10952 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	16.2	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 34.1	EDS
Estimated twinning fraction	0.012 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 217336 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16216	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BCT, PO4, LLP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/7653	0.52	0/10407
1	B	0.36	0/7509	0.52	1/10212 (0.0%)
All	All	0.35	0/15162	0.52	1/20619 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	48	MET	CG-SD-CE	-5.79	90.94	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7424	0	0	17	0
1	B	7298	0	0	17	0
2	A	20	0	8	3	0
2	B	10	0	4	1	0
3	A	12	0	0	0	0
3	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	10	0	0	0	0
4	B	10	0	0	0	0
5	A	32	0	48	1	0
5	B	20	0	30	1	0
6	A	698	0	0	8	0
6	B	678	0	0	5	0
All	All	16216	0	90	33	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (33) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:210[B]:ASN:ND2	6:B:1709:HOH:O	2.25	0.69
1:B:395:ARG:NH2	1:B:476:GLU:OE2	2.26	0.69
1:B:564[B]:GLN:NE2	6:B:1395:HOH:O	2.28	0.67
1:B:604:TYR:OH	1:B:608:ARG:NH1	2.30	0.65
1:B:89:GLN:NE2	6:B:1769:HOH:O	2.28	0.65
1:A:623:HIS:NE2	2:A:1001:GLY:HA2	2.13	0.63
1:B:160:GLU:OE1	1:B:309:ALA:N	2.33	0.61
1:A:381:ARG:NH2	1:B:24:GLU:OE2	2.35	0.60
1:A:818:GLU:OE2	1:A:826:ARG:NH2	2.35	0.59
1:B:416:GLN:NE2	6:B:1473:HOH:O	2.34	0.59
1:A:96:LYS:NZ	6:A:1670:HOH:O	2.37	0.57
1:A:89:GLN:NE2	6:A:1489:HOH:O	2.38	0.56
1:B:81:GLN:NE2	1:B:89:GLN:OE1	2.40	0.55
1:B:726:LLP:O3	1:B:726:LLP:NZ	2.40	0.55
1:A:81:GLN:NE2	1:A:89:GLN:OE1	2.40	0.54
1:A:623:HIS:N	6:A:1778:HOH:O	2.40	0.54
1:A:91:LYS:NZ	1:A:547:GLU:OE1	2.44	0.51
1:A:726:LLP:NZ	1:A:726:LLP:O3	2.40	0.50
1:A:55:ASP:N	5:A:1014:EDO:HO2	2.09	0.50
1:B:210[B]:ASN:ND2	6:B:1749:HOH:O	2.45	0.49
1:B:610:GLU:OE1	1:B:612:GLN:NE2	2.46	0.49
1:B:643:ASP:N	1:B:647:ASN:O	2.46	0.48
1:A:394:LYS:NZ	6:A:1598:HOH:O	2.47	0.48
1:B:714:ASP:OD2	5:B:1009:EDO:O1	2.24	0.46
1:A:131:TYR:OH	1:A:354:THR:OG1	2.35	0.45
1:A:416:GLN:CB	2:A:1003:GLY:HA2	2.46	0.45
1:A:24:GLU:OE2	1:B:381:ARG:NH2	2.49	0.45
1:A:647:ASN:ND2	6:A:1688:HOH:O	2.50	0.45
1:B:612:GLN:OE1	1:B:612:GLN:N	2.51	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:46:ARG:NH2	6:A:1622:HOH:O	2.53	0.41
2:A:1002:GLY:N	6:A:1551:HOH:O	2.52	0.41
1:B:155:ASP:OD1	2:B:1002:GLY:N	2.52	0.41
1:A:603:GLN:NE2	6:A:1555:HOH:O	2.52	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	969/983 (99%)	942 (97%)	26 (3%)	1 (0%)	59	41
1	B	950/983 (97%)	925 (97%)	24 (2%)	1 (0%)	59	41
All	All	1919/1966 (98%)	1867 (97%)	50 (3%)	2 (0%)	59	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	165	SER
1	B	165	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	794/798 (100%)	787 (99%)	7 (1%)	87	83
1	B	778/798 (98%)	771 (99%)	7 (1%)	87	83
All	All	1572/1596 (98%)	1558 (99%)	14 (1%)	87	83

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	A	315	MET
1	A	347	LYS
1	A	465	ASN
1	A	529	SER
1	A	755	THR
1	A	977	MET
1	B	315	MET
1	B	400	LEU
1	B	469	THR
1	B	478	LYS
1	B	529	SER
1	B	550	LYS
1	B	824	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LLP	A	726	1	24,24,25	3.87	6 (25%)	30,32,34	2.01	3 (10%)
1	LLP	B	726	1	24,24,25	3.98	5 (20%)	30,32,34	2.08	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	726	1	-	0/15/17/19	0/1/1/1
1	LLP	B	726	1	-	0/15/17/19	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	726	LLP	O-C	17.83	1.23	1.11
1	A	726	LLP	O-C	17.18	1.23	1.11
1	B	726	LLP	O3-C3	-5.40	1.24	1.37
1	A	726	LLP	O3-C3	-5.20	1.24	1.37
1	A	726	LLP	CA-C	2.52	1.53	1.48
1	B	726	LLP	C2-N1	2.51	1.38	1.33
1	A	726	LLP	C2-N1	2.45	1.38	1.33
1	B	726	LLP	CA-C	2.40	1.52	1.48
1	A	726	LLP	CB-CA	2.40	1.55	1.53
1	A	726	LLP	C4'-NZ	-2.20	1.34	1.45
1	B	726	LLP	C4'-NZ	-2.20	1.34	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	726	LLP	OP4-C5'-C5	8.84	127.25	109.26
1	A	726	LLP	OP4-C5'-C5	7.69	124.91	109.26
1	A	726	LLP	C-CA-N	-5.69	108.14	113.83
1	B	726	LLP	C-CA-N	-5.11	108.72	113.83
1	A	726	LLP	C4'-C4-C5	2.17	121.69	119.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLY	A	1001	-	4,4,4	1.03	0	4,4,4	1.20	0
2	GLY	A	1002	-	4,4,4	1.04	0	4,4,4	1.29	1 (25%)
2	GLY	A	1003	-	4,4,4	1.08	0	4,4,4	1.31	0
2	GLY	A	1004	-	4,4,4	1.01	0	4,4,4	1.40	1 (25%)
3	BCT	A	1005	-	0,3,3	0.00	-	0,3,3	0.00	-
3	BCT	A	1006	-	0,3,3	0.00	-	0,3,3	0.00	-
3	BCT	A	1007	-	0,3,3	0.00	-	0,3,3	0.00	-
4	PO4	A	1008	-	4,4,4	0.35	0	6,6,6	0.31	0
4	PO4	A	1009	-	4,4,4	0.25	0	6,6,6	0.31	0
5	EDO	A	1010	-	3,3,3	0.59	0	2,2,2	0.50	0
5	EDO	A	1011	-	3,3,3	0.53	0	2,2,2	0.24	0
5	EDO	A	1012	-	3,3,3	0.54	0	2,2,2	0.51	0
5	EDO	A	1013	-	3,3,3	0.61	0	2,2,2	0.15	0
5	EDO	A	1014	-	3,3,3	0.56	0	2,2,2	0.30	0
5	EDO	A	1015	-	3,3,3	0.56	0	2,2,2	0.40	0
5	EDO	A	1016	-	3,3,3	0.48	0	2,2,2	0.60	0
5	EDO	A	1017	-	3,3,3	0.53	0	2,2,2	0.50	0
2	GLY	B	1001	-	4,4,4	1.03	0	4,4,4	1.36	1 (25%)
2	GLY	B	1002	-	4,4,4	1.03	0	4,4,4	1.12	0
3	BCT	B	1003	-	0,3,3	0.00	-	0,3,3	0.00	-
4	PO4	B	1004	-	4,4,4	0.26	0	6,6,6	0.31	0
4	PO4	B	1005	-	4,4,4	0.24	0	6,6,6	0.31	0
5	EDO	B	1006	-	3,3,3	0.57	0	2,2,2	0.19	0
5	EDO	B	1007	-	3,3,3	0.61	0	2,2,2	0.43	0
5	EDO	B	1008	-	3,3,3	0.56	0	2,2,2	0.39	0
5	EDO	B	1009	-	3,3,3	0.52	0	2,2,2	0.50	0
5	EDO	B	1010	-	3,3,3	0.59	0	2,2,2	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLY	A	1001	-	-	0/2/2/2	0/0/0/0
2	GLY	A	1002	-	-	0/2/2/2	0/0/0/0
2	GLY	A	1003	-	-	0/2/2/2	0/0/0/0
2	GLY	A	1004	-	-	0/2/2/2	0/0/0/0
3	BCT	A	1005	-	-	0/0/0/0	0/0/0/0
3	BCT	A	1006	-	-	0/0/0/0	0/0/0/0
3	BCT	A	1007	-	-	0/0/0/0	0/0/0/0
4	PO4	A	1008	-	-	0/0/0/0	0/0/0/0
4	PO4	A	1009	-	-	0/0/0/0	0/0/0/0
5	EDO	A	1010	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1011	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1012	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1013	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1014	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1015	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1016	-	-	0/1/1/1	0/0/0/0
5	EDO	A	1017	-	-	0/1/1/1	0/0/0/0
2	GLY	B	1001	-	-	0/2/2/2	0/0/0/0
2	GLY	B	1002	-	-	0/2/2/2	0/0/0/0
3	BCT	B	1003	-	-	0/0/0/0	0/0/0/0
4	PO4	B	1004	-	-	0/0/0/0	0/0/0/0
4	PO4	B	1005	-	-	0/0/0/0	0/0/0/0
5	EDO	B	1006	-	-	0/1/1/1	0/0/0/0
5	EDO	B	1007	-	-	0/1/1/1	0/0/0/0
5	EDO	B	1008	-	-	0/1/1/1	0/0/0/0
5	EDO	B	1009	-	-	0/1/1/1	0/0/0/0
5	EDO	B	1010	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1004	GLY	OXT-C-O	-2.25	117.57	123.30
2	B	1001	GLY	OXT-C-O	-2.19	117.72	123.30
2	A	1002	GLY	OXT-C-O	-2.12	117.89	123.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	952/983 (96%)	-0.21	24 (2%)	54 47	7, 15, 37, 66	0
1	B	940/983 (95%)	-0.14	38 (4%)	36 29	8, 16, 40, 68	1 (0%)
All	All	1892/1966 (96%)	-0.18	62 (3%)	42 36	7, 16, 39, 68	1 (0%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	16	ILE	10.5
1	A	16	ILE	5.8
1	B	187	LYS	4.7
1	B	644	GLY	4.2
1	B	185	LYS	4.1
1	A	969	HIS	4.1
1	B	15	ALA	3.9
1	A	474	TRP	3.7
1	A	469	THR	3.7
1	A	17	ALA	3.6
1	B	977	MET	3.5
1	B	18	VAL	3.4
1	A	343	ILE	3.3
1	A	622	ALA	3.3
1	B	465	ASN	3.3
1	B	828	ASN	3.2
1	B	463	LYS	3.2
1	B	968	ARG	3.2
1	B	969	HIS	3.1
1	B	469	THR	3.1
1	A	470	PRO	3.0
1	A	18	VAL	3.0
1	A	755	THR	3.0
1	B	975	GLU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	22	LYS	3.0
1	B	478	LYS	2.9
1	B	17	ALA	2.9
1	B	20	LEU	2.8
1	A	477	VAL	2.8
1	B	976	GLY	2.8
1	A	15	ALA	2.7
1	B	466	LEU	2.7
1	B	470	PRO	2.7
1	A	347	LYS	2.6
1	B	19	ASP	2.5
1	A	344	ARG	2.5
1	B	313	ARG	2.4
1	B	965	TYR	2.4
1	B	474	TRP	2.4
1	B	464	ASP	2.4
1	A	22	LYS	2.4
1	A	621	SER	2.3
1	B	692	PHE	2.3
1	B	660	TYR	2.3
1	A	662	ASP	2.3
1	B	467	PRO	2.3
1	B	327	HIS	2.3
1	A	467	PRO	2.2
1	A	471	GLU	2.2
1	B	473	LEU	2.2
1	A	475	SER	2.2
1	B	23	LEU	2.1
1	A	660	TYR	2.1
1	B	477	VAL	2.1
1	B	662	ASP	2.1
1	B	461	ALA	2.1
1	A	19	ASP	2.1
1	B	181	TYR	2.1
1	B	479	THR	2.1
1	A	658	GLU	2.0
1	A	349	THR	2.0
1	B	622	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	LLP	B	726	24/25	0.14	2.71	11,27,36,40	11
1	LLP	A	726	24/25	0.09	0.26	8,20,32,36	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GLY	A	1003	5/5	0.26	19.45	35,35,41,42	5
5	EDO	B	1010	4/4	0.24	15.64	34,35,39,42	4
3	BCT	A	1006	4/4	0.14	15.62	37,37,41,45	4
3	BCT	B	1003	4/4	0.18	13.59	18,33,35,38	4
5	EDO	B	1009	4/4	0.24	12.20	9,13,15,15	4
5	EDO	B	1007	4/4	0.12	11.71	29,30,30,33	0
5	EDO	A	1010	4/4	0.14	10.39	27,28,30,33	0
5	EDO	A	1013	4/4	0.21	10.38	39,43,44,44	0
2	GLY	B	1001	5/5	0.27	10.27	55,59,64,65	0
5	EDO	A	1011	4/4	0.25	10.04	16,17,21,26	4
5	EDO	A	1012	4/4	0.26	8.41	12,12,17,27	4
5	EDO	A	1016	4/4	0.16	7.71	22,25,28,30	4
2	GLY	A	1001	5/5	0.24	7.07	33,35,49,53	0
4	PO4	A	1008	5/5	0.14	6.63	12,36,38,42	5
3	BCT	A	1005	4/4	0.12	5.95	15,27,30,31	4
2	GLY	A	1004	5/5	0.18	5.26	39,39,41,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	EDO	A	1014	4/4	0.11	5.11	33,42,46,46	0
5	EDO	A	1017	4/4	0.13	4.71	20,29,32,32	4
5	EDO	B	1008	4/4	0.27	4.58	42,43,47,49	0
5	EDO	A	1015	4/4	0.12	2.96	19,23,34,36	4
2	GLY	A	1002	5/5	0.22	2.54	35,36,39,43	5
4	PO4	A	1009	5/5	0.12	2.31	20,24,26,32	5
4	PO4	B	1004	5/5	0.11	1.99	28,50,52,56	0
2	GLY	B	1002	5/5	0.24	1.42	34,43,45,49	0
4	PO4	B	1005	5/5	0.11	1.21	21,27,30,31	5
3	BCT	A	1007	4/4	0.13	0.74	20,26,26,34	4
5	EDO	B	1006	4/4	0.10	0.19	21,23,30,40	0

6.5 Other polymers ⓘ

There are no such residues in this entry.