



# Full wwPDB X-ray Structure Validation Report

Jun 17, 2014 – 08:13 AM EDT

PDB ID : 4LPU  
Title : Crystal structure of TENCON variant P40AR2-32R2  
Authors : Teplyakov, A.; Obmolova, G.; Gilliland, G.L.  
Deposited on : 2013-07-16  
Resolution : 3.40 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : **FAILED**  
Xtriage (Phenix) : dev-1439  
EDS : stable23161  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23161

# 1 Overall quality at a glance



The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1017 (3.52-3.28)
Clashscore	79885	1214 (3.50-3.30)
Ramachandran outliers	78287	1177 (3.50-3.30)
Sidechain outliers	78261	1177 (3.50-3.30)
RSRZ outliers	66119	1017 (3.52-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	98	
1	B	98	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1390 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TENCON variant P40AR2-32R2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	93	Total	C	N	O	S	0	0	0
			706	455	109	141	1			
1	B	93	Total	C	N	O		0	0	0
			684	439	106	139				



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.73Å 84.73Å 127.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 3.40 27.77 – 3.40	Depositor EDS
% Data completeness (in resolution range)	93.8 (15.00-3.40) 94.1 (27.77-3.40)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.55 (at 3.38Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.277 , 0.335 0.273 , 0.330	Depositor DCC
$R_{free}$ test set	170 reflections (4.65%)	DCC
Wilson B-factor (Å <sup>2</sup> )	89.6	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 28.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 3822 reflections	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	1390	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/725	0.66	0/995
1	B	0.47	0/701	0.62	0/965
All	All	0.48	0/1426	0.64	0/1960

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	706	0	665	23	0
1	B	684	0	628	27	0
All	All	1390	0	1293	42	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

All (42) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:77:TYR:HB3	1:B:78:VAL:HB	1.74	0.69
1:A:25:PRO:HB2	1:A:28:TRP:HD1	1.58	0.68

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:19:ARG:HG2	1:B:19:ARG:O	1.98	0.63
1:B:59:LEU:HG	1:B:62:LEU:HD21	1.81	0.62
1:A:32:PHE:HB2	1:A:50:VAL:HG12	1.84	0.59
1:A:11:SER:O	1:A:13:VAL:HG23	2.05	0.56
1:A:35:GLN:HE21	1:A:71:SER:HB2	1.72	0.55
1:B:30:ASP:OD2	1:B:76:TYR:HA	2.08	0.53
1:B:82:ASN:HB3	1:B:83:PRO:HD2	1.91	0.53
1:A:35:GLN:HG3	1:A:45:ALA:HB1	1.90	0.53
1:A:64:PRO:O	1:A:66:THR:OG1	2.26	0.52
1:A:77:TYR:HB2	1:B:79:ALA:HB3	1.92	0.52
1:B:15:GLU:HB3	1:B:64:PRO:HG3	1.92	0.52
1:A:30:ASP:HA	1:A:52:GLY:HA3	1.93	0.50
1:B:32:PHE:HB2	1:B:50:VAL:HG13	1.93	0.49
1:B:21:SER:OG	1:B:56:SER:HB3	2.12	0.49
1:A:10:VAL:HA	1:A:19:ARG:O	2.13	0.48
1:A:7:ASN:O	1:A:22:TRP:HA	2.13	0.48
1:A:14:THR:OG1	1:A:17:SER:N	2.44	0.47
1:A:50:VAL:HG22	1:A:51:PRO:HD2	1.96	0.47
1:A:50:VAL:HG22	1:A:51:PRO:CD	2.44	0.47
1:B:37:GLN:O	1:B:69:THR:N	2.40	0.46
1:B:2:LEU:HA	1:B:3:PRO:HD2	1.79	0.46
1:A:88:PHE:CE1	1:B:10:VAL:HG22	2.51	0.45
1:B:36:TYR:CZ	1:B:59:LEU:HD11	2.52	0.45
1:A:36:TYR:HA	1:A:69:THR:O	2.16	0.45
1:B:39:SER:HA	1:B:69:THR:OG1	2.16	0.45
1:A:77:TYR:HB2	1:B:79:ALA:CB	2.47	0.44
1:B:36:TYR:CE1	1:B:46:ILE:HB	2.52	0.44
1:B:16:ASP:N	1:B:16:ASP:OD1	2.51	0.44
1:A:77:TYR:N	1:B:79:ALA:HB3	2.33	0.44
1:B:19:ARG:O	1:B:19:ARG:CG	2.66	0.43
1:B:19:ARG:O	1:B:20:LEU:C	2.57	0.43
1:B:42:VAL:O	1:B:43:GLY:C	2.57	0.43
1:B:22:TRP:NE1	1:B:55:ARG:HA	2.34	0.42
1:B:38:GLU:C	1:B:40:GLU:N	2.73	0.42
1:A:86:ALA:HB2	1:B:8:LEU:HD23	2.00	0.42
1:A:21:SER:OG	1:A:56:SER:HB3	2.20	0.42
1:A:77:TYR:H	1:B:79:ALA:HB3	1.86	0.41
1:A:25:PRO:HA	1:A:26:PRO:HD2	1.70	0.40
1:B:52:GLY:C	1:B:54:GLU:H	2.24	0.40
1:A:71:SER:HA	1:B:84:LEU:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	91/98 (93%)	82 (90%)	6 (7%)	3 (3%)	6	50
1	B	91/98 (93%)	79 (87%)	10 (11%)	2 (2%)	10	60
All	All	182/196 (93%)	161 (88%)	16 (9%)	5 (3%)	8	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	GLY
1	B	40	GLU
1	B	43	GLY
1	A	65	GLY
1	A	61	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	75/87 (86%)	59 (79%)	16 (21%)	1	8
1	B	70/87 (80%)	66 (94%)	4 (6%)	29	75
All	All	145/174 (83%)	125 (86%)	20 (14%)	5	28

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	9	VAL
1	A	11	SER
1	A	28	TRP

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	30	ASP
1	A	35	GLN
1	A	47	ASN
1	A	50	VAL
1	A	56	SER
1	A	58	ASP
1	A	63	LYS
1	A	66	THR
1	A	72	ILE
1	A	77	TYR
1	A	81	SER
1	A	90	THR
1	B	2	LEU
1	B	49	THR
1	B	69	THR
1	B	75	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	37	GLN
1	A	47	ASN
1	B	35	GLN
1	B	47	ASN
1	B	82	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

Mogul failed to run properly - this section will therefore be empty.

## 5.5 Carbohydrates ⓘ

Mogul failed to run properly - this section will therefore be empty.

## 5.6 Ligand geometry ⓘ

Mogul failed to run properly - this section will therefore be empty.

## 5.7 Other polymers ⓘ

Mogul failed to run properly - this section will therefore be empty.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	93/98 (94%)	-0.17	0 100 100	31, 54, 85, 127	0
1	B	93/98 (94%)	-0.07	0 100 100	37, 68, 98, 121	0
All	All	186/196 (94%)	-0.12	0 100 100	31, 62, 93, 127	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

### 6.5 Other polymers ⓘ

There are no such residues in this entry.