



Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 06:13 PM GMT

PDB ID : 4LXA
Title : Crystal Structure of Human Beta Secretase in Complex with Compound 11a
Authors : Rondeau, J.M.; Bourgier, E.
Deposited on : 2013-07-29
Resolution : 1.95 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1321 (1.96-1.96)
Clashscore	79885	1488 (1.96-1.96)
Ramachandran outliers	78287	1475 (1.96-1.96)
Sidechain outliers	78261	1475 (1.96-1.96)
RSRZ outliers	66119	1321 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	402	
1	B	402	
1	C	402	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9566 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	376	Total	C	N	O	S	0	0	0
			2961	1895	492	560	14			
1	B	378	Total	C	N	O	S	0	0	0
			2971	1901	494	562	14			
1	C	381	Total	C	N	O	S	0	0	0
			2993	1917	497	565	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33P	GLY	-	EXPRESSION TAG	UNP P56817
A	34P	PRO	-	EXPRESSION TAG	UNP P56817
B	33P	GLY	-	EXPRESSION TAG	UNP P56817
B	34P	PRO	-	EXPRESSION TAG	UNP P56817
C	33P	GLY	-	EXPRESSION TAG	UNP P56817
C	34P	PRO	-	EXPRESSION TAG	UNP P56817

- Molecule 2 is (1R,3S,4S,5R)-3-{4-AMINO-3-FLUORO-5-[(1,1,1,3,3,3-HEXAFLUOROPROPAN-2-YL)OXY]BENZYL}-5-[(3-TERT-BUTYLBENZYL)AMINO]TETRAHYDRO-2H-THIOPYRAN-4-OL1-OXIDE (three-letter code: 1YS) (formula: C₂₆H₃₁F₇N₂O₃S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			39	26	7	2	3	1		
2	B	1	Total	C	F	N	O	S	0	0
			39	26	7	2	3	1		
2	C	1	Total	C	F	N	O	S	0	0
			39	26	7	2	3	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	172	Total	O	0	0
			172	172		
5	B	168	Total	O	0	0
			168	168		
5	C	173	Total	O	0	0
			173	173		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Beta-secretase 1



• Molecule 1: Beta-secretase 1



• Molecule 1: Beta-secretase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.93Å 103.07Å 100.23Å 90.00° 104.33° 90.00°	Depositor
Resolution (Å)	37.65 – 1.95 37.65 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.4 (37.65-1.95) 99.6 (37.65-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.95Å)	Xtriage
Refinement program	CNX	Depositor
R, R_{free}	0.199 , 0.222 0.195 , 0.217	Depositor DCC
R_{free} test set	5880 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 116548 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9566	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, 1YS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/3036	0.46	0/4126
1	B	0.27	0/3046	0.46	0/4140
1	C	0.26	0/3070	0.46	0/4173
All	All	0.27	0/9152	0.46	0/12439

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2961	0	2870	17	0
1	B	2971	0	2880	19	0
1	C	2993	0	2899	16	0
2	A	39	0	31	0	0
2	B	39	0	31	0	0
2	C	39	0	31	0	0
3	B	6	0	8	0	0
4	C	5	0	0	0	0
5	A	172	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	168	0	0	0	0
5	C	173	0	0	0	0
All	All	9566	0	8750	52	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (52) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:267:LEU:HD13	1:B:319:CYS:HB3	1.71	0.71
1:A:267:LEU:HD13	1:A:319:CYS:HB3	1.75	0.67
1:A:254:THR:HG23	1:A:255:GLU:HG3	1.81	0.62
1:B:267:LEU:HD22	1:B:309:VAL:HG21	1.82	0.62
1:A:2:MET:HG2	1:A:90:GLY:HA2	1.83	0.61
1:B:357:SER:O	1:B:360:HIS:HB3	2.01	0.60
1:B:9:LYS:HZ1	1:B:168:ALA:N	1.99	0.60
1:C:335:ALA:O	1:C:339:GLU:HG3	2.02	0.60
1:B:264:GLY:O	1:B:321:LYS:HE3	2.03	0.58
1:B:232:THR:O	1:B:336:VAL:HG13	2.03	0.58
1:B:64:ARG:HH11	1:B:64:ARG:HB3	1.69	0.58
1:A:278:ASN:H	1:A:278:ASN:HD22	1.52	0.57
1:A:125:GLU:HG3	5:A:744:HOH:O	2.05	0.56
1:B:276:PRO:O	1:B:279:ILE:HG12	2.06	0.56
1:C:249:LYS:HE2	1:C:262:TRP:CD1	2.41	0.55
1:C:267:LEU:HD13	1:C:319:CYS:HB3	1.88	0.55
1:C:363:ASP:HB3	1:C:366:ARG:O	2.05	0.55
1:A:335:ALA:O	1:A:339:GLU:HG3	2.08	0.54
1:C:267:LEU:HD22	1:C:309:VAL:HG21	1.89	0.54
1:B:63:LEU:HG	1:B:81:GLY:HA2	1.92	0.52
1:A:365:PHE:HB2	5:A:768:HOH:O	2.08	0.52
1:C:63:LEU:HG	1:C:81:GLY:HA2	1.93	0.50
1:A:267:LEU:HD22	1:A:309:VAL:HG21	1.94	0.49
1:C:110:ILE:HB	1:C:113:SER:HB3	1.94	0.49
1:B:205:ARG:NH2	1:B:212:ASP:HB2	2.28	0.49
1:B:267:LEU:HD21	1:B:312:VAL:HG13	1.94	0.49
1:C:304:GLN:HG3	1:C:361:VAL:HG21	1.94	0.49
1:B:267:LEU:HD22	1:B:309:VAL:CG2	2.42	0.48
1:B:2:MET:HG2	1:B:90:GLY:HA2	1.96	0.48
1:A:363:ASP:HB3	1:A:366:ARG:O	2.14	0.48
1:B:110:ILE:HB	1:B:113:SER:HB3	1.96	0.47
1:C:12:GLN:OE1	1:C:113:SER:HA	2.15	0.47
1:C:267:LEU:HD22	1:C:309:VAL:CG2	2.44	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:205:ARG:HB3	1:A:286:TYR:HB2	1.98	0.46
1:B:222:TYR:HA	1:B:223:ASP:HA	1.62	0.45
1:C:276:PRO:O	1:C:279:ILE:HG12	2.17	0.45
1:A:282:VAL:HG12	1:A:301:LEU:HD23	1.99	0.44
1:A:244:ALA:O	1:A:248:ILE:HG13	2.17	0.44
1:C:293:ASN:HA	1:C:375:VAL:HA	1.99	0.43
1:A:110:ILE:HB	1:A:113:SER:HB3	2.01	0.43
1:C:222:TYR:HA	1:C:223:ASP:HA	1.65	0.43
1:A:211:GLN:HB2	1:A:211:GLN:HE21	1.62	0.43
1:C:2:MET:HG2	1:C:90:GLY:HA2	2.00	0.43
1:A:222:TYR:HA	1:A:223:ASP:HA	1.70	0.43
1:A:9:LYS:HE2	1:A:9:LYS:HB3	1.83	0.42
1:B:249:LYS:HE2	1:B:262:TRP:CD1	2.54	0.42
1:B:364:GLU:HG3	1:B:365:PHE:CD2	2.55	0.42
1:C:304:GLN:CG	1:C:361:VAL:HG21	2.50	0.41
1:B:314:THR:O	1:B:314:THR:HG23	2.20	0.41
1:C:270:TRP:O	1:C:317:ASP:HB3	2.21	0.41
1:A:23:PRO:HA	1:A:24:PRO:HD3	1.93	0.40
1:B:125:GLU:O	1:B:125:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/402 (92%)	366 (98%)	6 (2%)	0	100	100
1	B	374/402 (93%)	366 (98%)	8 (2%)	0	100	100
1	C	377/402 (94%)	368 (98%)	9 (2%)	0	100	100
All	All	1123/1206 (93%)	1100 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/342 (94%)	318 (99%)	4 (1%)	82	79
1	B	322/342 (94%)	319 (99%)	3 (1%)	87	86
1	C	324/342 (95%)	322 (99%)	2 (1%)	92	92
All	All	968/1026 (94%)	959 (99%)	9 (1%)	87	86

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	GLN
1	A	197	TRP
1	A	211	GLN
1	A	278	ASN
1	B	125	GLU
1	B	197	TRP
1	B	209	ASN
1	C	197	TRP
1	C	362	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	ASN
1	A	278	ASN
1	A	326	GLN
1	B	114	ASN
1	B	278	ASN
1	B	293	ASN
1	B	326	GLN
1	C	114	ASN
1	C	326	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1YS	A	501	-	41,41,41	1.57	6 (14%)	63,63,63	1.43	10 (15%)
3	GOL	B	501	-	5,5,5	0.62	0	5,5,5	0.56	0
2	1YS	B	502	-	41,41,41	1.63	7 (17%)	63,63,63	1.42	10 (15%)
2	1YS	C	501	-	41,41,41	1.58	6 (14%)	63,63,63	1.40	9 (14%)
4	PO4	C	502	-	4,4,4	0.85	0	6,6,6	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1YS	A	501	-	-	0/31/47/47	0/3/3/3
3	GOL	B	501	-	-	0/4/4/4	0/0/0/0
2	1YS	B	502	-	-	0/31/47/47	0/3/3/3
2	1YS	C	501	-	-	0/31/47/47	0/3/3/3
4	PO4	C	502	-	-	0/0/0/0	0/0/0/0

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	1YS	C25-C23	4.21	1.56	1.52
2	B	502	1YS	C25-C23	4.05	1.56	1.52
2	C	501	1YS	C25-C23	3.92	1.56	1.52
2	B	502	1YS	C19-C21	2.99	1.56	1.53
2	C	501	1YS	C19-C21	2.93	1.56	1.53
2	A	501	1YS	C16-C19	2.69	1.57	1.53
2	A	501	1YS	C19-C21	2.69	1.56	1.53
2	B	502	1YS	C29-S28	-2.64	1.78	1.81
2	C	501	1YS	C42-C41	2.50	1.43	1.39
2	B	502	1YS	C11-C10	2.50	1.42	1.37
2	A	501	1YS	C42-C41	2.49	1.43	1.39
2	C	501	1YS	C16-C19	2.49	1.57	1.53
2	B	502	1YS	C42-C41	2.49	1.43	1.39
2	C	501	1YS	C11-C10	2.46	1.42	1.37
2	B	502	1YS	C16-C19	2.46	1.57	1.53
2	A	501	1YS	C39-C41	2.20	1.43	1.39
2	B	502	1YS	C39-C41	2.19	1.43	1.39
2	C	501	1YS	C39-C41	2.18	1.43	1.39
2	A	501	1YS	C11-C10	2.15	1.41	1.37

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	1YS	O32-S28-C25	4.57	109.53	105.64
2	B	502	1YS	C35-N33-C23	4.53	122.27	114.94
2	B	502	1YS	C8-C9-C10	4.45	119.46	115.94
2	A	501	1YS	C35-N33-C23	4.43	122.10	114.94
2	C	501	1YS	C8-C9-C10	4.40	119.42	115.94
2	A	501	1YS	C8-C9-C10	4.17	119.24	115.94
2	C	501	1YS	C35-N33-C23	3.95	121.32	114.94
2	A	501	1YS	F63-C10-C9	3.66	120.81	117.70
2	B	502	1YS	O32-S28-C25	3.65	108.76	105.64
2	A	501	1YS	O32-S28-C25	3.05	108.24	105.64
2	B	502	1YS	F63-C10-C9	2.94	120.20	117.70
2	C	501	1YS	C11-C10-C9	-2.91	120.34	123.58
2	A	501	1YS	C29-C19-C16	2.85	114.69	109.58
2	A	501	1YS	C11-C10-C9	-2.83	120.42	123.58
2	A	501	1YS	C13-C16-C19	-2.83	108.58	113.71
2	B	502	1YS	C11-C10-C9	-2.78	120.48	123.58
2	C	501	1YS	C29-C19-C16	2.77	114.55	109.58
2	B	502	1YS	C29-C19-C16	2.71	114.45	109.58
2	B	502	1YS	C25-C23-C21	2.65	115.11	109.95
2	B	502	1YS	C13-C16-C19	-2.44	109.27	113.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	1YS	C53-C48-C41	2.41	116.25	110.39
2	A	501	1YS	C25-C23-C21	2.39	114.59	109.95
2	C	501	1YS	C25-C23-C21	2.38	114.59	109.95
2	C	501	1YS	C13-C16-C19	-2.34	109.47	113.71
2	C	501	1YS	F63-C10-C9	2.33	119.69	117.70
2	A	501	1YS	C2-C5-C67	2.18	116.38	112.47
2	B	502	1YS	C2-C5-C67	2.18	116.37	112.47
2	C	501	1YS	C53-C48-C41	2.13	115.58	110.39
2	B	502	1YS	C53-C48-C41	2.04	115.36	110.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	376/402 (93%)	0.27	22 (5%) 22 28	24, 38, 71, 106	0
1	B	378/402 (94%)	0.55	41 (10%) 6 7	23, 39, 76, 110	0
1	C	381/402 (94%)	0.40	29 (7%) 14 17	26, 40, 73, 105	0
All	All	1135/1206 (94%)	0.41	92 (8%) 12 15	23, 39, 74, 110	0

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	314	THR	12.5
1	B	312	VAL	12.1
1	A	312	VAL	10.6
1	A	314	THR	9.2
1	B	313	ALA	9.1
1	C	312	VAL	9.0
1	C	315	SER	8.3
1	C	313	ALA	7.8
1	A	313	ALA	7.7
1	C	314	THR	7.2
1	A	311	ASP	6.9
1	B	311	ASP	6.4
1	B	310	GLU	5.8
1	B	315	SER	5.6
1	A	316	GLN	5.5
1	B	256	LYS	5.4
1	C	310	GLU	5.4
1	A	315	SER	5.3
1	C	168	ALA	5.3
1	B	365	PHE	5.0
1	B	316	GLN	4.7
1	A	310	GLU	4.6
1	C	259	ASP	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	160	PRO	4.4
1	C	316	GLN	4.4
1	C	311	ASP	4.1
1	C	272	ALA	4.0
1	B	364	GLU	3.9
1	B	254	THR	3.6
1	B	266	GLN	3.6
1	B	309	VAL	3.5
1	B	361	VAL	3.3
1	A	256	LYS	3.3
1	B	253	SER	3.3
1	C	309	VAL	3.3
1	B	317	ASP	3.3
1	B	145	HIS	3.1
1	A	317	ASP	3.1
1	B	51	TYR	3.1
1	C	267	LEU	3.1
1	B	252	SER	3.0
1	B	257	PHE	3.0
1	A	145	HIS	3.0
1	B	267	LEU	3.0
1	A	364	GLU	3.0
1	B	274	THR	3.0
1	A	259	ASP	3.0
1	C	49	HIS	3.0
1	B	168	ALA	2.9
1	B	258	PRO	2.9
1	C	64	ARG	2.9
1	B	49	HIS	2.8
1	C	273	GLY	2.8
1	B	250	ALA	2.8
1	C	145	HIS	2.8
1	B	46	PRO	2.8
1	C	159	PHE	2.8
1	A	64	ARG	2.7
1	C	46(P)	SER	2.7
1	B	265	GLU	2.7
1	B	268	VAL	2.7
1	A	267	LEU	2.7
1	A	365	PHE	2.7
1	C	158	GLY	2.6
1	A	384	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	270	TRP	2.5
1	C	317	ASP	2.5
1	B	362	HIS	2.5
1	B	46(P)	SER	2.5
1	A	319	CYS	2.4
1	B	260	GLY	2.4
1	C	269	CYS	2.4
1	C	271	GLN	2.4
1	B	280	PHE	2.4
1	C	256	LYS	2.3
1	A	272	ALA	2.3
1	B	271	GLN	2.3
1	A	266	GLN	2.2
1	B	55	GLN	2.2
1	B	262	TRP	2.2
1	B	56	LEU	2.2
1	B	157	ALA	2.2
1	C	365	PHE	2.2
1	B	255	GLU	2.1
1	A	362	HIS	2.1
1	A	385	ASN	2.1
1	C	362	HIS	2.1
1	A	271	GLN	2.1
1	C	56	LEU	2.1
1	C	55	GLN	2.1
1	C	266	GLN	2.1
1	B	211	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	1YS	B	502	39/39	0.11	-	26,32,44,46	0
3	GOL	B	501	6/6	0.11	-	40,43,44,46	0
4	PO4	C	502	5/5	0.13	-	45,46,47,49	0
2	1YS	A	501	39/39	0.12	-	27,33,41,44	0
2	1YS	C	501	39/39	0.12	-	28,34,44,46	0

6.5 Other polymers ⓘ

There are no such residues in this entry.