



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 04:29 AM GMT

PDB ID : 3LZK
Title : The crystal structure of a probably aromatic amino acid degradation protein from *Sinorhizobium meliloti* 1021
Authors : Tan, K.; Xu, X.; Cui, H.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2010-03-01
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	359	
1	B	359	
1	C	359	
1	D	359	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	CA	B	339	-	X
2	CA	D	339	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10754 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fumarylacetoacetate hydrolase family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	S	Se	0	1	0
			2581	1626	456	487	2	10			
1	B	339	Total	C	N	O	S	Se	0	0	0
			2562	1612	452	486	2	10			
1	C	343	Total	C	N	O	S	Se	0	1	0
			2609	1645	460	492	2	10			
1	D	340	Total	C	N	O	S	Se	0	1	0
			2573	1618	454	489	2	10			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	GLY	-	EXPRESSION TAG	UNP Q92LT4
A	-18	SER	-	EXPRESSION TAG	UNP Q92LT4
A	-17	SER	-	EXPRESSION TAG	UNP Q92LT4
A	-16	HIS	-	EXPRESSION TAG	UNP Q92LT4
A	-15	HIS	-	EXPRESSION TAG	UNP Q92LT4
A	-14	HIS	-	EXPRESSION TAG	UNP Q92LT4
A	-13	HIS	-	EXPRESSION TAG	UNP Q92LT4
A	-12	HIS	-	EXPRESSION TAG	UNP Q92LT4
A	-11	HIS	-	EXPRESSION TAG	UNP Q92LT4
A	-10	SER	-	EXPRESSION TAG	UNP Q92LT4
A	-9	SER	-	EXPRESSION TAG	UNP Q92LT4
A	-8	GLY	-	EXPRESSION TAG	UNP Q92LT4
A	-7	ARG	-	EXPRESSION TAG	UNP Q92LT4
A	-6	GLU	-	EXPRESSION TAG	UNP Q92LT4
A	-5	ASN	-	EXPRESSION TAG	UNP Q92LT4
A	-4	LEU	-	EXPRESSION TAG	UNP Q92LT4
A	-3	TYR	-	EXPRESSION TAG	UNP Q92LT4
A	-2	PHE	-	EXPRESSION TAG	UNP Q92LT4
A	-1	GLN	-	EXPRESSION TAG	UNP Q92LT4
A	0	GLY	-	EXPRESSION TAG	UNP Q92LT4
A	1	MSE	-	EXPRESSION TAG	UNP Q92LT4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	GLY	-	EXPRESSION TAG	UNP Q92LT4
B	-18	SER	-	EXPRESSION TAG	UNP Q92LT4
B	-17	SER	-	EXPRESSION TAG	UNP Q92LT4
B	-16	HIS	-	EXPRESSION TAG	UNP Q92LT4
B	-15	HIS	-	EXPRESSION TAG	UNP Q92LT4
B	-14	HIS	-	EXPRESSION TAG	UNP Q92LT4
B	-13	HIS	-	EXPRESSION TAG	UNP Q92LT4
B	-12	HIS	-	EXPRESSION TAG	UNP Q92LT4
B	-11	HIS	-	EXPRESSION TAG	UNP Q92LT4
B	-10	SER	-	EXPRESSION TAG	UNP Q92LT4
B	-9	SER	-	EXPRESSION TAG	UNP Q92LT4
B	-8	GLY	-	EXPRESSION TAG	UNP Q92LT4
B	-7	ARG	-	EXPRESSION TAG	UNP Q92LT4
B	-6	GLU	-	EXPRESSION TAG	UNP Q92LT4
B	-5	ASN	-	EXPRESSION TAG	UNP Q92LT4
B	-4	LEU	-	EXPRESSION TAG	UNP Q92LT4
B	-3	TYR	-	EXPRESSION TAG	UNP Q92LT4
B	-2	PHE	-	EXPRESSION TAG	UNP Q92LT4
B	-1	GLN	-	EXPRESSION TAG	UNP Q92LT4
B	0	GLY	-	EXPRESSION TAG	UNP Q92LT4
B	1	MSE	-	EXPRESSION TAG	UNP Q92LT4
C	-19	GLY	-	EXPRESSION TAG	UNP Q92LT4
C	-18	SER	-	EXPRESSION TAG	UNP Q92LT4
C	-17	SER	-	EXPRESSION TAG	UNP Q92LT4
C	-16	HIS	-	EXPRESSION TAG	UNP Q92LT4
C	-15	HIS	-	EXPRESSION TAG	UNP Q92LT4
C	-14	HIS	-	EXPRESSION TAG	UNP Q92LT4
C	-13	HIS	-	EXPRESSION TAG	UNP Q92LT4
C	-12	HIS	-	EXPRESSION TAG	UNP Q92LT4
C	-11	HIS	-	EXPRESSION TAG	UNP Q92LT4
C	-10	SER	-	EXPRESSION TAG	UNP Q92LT4
C	-9	SER	-	EXPRESSION TAG	UNP Q92LT4
C	-8	GLY	-	EXPRESSION TAG	UNP Q92LT4
C	-7	ARG	-	EXPRESSION TAG	UNP Q92LT4
C	-6	GLU	-	EXPRESSION TAG	UNP Q92LT4
C	-5	ASN	-	EXPRESSION TAG	UNP Q92LT4
C	-4	LEU	-	EXPRESSION TAG	UNP Q92LT4
C	-3	TYR	-	EXPRESSION TAG	UNP Q92LT4
C	-2	PHE	-	EXPRESSION TAG	UNP Q92LT4
C	-1	GLN	-	EXPRESSION TAG	UNP Q92LT4
C	0	GLY	-	EXPRESSION TAG	UNP Q92LT4
C	1	MSE	-	EXPRESSION TAG	UNP Q92LT4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	GLY	-	EXPRESSION TAG	UNP Q92LT4
D	-18	SER	-	EXPRESSION TAG	UNP Q92LT4
D	-17	SER	-	EXPRESSION TAG	UNP Q92LT4
D	-16	HIS	-	EXPRESSION TAG	UNP Q92LT4
D	-15	HIS	-	EXPRESSION TAG	UNP Q92LT4
D	-14	HIS	-	EXPRESSION TAG	UNP Q92LT4
D	-13	HIS	-	EXPRESSION TAG	UNP Q92LT4
D	-12	HIS	-	EXPRESSION TAG	UNP Q92LT4
D	-11	HIS	-	EXPRESSION TAG	UNP Q92LT4
D	-10	SER	-	EXPRESSION TAG	UNP Q92LT4
D	-9	SER	-	EXPRESSION TAG	UNP Q92LT4
D	-8	GLY	-	EXPRESSION TAG	UNP Q92LT4
D	-7	ARG	-	EXPRESSION TAG	UNP Q92LT4
D	-6	GLU	-	EXPRESSION TAG	UNP Q92LT4
D	-5	ASN	-	EXPRESSION TAG	UNP Q92LT4
D	-4	LEU	-	EXPRESSION TAG	UNP Q92LT4
D	-3	TYR	-	EXPRESSION TAG	UNP Q92LT4
D	-2	PHE	-	EXPRESSION TAG	UNP Q92LT4
D	-1	GLN	-	EXPRESSION TAG	UNP Q92LT4
D	0	GLY	-	EXPRESSION TAG	UNP Q92LT4
D	1	MSE	-	EXPRESSION TAG	UNP Q92LT4

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	108	Total O 108 108	0	0
3	B	74	Total O 74 74	0	0
3	C	145	Total O 145 145	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	98	Total	O	0	0
			98	98		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fumarylacetoacetate hydrolase family protein

Chain A: 



- Molecule 1: Fumarylacetoacetate hydrolase family protein

Chain B: 




- Molecule 1: Fumarylacetoacetate hydrolase family protein

Chain C: 



- Molecule 1: Fumarylacetoacetate hydrolase family protein

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.77Å 188.89Å 67.74Å 90.00° 115.17° 90.00°	Depositor
Resolution (Å)	33.60 – 1.90 47.47 – 1.90	Depositor EDS
% Data completeness (in resolution range)	90.2 (33.60-1.90) 96.2 (47.47-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 1.90Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, R_{free}	0.185 , 0.227 0.184 , 0.224	Depositor DCC
R_{free} test set	5365 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.475	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 42.3	EDS
Estimated twinning fraction	0.027 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 107578 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10754	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2631	0.55	0/3546
1	B	0.33	0/2608	0.52	0/3516
1	C	0.40	0/2660	0.56	0/3586
1	D	0.36	0/2622	0.51	0/3533
All	All	0.37	0/10521	0.53	0/14181

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2581	0	0	5	0
1	B	2562	0	0	5	0
1	C	2609	0	0	2	0
1	D	2573	0	0	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	108	0	0	1	0
3	B	74	0	0	0	0
3	C	145	0	0	0	0
3	D	98	0	0	1	0
All	All	10754	0	0	19	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (19) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:95[A]:ARG:CG	1:A:95[A]:ARG:NH1	2.68	0.56
1:B:137:ASP:OD2	1:B:174:ARG:NH1	2.39	0.55
1:D:129:MSE:CE	1:D:134:TRP:CZ3	2.89	0.55
1:D:211:ASP:OD1	1:D:211:ASP:C	2.48	0.52
1:D:24:ARG:NE	1:D:63:THR:OG1	2.43	0.50
1:D:220:HIS:CD2	3:D:350:HOH:O	2.65	0.50
1:A:294:GLU:O	1:A:298:GLY:N	2.45	0.50
1:A:211:ASP:OD2	1:A:216:HIS:NE2	2.46	0.48
1:B:320:THR:OG1	1:B:322:HIS:CD2	2.67	0.47
1:A:139:GLU:OE1	1:A:174:ARG:NH1	2.47	0.47
1:D:320:THR:OG1	1:D:322:HIS:CD2	2.69	0.46
1:B:211:ASP:OD2	1:B:216:HIS:NE2	2.49	0.46
1:A:20:LYS:NZ	3:A:393:HOH:O	2.51	0.44
1:B:98:ARG:O	1:B:99:ASN:CB	2.66	0.43
1:D:275:ARG:NH1	1:D:297:GLU:OE2	2.51	0.43
1:D:129:MSE:CE	1:D:134:TRP:CE3	3.03	0.42
1:C:21:ASP:OD2	1:C:23:THR:OG1	2.39	0.41
1:C:68:GLU:OE1	1:C:124:ARG:NH2	2.54	0.41
1:B:276:PRO:O	1:B:279:GLU:N	2.53	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/359 (94%)	326 (96%)	13 (4%)	0	100	100
1	B	337/359 (94%)	318 (94%)	18 (5%)	1 (0%)	50	37
1	C	342/359 (95%)	330 (96%)	12 (4%)	0	100	100
1	D	339/359 (94%)	325 (96%)	13 (4%)	1 (0%)	50	37
All	All	1357/1436 (94%)	1299 (96%)	56 (4%)	2 (0%)	59	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	99	ASN
1	D	298	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/271 (98%)	257 (97%)	9 (3%)	49	36
1	B	264/271 (97%)	254 (96%)	10 (4%)	44	31
1	C	269/271 (99%)	263 (98%)	6 (2%)	64	57
1	D	265/271 (98%)	252 (95%)	13 (5%)	35	21
All	All	1064/1084 (98%)	1026 (96%)	38 (4%)	48	33

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ARG
1	A	65	ARG
1	A	95[A]	ARG
1	A	95[B]	ARG
1	A	138	MSE
1	A	238	THR
1	A	269	LEU
1	A	297	GLU
1	A	337	ARG
1	B	57	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	77	ARG
1	B	98	ARG
1	B	102	MSE
1	B	138	MSE
1	B	226	GLU
1	B	238	THR
1	B	265	VAL
1	B	269	LEU
1	B	306	LEU
1	C	57	GLU
1	C	138	MSE
1	C	214	LYS
1	C	297	GLU
1	C	306	LEU
1	C	319	ARG
1	D	-1	GLN
1	D	33	ARG
1	D	57	GLU
1	D	99	ASN
1	D	101	GLU
1	D	138	MSE
1	D	174	ARG
1	D	215	LEU
1	D	238	THR
1	D	265	VAL
1	D	275	ARG
1	D	291	ARG
1	D	306	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/359 (94%)	-0.12	4 (1%) 75 78	33, 49, 79, 128	0
1	B	339/359 (94%)	0.30	16 (4%) 30 30	38, 59, 101, 127	0
1	C	343/359 (95%)	-0.07	2 (0%) 86 88	26, 41, 64, 105	0
1	D	340/359 (94%)	0.13	16 (4%) 30 30	28, 54, 105, 158	0
All	All	1362/1436 (94%)	0.06	38 (2%) 50 51	26, 50, 89, 158	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	296	ILE	10.0
1	D	293	ILE	8.8
1	D	338	GLY	8.2
1	B	273	PRO	5.6
1	B	296	ILE	5.4
1	B	97	ALA	5.3
1	D	96	LYS	5.1
1	B	100	ALA	4.9
1	B	269	LEU	4.7
1	B	93	LEU	4.7
1	D	295	THR	4.6
1	C	104	ALA	4.3
1	B	293	ILE	4.0
1	A	212	GLY	4.0
1	B	96	LYS	3.9
1	B	94	VAL	3.6
1	D	93	LEU	3.5
1	B	134	TRP	3.2
1	D	95	ARG	3.1
1	A	296	ILE	3.0
1	C	-5	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	130	ALA	2.9
1	D	97	ALA	2.9
1	D	298	GLY	2.9
1	B	299	GLY	2.8
1	A	96	LYS	2.8
1	B	268	LYS	2.8
1	D	304[A]	GLN	2.7
1	B	57	GLU	2.6
1	B	275	ARG	2.5
1	D	134	TRP	2.5
1	B	95	ARG	2.3
1	D	269	LEU	2.3
1	A	100	ALA	2.3
1	D	335	TYR	2.2
1	D	308	PHE	2.2
1	D	336	GLU	2.1
1	B	101	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	D	339	1/1	0.20	4.78	49,49,49,49	0
2	CA	B	339	1/1	0.12	2.82	54,54,54,54	0
2	CA	C	339	1/1	0.10	-0.28	35,35,35,35	0
2	CA	A	339	1/1	0.09	-0.89	42,42,42,42	0

6.5 Other polymers ⓘ

There are no such residues in this entry.