



# Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 02:49 PM GMT

PDB ID : 4M1D  
Title : Crystal structure of anti-HIV-1 Fab 447-52D in complex with V3 cyclic peptide MN  
Authors : Killikelly, A.; Kong, X.P.  
Deposited on : 2013-08-02  
Resolution : 1.80 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	L	216	
1	M	216	
2	H	231	
2	I	231	
3	P	14	
3	Q	14	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8039 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab mAb 447-52D Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	216	Total	C	N	O	S	0	0	0
			1603	1007	266	325	5			
1	M	213	Total	C	N	O	S	0	0	0
			1579	994	262	319	4			

- Molecule 2 is a protein called Fab mAb 447-52D Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	231	Total	C	N	O	S	0	0	0
			1735	1092	290	345	8			
2	I	231	Total	C	N	O	S	0	0	0
			1739	1095	291	345	8			

- Molecule 3 is a protein called Cyclic V3 Arch Peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	14	Total	C	N	O	S	0	0	0
			111	70	22	17	2			
3	Q	14	Total	C	N	O	S	0	0	0
			111	70	22	17	2			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			5	3	2		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	H	1	Total	C	O	0	0
			6	3	3		
4	M	1	Total	C	O	0	0
			6	3	3		
4	M	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	M	1	Total	C	O	0	0
			6	3	3		
4	I	1	Total	C	O	0	0
			6	3	3		
4	I	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	291	Total	O	0	0
			291	291		
5	H	311	Total	O	0	0
			311	311		
5	P	18	Total	O	0	0
			18	18		
5	M	213	Total	O	0	0
			213	213		
5	I	213	Total	O	0	0
			213	213		
5	Q	14	Total	O	0	0
			14	14		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab mAb 447-52D Light Chain

Chain L: 



- Molecule 1: Fab mAb 447-52D Light Chain

Chain M: 




- Molecule 2: Fab mAb 447-52D Heavy Chain

Chain H: 



- Molecule 2: Fab mAb 447-52D Heavy Chain

Chain I: 



- Molecule 3: Cyclic V3 Arch Peptide

Chain P: 

There are no outlier residues recorded for this chain.

- Molecule 3: Cyclic V3 Arch Peptide

Chain Q: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	179.84Å 59.78Å 137.07Å 90.00° 127.35° 90.00°	Depositor
Resolution (Å)	45.34 – 1.80 45.34 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.34-1.80) 98.9 (45.34-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.186 , 0.216 0.186 , 0.216	Depositor DCC
$R_{free}$ test set	5299 reflections (4.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.0	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 38.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 106417 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8039	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.42	0/1644	0.60	0/2247
1	M	0.32	0/1620	0.52	0/2215
2	H	0.40	0/1774	0.60	1/2417 (0.0%)
2	I	0.34	0/1778	0.54	0/2421
3	P	0.45	0/114	0.64	0/151
3	Q	0.37	0/114	0.49	0/151
All	All	0.38	0/7044	0.57	1/9602 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	138	LEU	CA-CB-CG	6.50	130.24	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1603	0	1556	13	1
1	M	1579	0	1535	14	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	H	1735	0	1694	20	0
2	I	1739	0	1706	12	1
3	P	111	0	105	0	0
3	Q	111	0	105	0	0
4	H	36	0	48	6	0
4	I	12	0	16	5	0
4	L	35	0	45	6	0
4	M	18	0	24	3	0
5	H	311	0	0	10	4
5	I	213	0	0	3	3
5	L	291	0	0	4	5
5	M	213	0	0	7	3
5	P	18	0	0	0	0
5	Q	14	0	0	0	1
All	All	8039	0	6834	60	9

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (60) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:L:54:ARG:NH1	5:L:628:HOH:O	2.06	0.87
1:M:60:ASP:OD2	5:M:569:HOH:O	2.09	0.69
2:H:192:GLN:NE2	5:H:661:HOH:O	1.90	0.68
1:M:45:LYS:HE2	5:M:602:HOH:O	1.92	0.68
2:H:28:THR:H	4:H:302:GOL:H12	1.60	0.67
2:H:115:SER:H	1:M:126:GLN:NE2	1.93	0.66
2:I:144:ASP:OD2	5:I:553:HOH:O	2.14	0.65
1:M:61:ARG:NH2	4:M:303:GOL:O3	2.18	0.65
1:M:61:ARG:HH21	4:M:303:GOL:HO3	1.43	0.64
2:H:100(E):GLU:OE1	5:H:564:HOH:O	2.15	0.64
4:L:302:GOL:H2	5:H:425:HOH:O	1.99	0.62
1:L:126:GLN:NE2	2:I:115:SER:H	1.99	0.61
2:H:27:PHE:HB2	4:H:302:GOL:H11	1.85	0.59
1:L:118:PHE:CE2	1:L:135:LEU:HD12	2.37	0.59
2:H:45:LEU:HB2	4:H:301:GOL:H31	1.84	0.58
2:H:21:THR:HG23	5:H:493:HOH:O	2.03	0.58
2:H:28:THR:H	4:H:302:GOL:C1	2.17	0.57
2:H:199:ASN:OD1	5:H:514:HOH:O	2.18	0.56
1:L:182:PRO:O	1:L:186:LYS:HG2	2.06	0.55
1:L:118:PHE:HE2	1:L:135:LEU:HD12	1.70	0.55
1:L:79:GLN:HE22	4:L:301:GOL:H2	1.70	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:I:1:GLU:O	5:I:588:HOH:O	2.18	0.55
1:L:132:LEU:HD12	1:L:178:LEU:HD23	1.90	0.54
1:M:2:SER:HA	5:M:547:HOH:O	2.07	0.54
2:I:105:LYS:NZ	4:I:302:GOL:H32	2.23	0.53
1:L:20:THR:H	4:L:303:GOL:C3	2.22	0.53
2:I:12:VAL:HG11	2:I:82(C):LEU:HD13	1.90	0.52
4:L:303:GOL:H11	5:L:493:HOH:O	2.09	0.52
2:I:159:LEU:HD21	2:I:182:VAL:HG21	1.92	0.51
2:I:128:SER:O	2:I:130:SER:N	2.43	0.50
2:H:171:GLN:NE2	5:H:646:HOH:O	2.46	0.49
2:I:107:THR:OG1	4:I:302:GOL:H31	2.13	0.49
2:I:126:PRO:HD2	2:I:213:LEU:HD22	1.94	0.48
2:H:171:GLN:CD	5:H:681:HOH:O	2.51	0.48
4:L:302:GOL:H11	2:H:45:LEU:HD13	1.97	0.47
1:M:3:VAL:N	5:M:547:HOH:O	2.24	0.47
1:M:103:LYS:HE2	5:M:595:HOH:O	2.14	0.46
1:L:79:GLN:NE2	4:L:301:GOL:H2	2.29	0.46
2:I:138:LEU:HB3	2:I:213:LEU:HD11	1.98	0.46
1:M:45:LYS:HE3	1:M:58:ILE:HD11	1.98	0.45
2:I:45:LEU:HB2	4:I:301:GOL:H32	1.98	0.45
1:L:54:ARG:NH1	5:L:637:HOH:O	2.27	0.45
1:L:54:ARG:CZ	5:L:628:HOH:O	2.59	0.44
2:H:64:LYS:HG3	5:H:642:HOH:O	2.18	0.44
1:M:59:PRO:HG3	4:M:303:GOL:H2	2.00	0.44
1:L:135:LEU:HD13	2:H:181:VAL:HG11	1.99	0.44
2:H:71:ARG:HG2	4:H:304:GOL:H31	2.00	0.44
4:I:302:GOL:O3	5:I:558:HOH:O	2.00	0.43
2:H:12:VAL:HG11	2:H:82(C):LEU:HD13	2.00	0.43
1:M:194:GLN:HG2	1:M:203:GLU:HG3	1.99	0.43
1:M:3:VAL:HG23	5:M:547:HOH:O	2.19	0.42
2:H:171:GLN:NE2	5:H:681:HOH:O	2.53	0.42
1:M:167:GLN:OE1	1:M:173:ALA:HB2	2.20	0.42
2:I:64:LYS:HA	2:I:64:LYS:HD2	1.87	0.42
2:H:71:ARG:HG3	4:H:304:GOL:H12	2.01	0.41
2:H:126:PRO:HG3	5:H:605:HOH:O	2.21	0.41
1:M:54:ARG:HD3	1:M:62:PHE:O	2.19	0.41
5:M:421:HOH:O	4:I:301:GOL:H11	2.20	0.41
2:H:36:TRP:CE2	2:H:80:LEU:HB2	2.55	0.41
1:L:19:VAL:HG13	1:L:78:LEU:HD11	2.01	0.41

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:L:685:HOH:O	5:H:691:HOH:O[4_546]	1.86	0.34
5:L:681:HOH:O	5:I:600:HOH:O[3_556]	1.88	0.32
5:L:630:HOH:O	5:Q:412:HOH:O[3_556]	1.94	0.26
5:H:676:HOH:O	5:M:591:HOH:O[3_556]	2.02	0.18
5:L:672:HOH:O	5:I:601:HOH:O[3_556]	2.05	0.15
5:H:643:HOH:O	5:M:588:HOH:O[3_556]	2.09	0.11
2:I:3:GLN:NE2	5:I:503:HOH:O[4_445]	2.11	0.09
1:L:30:ASN:ND2	5:L:583:HOH:O[4_556]	2.13	0.07
5:H:629:HOH:O	5:M:561:HOH:O[3_556]	2.16	0.04

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	214/216 (99%)	203 (95%)	11 (5%)	0	100	100
1	M	211/216 (98%)	204 (97%)	6 (3%)	1 (0%)	38	19
2	H	229/231 (99%)	226 (99%)	3 (1%)	0	100	100
2	I	229/231 (99%)	222 (97%)	6 (3%)	1 (0%)	43	25
3	P	12/14 (86%)	12 (100%)	0	0	100	100
3	Q	12/14 (86%)	12 (100%)	0	0	100	100
All	All	907/922 (98%)	879 (97%)	26 (3%)	2 (0%)	56	38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	I	129	ARG
1	M	151	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	180/180 (100%)	178 (99%)	2 (1%)	84	77
1	M	177/180 (98%)	174 (98%)	3 (2%)	73	61
2	H	195/196 (100%)	191 (98%)	4 (2%)	66	51
2	I	196/196 (100%)	191 (97%)	5 (3%)	59	41
3	P	11/11 (100%)	11 (100%)	0	100	100
3	Q	11/11 (100%)	11 (100%)	0	100	100
All	All	770/774 (100%)	756 (98%)	14 (2%)	71	58

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	95(C)	ASP
1	L	103	LYS
2	H	76	ASN
2	H	127	CYS
2	H	131	THR
2	H	138	LEU
1	M	95(C)	ASP
1	M	163	THR
1	M	168	SER
2	I	1	GLU
2	I	105	LYS
2	I	129	ARG
2	I	165	THR
2	I	199	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	30	ASN
1	L	126	GLN
2	H	76	ASN
1	M	126	GLN
2	I	76	ASN
2	I	164	HIS
2	I	199	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	H	301	-	5,5,5	0.12	0	5,5,5	0.70	0
4	GOL	H	302	-	5,5,5	0.29	0	5,5,5	0.64	0
4	GOL	H	303	-	5,5,5	0.30	0	5,5,5	0.53	0
4	GOL	H	304	-	5,5,5	0.40	0	5,5,5	0.67	0
4	GOL	H	305	-	5,5,5	0.37	0	5,5,5	0.31	0
4	GOL	H	306	-	5,5,5	0.60	0	5,5,5	0.35	0
4	GOL	I	301	-	5,5,5	0.50	0	5,5,5	0.69	0
4	GOL	I	302	-	5,5,5	0.30	0	5,5,5	0.50	0
4	GOL	L	301	-	5,5,5	0.52	0	5,5,5	0.70	0
4	GOL	L	302	-	5,5,5	0.49	0	5,5,5	0.52	0
4	GOL	L	303	-	3,4,5	0.20	0	2,4,5	0.46	0
4	GOL	L	304	-	5,5,5	0.30	0	5,5,5	0.29	0
4	GOL	L	305	-	5,5,5	0.35	0	5,5,5	0.43	0
4	GOL	L	306	-	5,5,5	0.25	0	5,5,5	0.40	0
4	GOL	M	301	-	5,5,5	0.25	0	5,5,5	0.28	0
4	GOL	M	302	-	5,5,5	0.32	0	5,5,5	0.33	0
4	GOL	M	303	-	5,5,5	0.35	0	5,5,5	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	H	301	-	-	0/4/4/4	0/0/0/0
4	GOL	H	302	-	-	0/4/4/4	0/0/0/0
4	GOL	H	303	-	-	0/4/4/4	0/0/0/0
4	GOL	H	304	-	-	0/4/4/4	0/0/0/0
4	GOL	H	305	-	-	0/4/4/4	0/0/0/0
4	GOL	H	306	-	-	0/4/4/4	0/0/0/0
4	GOL	I	301	-	-	0/4/4/4	0/0/0/0
4	GOL	I	302	-	-	0/4/4/4	0/0/0/0
4	GOL	L	301	-	-	0/4/4/4	0/0/0/0
4	GOL	L	302	-	-	0/4/4/4	0/0/0/0
4	GOL	L	303	-	-	0/2/2/4	0/0/0/0
4	GOL	L	304	-	-	0/4/4/4	0/0/0/0
4	GOL	L	305	-	-	0/4/4/4	0/0/0/0
4	GOL	L	306	-	-	0/4/4/4	0/0/0/0
4	GOL	M	301	-	-	0/4/4/4	0/0/0/0
4	GOL	M	302	-	-	0/4/4/4	0/0/0/0
4	GOL	M	303	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	216/216 (100%)	0.08	5 (2%) 57 51	7, 20, 44, 66	0
1	M	213/216 (98%)	0.07	11 (5%) 26 20	13, 30, 51, 70	0
2	H	231/231 (100%)	-0.03	6 (2%) 53 46	8, 23, 43, 55	0
2	I	231/231 (100%)	-0.02	11 (4%) 29 23	14, 28, 50, 101	0
3	P	14/14 (100%)	0.62	0 100 100	9, 13, 31, 32	0
3	Q	14/14 (100%)	0.02	1 (7%) 16 11	17, 24, 34, 37	0
All	All	919/922 (99%)	0.03	34 (3%) 41 32	7, 26, 47, 101	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	131	THR	7.7
2	I	130	SER	6.6
2	H	127	CYS	5.2
1	L	211	CYS	5.1
2	I	127	CYS	4.4
1	M	209	THR	4.1
1	L	1	GLN	4.1
2	I	126	PRO	3.9
1	L	209	THR	3.9
2	I	129	ARG	3.9
2	H	132	SER	3.8
2	H	128	SER	3.6
1	L	210	GLU	3.6
2	H	131	THR	3.5
2	I	213	LEU	3.4
1	M	155	VAL	3.4
1	M	189	ARG	3.4
3	Q	320	CYS	3.3
2	I	128	SER	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	M	156	LYS	3.2
1	M	186	LYS	3.2
2	I	1	GLU	3.0
2	I	133	GLY	2.9
1	M	187	SER	2.9
2	I	132	SER	2.7
2	H	130	SER	2.5
1	M	150	ALA	2.4
1	M	151	ASP	2.2
1	M	183	GLU	2.2
2	I	100(C)	VAL	2.2
1	L	168	SER	2.2
1	M	154	PRO	2.2
2	H	126	PRO	2.1
1	M	188	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	L	303	5/6	0.20	-	10,33,35,36	0
4	GOL	H	303	6/6	0.10	-	14,20,22,28	0
4	GOL	H	304	6/6	0.24	-	21,22,27,29	0
4	GOL	M	302	6/6	0.12	-	39,41,43,48	0
4	GOL	H	301	6/6	0.10	-	17,27,32,33	0
4	GOL	L	304	6/6	0.11	-	19,24,31,37	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	I	302	6/6	0.10	-	33,34,43,46	0
4	GOL	M	301	6/6	0.14	-	31,34,37,40	0
4	GOL	L	301	6/6	0.25	-	26,27,40,47	0
4	GOL	I	301	6/6	0.19	-	23,33,41,45	0
4	GOL	L	302	6/6	0.15	-	29,32,34,37	0
4	GOL	L	305	6/6	0.21	-	22,29,40,41	0
4	GOL	H	305	6/6	0.08	-	26,33,38,39	0
4	GOL	L	306	6/6	0.15	-	21,27,41,42	0
4	GOL	H	306	6/6	0.17	-	26,30,35,38	0
4	GOL	H	302	6/6	0.17	-	21,30,34,40	0
4	GOL	M	303	6/6	0.17	-	37,43,47,55	0

## 6.5 Other polymers

There are no such residues in this entry.