



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 01:49 AM GMT

PDB ID : 1M3D
Title : Structure of Type IV Collagen NC1 Domains
Authors : Sundaramoorthy, M.; Meiyappan, M.; Todd, P.; Hudson, B.G.
Deposited on : 2002-06-27
Resolution : 2.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	229	
1	B	229	
1	D	229	
1	E	229	
1	G	229	
1	H	229	
1	J	229	
1	K	229	
2	C	227	
2	F	227	
2	I	227	
2	L	227	

The following table lists non-polymeric compounds that are outliers for geometric or electron-

density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	GOL	A	4001	X	X
5	GOL	B	4002	X	-
5	GOL	D	4003	X	-
5	GOL	E	4004	X	X
5	GOL	G	4005	X	X
5	GOL	H	4006	X	X
5	GOL	J	4007	X	X
5	GOL	K	4008	X	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 22023 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type IV Collagen Noncollagenous Domain- Alpha1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	2	0
			1735	1093	303	318	21			
1	B	224	Total	C	N	O	S	0	2	0
			1747	1098	307	321	21			
1	D	225	Total	C	N	O	S	0	1	0
			1743	1098	304	320	21			
1	E	224	Total	C	N	O	S	0	0	0
			1734	1091	303	319	21			
1	G	225	Total	C	N	O	S	0	2	0
			1748	1102	305	320	21			
1	H	224	Total	C	N	O	S	0	1	0
			1738	1095	303	319	21			
1	J	223	Total	C	N	O	S	0	0	0
			1724	1085	300	318	21			
1	K	224	Total	C	N	O	S	0	0	0
			1738	1093	304	320	21			

- Molecule 2 is a protein called Type IV Collagen Noncollagenous Domain- Alpha2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	222	Total	C	N	O	S	0	0	0
			1720	1089	291	321	19			
2	F	223	Total	C	N	O	S	0	0	0
			1724	1091	292	322	19			
2	I	222	Total	C	N	O	S	0	0	0
			1720	1089	291	321	19			
2	L	222	Total	C	N	O	S	0	1	0
			1726	1093	293	321	19			

- Molecule 3 is LUTETIUM (III) ION (three-letter code: LU) (formula: Lu).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	I	1	Total Lu 1 1	0	0
3	L	1	Total Lu 1 1	0	0
3	C	1	Total Lu 1 1	0	0
3	F	1	Total Lu 1 1	0	0

- Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	3	Total Br 3 3	0	0
4	J	2	Total Br 2 2	0	0
4	D	1	Total Br 1 1	0	0
4	K	3	Total Br 3 3	0	0
4	E	2	Total Br 2 2	0	0
4	H	4	Total Br 4 4	0	0
4	B	5	Total Br 5 5	0	0
4	I	3	Total Br 3 3	0	0
4	C	4	Total Br 4 4	0	0
4	A	4	Total Br 4 4	0	0
4	L	3	Total Br 3 3	0	0
4	F	2	Total Br 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	D	1	Total	C	O	0	0
			6	3	3		
5	E	1	Total	C	O	0	0
			6	3	3		
5	G	1	Total	C	O	0	0
			6	3	3		
5	H	1	Total	C	O	0	0
			6	3	3		
5	J	1	Total	C	O	0	0
			6	3	3		
5	K	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	108	Total	O	0	0
			108	108		
6	B	87	Total	O	0	0
			87	87		
6	C	103	Total	O	0	0
			103	103		
6	D	103	Total	O	0	0
			103	103		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	E	83	Total 83	O 83	0	0
6	F	111	Total 111	O 111	0	0
6	G	85	Total 85	O 85	0	0
6	H	88	Total 88	O 88	0	0
6	I	99	Total 99	O 99	0	0
6	J	94	Total 94	O 94	0	0
6	K	85	Total 85	O 85	0	0
6	L	92	Total 92	O 92	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain A:




- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain B:



- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain D:




- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain E:




- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain G: 




- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain H: 




- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain J: 




- Molecule 1: Type IV Collagen Noncollagenous Domain- Alpha1

Chain K: 




- Molecule 2: Type IV Collagen Noncollagenous Domain- Alpha2

Chain C: 




- Molecule 2: Type IV Collagen Noncollagenous Domain- Alpha2

Chain F: 



• Molecule 2: Type IV Collagen Noncollagenous Domain- Alpha2

Chain I:



• Molecule 2: Type IV Collagen Noncollagenous Domain- Alpha2

Chain L:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.07Å 137.96Å 127.13Å 90.00° 90.31° 90.00°	Depositor
Resolution (Å)	8.00 – 2.00 93.49 – 1.99	Depositor EDS
% Data completeness (in resolution range)	95.9 (8.00-2.00) 97.2 (93.49-1.99)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.45 (at 2.00Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.169 , 0.197 0.174 , 0.199	Depositor DCC
R_{free} test set	9094 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	14.8	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 40.5	EDS
Estimated twinning fraction	0.018 for h,-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 184668 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	22023	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, LU, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1793	0.60	0/2435
1	B	0.32	0/1805	0.59	0/2452
1	D	0.32	0/1798	0.59	0/2443
1	E	0.32	0/1785	0.59	0/2425
1	G	0.31	0/1806	0.59	0/2453
1	H	0.31	0/1793	0.58	0/2436
1	J	0.31	0/1774	0.58	0/2410
1	K	0.31	0/1788	0.59	0/2429
2	C	0.31	0/1771	0.61	0/2410
2	F	0.32	0/1775	0.62	0/2415
2	I	0.30	0/1771	0.61	0/2410
2	L	0.31	0/1782	0.61	0/2425
All	All	0.31	0/21441	0.60	0/29143

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1735	0	1657	62	0
1	B	1747	0	1668	70	0
1	D	1743	0	1657	63	0
1	E	1734	0	1648	50	0
1	G	1748	0	1670	54	0
1	H	1738	0	1655	73	0
1	J	1724	0	1641	55	0
1	K	1738	0	1658	71	0
2	C	1720	0	1629	61	0
2	F	1724	0	1632	65	0
2	I	1720	0	1629	59	0
2	L	1726	0	1632	43	0
3	C	1	0	0	0	0
3	F	1	0	0	0	0
3	I	1	0	0	0	0
3	L	1	0	0	0	0
4	A	4	0	0	3	0
4	B	5	0	0	5	0
4	C	4	0	0	1	0
4	D	1	0	0	0	0
4	E	2	0	0	3	0
4	F	2	0	0	1	0
4	G	3	0	0	2	0
4	H	4	0	0	2	0
4	I	3	0	0	1	0
4	J	2	0	0	2	0
4	K	3	0	0	4	0
4	L	3	0	0	1	0
5	A	6	0	4	0	0
5	B	6	0	4	0	0
5	D	6	0	4	0	0
5	E	6	0	4	0	0
5	G	6	0	4	0	0
5	H	6	0	4	0	0
5	J	6	0	4	0	0
5	K	6	0	4	0	0
6	A	108	0	0	1	0
6	B	87	0	0	4	0
6	C	103	0	0	3	0
6	D	103	0	0	1	0
6	E	83	0	0	3	0
6	F	111	0	0	2	0
6	G	85	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	H	88	0	0	3	0
6	I	99	0	0	2	0
6	J	94	0	0	1	0
6	K	85	0	0	1	0
6	L	92	0	0	2	0
All	All	22023	0	19808	588	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

The worst 5 of 588 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:L:98:ILE:HG22	2:L:102:ILE:HD11	1.29	1.10
1:J:101:ILE:HG22	1:J:105:ILE:HD11	1.37	1.04
2:F:214:ARG:HA	2:F:217:ILE:HD12	1.43	1.00
1:B:90:PRO:HG2	1:B:96:ILE:HD11	1.44	0.99
1:D:10:ARG:HH12	1:H:129:GLN:NE2	1.62	0.95

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/229 (97%)	215 (96%)	8 (4%)	0	100	100
1	B	224/229 (98%)	216 (96%)	8 (4%)	0	100	100
1	D	224/229 (98%)	216 (96%)	7 (3%)	1 (0%)	43	36
1	E	222/229 (97%)	214 (96%)	8 (4%)	0	100	100
1	G	225/229 (98%)	216 (96%)	9 (4%)	0	100	100
1	H	223/229 (97%)	213 (96%)	10 (4%)	0	100	100
1	J	221/229 (96%)	213 (96%)	8 (4%)	0	100	100
1	K	222/229 (97%)	215 (97%)	7 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	220/227 (97%)	209 (95%)	9 (4%)	2 (1%)	25	14
2	F	221/227 (97%)	211 (96%)	9 (4%)	1 (0%)	38	29
2	I	220/227 (97%)	207 (94%)	9 (4%)	4 (2%)	13	5
2	L	221/227 (97%)	210 (95%)	10 (4%)	1 (0%)	38	29
All	All	2666/2740 (97%)	2555 (96%)	102 (4%)	9 (0%)	50	44

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	F	175	GLY
2	I	21	VAL
2	I	211	GLY
2	L	175	GLY
2	I	175	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/195 (98%)	183 (96%)	8 (4%)	40	34
1	B	193/195 (99%)	190 (98%)	3 (2%)	75	77
1	D	191/195 (98%)	184 (96%)	7 (4%)	45	40
1	E	190/195 (97%)	185 (97%)	5 (3%)	59	58
1	G	192/195 (98%)	188 (98%)	4 (2%)	66	67
1	H	191/195 (98%)	188 (98%)	3 (2%)	75	77
1	J	189/195 (97%)	183 (97%)	6 (3%)	51	47
1	K	191/195 (98%)	189 (99%)	2 (1%)	85	88
2	C	187/191 (98%)	182 (97%)	5 (3%)	57	56
2	F	187/191 (98%)	183 (98%)	4 (2%)	66	67
2	I	187/191 (98%)	181 (97%)	6 (3%)	51	47
2	L	188/191 (98%)	186 (99%)	2 (1%)	84	86
All	All	2277/2324 (98%)	2222 (98%)	55 (2%)	61	61

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	79	TYR
2	F	177	ARG
1	J	145	MET
1	E	102	ARG
1	E	227	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 97 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	153	GLN
1	G	156	GLN
2	L	38	GLN
2	F	221	GLN
1	G	45	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 48 ligands modelled in this entry, 40 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	4001	-	5,5,5	4.56	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	B	4002	-	5,5,5	4.55	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	D	4003	-	5,5,5	4.54	5 (100%)	5,5,5	5.75	3 (60%)
5	GOL	E	4004	-	5,5,5	4.54	5 (100%)	5,5,5	5.76	3 (60%)
5	GOL	G	4005	-	5,5,5	4.55	5 (100%)	5,5,5	5.72	3 (60%)
5	GOL	H	4006	-	5,5,5	4.57	5 (100%)	5,5,5	5.74	3 (60%)
5	GOL	J	4007	-	5,5,5	4.60	5 (100%)	5,5,5	5.74	3 (60%)
5	GOL	K	4008	-	5,5,5	4.56	5 (100%)	5,5,5	5.72	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	4001	-	-	0/4/4/4	0/0/0/0
5	GOL	B	4002	-	-	0/4/4/4	0/0/0/0
5	GOL	D	4003	-	-	0/4/4/4	0/0/0/0
5	GOL	E	4004	-	-	0/4/4/4	0/0/0/0
5	GOL	G	4005	-	-	0/4/4/4	0/0/0/0
5	GOL	H	4006	-	-	0/4/4/4	0/0/0/0
5	GOL	J	4007	-	-	0/4/4/4	0/0/0/0
5	GOL	K	4008	-	-	0/4/4/4	0/0/0/0

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	J	4007	GOL	C3-C2	-7.68	1.20	1.52
5	K	4008	GOL	C3-C2	-7.65	1.20	1.52
5	E	4004	GOL	C3-C2	-7.59	1.21	1.52
5	H	4006	GOL	C3-C2	-7.59	1.21	1.52
5	A	4001	GOL	C3-C2	-7.58	1.21	1.52

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	4003	GOL	O3-C3-C2	10.44	160.63	109.71
5	E	4004	GOL	O3-C3-C2	10.39	160.42	109.71
5	H	4006	GOL	O3-C3-C2	10.35	160.23	109.71
5	K	4008	GOL	O3-C3-C2	10.34	160.14	109.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	4002	GOL	O3-C3-C2	10.33	160.13	109.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/229 (97%)	-0.24	1 (0%) 90 91	6, 12, 25, 35	0
1	B	224/229 (97%)	-0.17	4 (1%) 65 66	6, 12, 29, 42	0
1	D	225/229 (98%)	-0.15	4 (1%) 65 66	7, 13, 27, 40	0
1	E	224/229 (97%)	-0.15	2 (0%) 81 82	6, 12, 29, 43	0
1	G	225/229 (98%)	-0.14	0 100 100	8, 15, 28, 39	0
1	H	224/229 (97%)	-0.11	4 (1%) 65 66	8, 14, 30, 41	0
1	J	223/229 (97%)	-0.13	2 (0%) 81 82	8, 14, 29, 39	0
1	K	224/229 (97%)	-0.10	5 (2%) 59 59	6, 14, 31, 45	0
2	C	222/227 (97%)	-0.13	3 (1%) 72 72	7, 13, 28, 39	0
2	F	223/227 (98%)	-0.07	3 (1%) 74 74	7, 12, 30, 41	0
2	I	222/227 (97%)	-0.05	7 (3%) 45 45	8, 15, 32, 44	0
2	L	222/227 (97%)	-0.12	1 (0%) 88 89	7, 14, 28, 37	0
All	All	2681/2740 (97%)	-0.13	36 (1%) 74 74	6, 14, 30, 45	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	212	LEU	7.0
2	I	212	LEU	5.2
2	C	21	VAL	4.2
2	C	22	GLY	4.0
1	E	4	HIS	3.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	GOL	H	4006	6/6	0.25	8.46	37,42,43,43	0
5	GOL	E	4004	6/6	0.25	7.40	25,34,35,36	0
5	GOL	A	4001	6/6	0.18	6.38	31,35,36,37	0
5	GOL	K	4008	6/6	0.28	5.35	43,46,46,47	0
5	GOL	J	4007	6/6	0.19	2.34	35,37,38,38	0
5	GOL	G	4005	6/6	0.26	2.12	38,42,44,45	0
5	GOL	B	4002	6/6	0.17	1.54	31,35,36,37	0
5	GOL	D	4003	6/6	0.20	1.17	42,42,42,43	0
4	BR	F	3013	1/1	0.05	-0.77	15,15,15,15	0
4	BR	D	3030	1/1	0.10	-1.05	39,39,39,39	1
4	BR	B	3012	1/1	0.06	-1.08	18,18,18,18	0
4	BR	H	3006	1/1	0.06	-1.18	17,17,17,17	0
4	BR	K	3009	1/1	0.06	-1.25	18,18,18,18	0
4	BR	B	3028	1/1	0.06	-1.43	29,29,29,29	1
4	BR	G	3031	1/1	0.08	-1.64	37,37,37,37	1
4	BR	K	3033	1/1	0.07	-1.71	46,46,46,46	1
4	BR	E	3003	1/1	0.05	-1.74	16,16,16,16	0
4	BR	B	3026	1/1	0.05	-2.03	32,32,32,32	1
4	BR	A	3004	1/1	0.06	-2.05	14,14,14,14	0
4	BR	C	3029	1/1	0.06	-2.06	30,30,30,30	1
4	BR	B	3034	1/1	0.08	-2.14	43,43,43,43	1
4	BR	A	3005	1/1	0.06	-2.16	14,14,14,14	0
4	BR	J	3032	1/1	0.09	-2.30	37,37,37,37	1
4	BR	H	3035	1/1	0.08	-2.49	53,53,53,53	1
4	BR	G	3016	1/1	0.05	-2.72	17,17,17,17	0
4	BR	E	3036	1/1	0.07	-2.81	53,53,53,53	1
3	LU	L	2004	1/1	0.06	-3.00	34,34,34,34	1
4	BR	F	3022	1/1	0.05	-3.04	20,20,20,20	1
4	BR	J	3008	1/1	0.06	-3.15	16,16,16,16	0
4	BR	A	3002	1/1	0.05	-3.19	15,15,15,15	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	LU	I	2003	1/1	0.04	-3.32	36,36,36,36	1
4	BR	I	3014	1/1	0.05	-3.34	18,18,18,18	0
4	BR	L	3017	1/1	0.06	-3.36	22,22,22,22	1
4	BR	K	3027	1/1	0.04	-3.39	34,34,34,34	1
3	LU	C	2001	1/1	0.03	-3.80	37,37,37,37	1
4	BR	G	3015	1/1	0.06	-4.29	15,15,15,15	0
4	BR	L	3010	1/1	0.04	-4.77	18,18,18,18	0
4	BR	A	3001	1/1	0.07	-5.17	16,16,16,16	0
4	BR	C	3019	1/1	0.04	-5.55	19,19,19,19	1
4	BR	I	3023	1/1	0.04	-5.63	24,24,24,24	1
4	BR	C	3018	1/1	0.04	-7.01	21,21,21,21	1
3	LU	F	2002	1/1	0.05	-7.30	42,42,42,42	1
4	BR	I	3024	1/1	0.04	-7.82	23,23,23,23	1
4	BR	C	3007	1/1	0.04	-7.84	15,15,15,15	0
4	BR	L	3025	1/1	0.04	-8.23	22,22,22,22	1
4	BR	B	3021	1/1	0.05	-8.40	23,23,23,23	1
4	BR	H	3011	1/1	0.05	-11.54	17,17,17,17	0
4	BR	H	3020	1/1	0.04	-14.69	20,20,20,20	1

6.5 Other polymers ⓘ

There are no such residues in this entry.