



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 11:30 AM GMT

PDB ID : 1M4Z
Title : Crystal structure of the N-terminal BAH domain of Orc1p
Authors : Zhang, Z.; Hayashi, M.K.; Merkel, O.; Stillman, B.; Xu, R.-M.
Deposited on : 2002-07-05
Resolution : 2.20 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2938 (2.20-2.20)
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	238	
1	B	238	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3586 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ORIGIN RECOGNITION COMPLEX SUBUNIT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	203	Total	C	N	O	S	0	0	0
			1658	1059	278	315	6			
1	B	203	Total	C	N	O	S	0	0	0
			1659	1059	278	316	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	ALA	-	EXPRESSION TAG	UNP P54784
A	-1	MET	-	EXPRESSION TAG	UNP P54784
A	0	THR	-	EXPRESSION TAG	UNP P54784
B	-2	ALA	-	EXPRESSION TAG	UNP P54784
B	-1	MET	-	EXPRESSION TAG	UNP P54784
B	0	THR	-	EXPRESSION TAG	UNP P54784

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	144	Total	O	0	0
			144	144		
3	B	122	Total	O	0	0
			122	122		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ORIGIN RECOGNITION COMPLEX SUBUNIT 1

Chain A: 



• Molecule 1: ORIGIN RECOGNITION COMPLEX SUBUNIT 1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	128.26Å 61.05Å 67.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 46.58 – 2.19	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.20) 95.2 (46.58-2.19)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.11 (at 2.20Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.200 , 0.225 0.203 , 0.226	Depositor DCC
R_{free} test set	2099 reflections (7.91%)	DCC
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.871	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 26605 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3586	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1692	0.61	0/2294
1	B	0.33	0/1693	0.67	2/2295 (0.1%)
All	All	0.33	0/3385	0.64	2/4589 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	ASP	N-CA-C	-6.74	92.81	111.00
1	B	107	PRO	N-CA-C	5.86	127.33	112.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1658	0	1630	30	0
1	B	1659	0	1630	47	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	144	0	0	2	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	122	0	0	2	0
All	All	3586	0	3260	76	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

All (76) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:103:ARG:HA	1:B:107:PRO:HG2	1.23	1.17
1:B:103:ARG:CA	1:B:107:PRO:HG2	2.02	0.89
1:B:111:ILE:HG22	1:B:112:LEU:N	1.96	0.81
1:B:106:ASN:HB2	1:B:107:PRO:HD3	1.66	0.77
1:B:111:ILE:HG22	1:B:112:LEU:H	1.50	0.75
1:B:102:TYR:O	1:B:107:PRO:HD2	1.94	0.67
1:A:174:ARG:HD2	3:A:568:HOH:O	2.00	0.61
1:B:113:ASN:O	1:B:114:ARG:C	2.40	0.60
1:B:111:ILE:CG2	1:B:112:LEU:H	2.15	0.59
1:B:15:THR:HG22	1:B:39:HIS:CD2	2.36	0.59
1:A:40:TYR:HA	1:A:74:LEU:CD1	2.33	0.59
1:B:15:THR:HG22	1:B:39:HIS:HD2	1.68	0.58
1:B:74:LEU:HD21	1:B:85:LEU:HD13	1.85	0.58
1:B:102:TYR:O	1:B:107:PRO:CD	2.52	0.58
1:B:16:THR:HB	1:B:20:GLY:HA2	1.86	0.57
1:B:214:PRO:O	1:B:215:SER:HB2	2.03	0.57
1:B:111:ILE:CG2	1:B:112:LEU:N	2.66	0.57
1:A:14:ILE:HD12	1:A:40:TYR:CE1	2.40	0.56
1:B:161:LEU:HD21	3:B:605:HOH:O	2.06	0.56
1:B:111:ILE:O	1:B:112:LEU:HD23	2.05	0.56
1:B:74:LEU:CD2	1:B:85:LEU:HD13	2.37	0.55
1:A:43:ARG:HD2	1:A:46:ASP:OD1	2.06	0.55
1:A:178:GLU:HG3	1:A:183:LYS:HD2	1.89	0.55
1:B:115:PRO:O	1:B:119:TYR:HD1	1.90	0.55
1:B:106:ASN:N	1:B:107:PRO:CD	2.70	0.54
1:A:16:THR:HB	1:A:20:GLY:O	2.08	0.54
1:A:79:LEU:HD22	1:B:116:LEU:HD22	1.90	0.53
1:B:77:ASN:HB3	1:B:80:ASN:HD21	1.74	0.52
1:B:106:ASN:H	1:B:107:PRO:HD2	1.74	0.52
1:B:16:THR:HB	1:B:20:GLY:CA	2.40	0.51
1:B:106:ASN:CB	1:B:107:PRO:HD3	2.39	0.50
1:A:110:ASN:ND2	3:A:601:HOH:O	2.44	0.50
1:B:43:ARG:HD3	1:B:45:SER:OG	2.12	0.50
1:A:199:VAL:HG23	1:A:204:ALA:HB2	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:42:LYS:HD2	1:A:49:LYS:HE2	1.94	0.49
1:B:114:ARG:O	1:B:115:PRO:O	2.31	0.49
1:A:16:THR:HG22	1:A:21:ASN:HA	1.95	0.48
1:B:106:ASN:N	1:B:107:PRO:HD2	2.27	0.48
1:A:74:LEU:CD2	1:A:85:LEU:HD13	2.43	0.47
1:A:80:ASN:C	1:A:80:ASN:HD22	2.15	0.47
1:B:96:VAL:O	1:B:98:PRO:HD3	2.14	0.47
1:B:176:ILE:HB	1:B:187:ILE:HD13	1.97	0.47
1:A:200:GLU:HG3	1:A:201:PRO:HD2	1.96	0.46
1:A:16:THR:HB	1:A:21:ASN:HA	1.98	0.46
1:A:80:ASN:ND2	1:A:82:VAL:H	2.14	0.45
1:B:80:ASN:C	1:B:80:ASN:HD22	2.19	0.45
1:B:17:ASP:C	1:B:19:GLN:H	2.20	0.45
1:B:17:ASP:OD1	1:B:19:GLN:N	2.50	0.45
1:B:196:ILE:HD13	1:B:204:ALA:HB1	1.98	0.45
1:A:213:LEU:O	1:A:213:LEU:HD12	2.17	0.44
1:B:106:ASN:H	1:B:107:PRO:CD	2.28	0.43
1:B:102:TYR:CD2	1:B:119:TYR:HE2	2.37	0.43
1:A:131:GLU:OE2	1:A:174:ARG:HD3	2.18	0.43
1:B:80:ASN:ND2	1:B:82:VAL:H	2.16	0.43
1:A:46:ASP:OD1	1:A:48:ILE:HG12	2.18	0.43
1:A:196:ILE:HD11	1:A:208:LEU:HD11	2.01	0.43
1:A:55:SER:HA	1:A:69:TYR:O	2.19	0.43
1:B:105:PHE:HZ	1:B:184:PHE:HB2	1.84	0.43
1:B:165:VAL:HG22	3:B:593:HOH:O	2.19	0.43
1:B:166:ASP:OD1	1:B:169:ARG:NH1	2.52	0.43
1:B:209:LYS:HB3	1:B:209:LYS:HE2	1.82	0.42
1:B:43:ARG:HD2	1:B:46:ASP:OD1	2.18	0.42
1:B:96:VAL:HG13	1:B:184:PHE:HZ	1.83	0.42
1:B:119:TYR:N	1:B:119:TYR:CD1	2.86	0.42
1:A:16:THR:CB	1:A:21:ASN:HA	2.49	0.42
1:A:162:LYS:HE3	1:A:162:LYS:HB2	1.92	0.41
1:A:199:VAL:CG2	1:A:204:ALA:HB2	2.50	0.41
1:B:105:PHE:CZ	1:B:184:PHE:HB2	2.55	0.41
1:B:17:ASP:OD1	1:B:19:GLN:HB2	2.20	0.41
1:A:192:VAL:O	1:A:196:ILE:HG12	2.19	0.41
1:A:16:THR:CG2	1:A:21:ASN:HA	2.50	0.41
1:A:182:GLU:O	1:A:183:LYS:HB2	2.21	0.41
1:A:17:ASP:O	1:A:19:GLN:N	2.54	0.41
1:A:75:ARG:HB3	1:A:84:GLU:HB2	2.03	0.40
1:B:16:THR:HB	1:B:20:GLY:C	2.41	0.40
1:A:161:LEU:O	1:A:162:LYS:C	2.60	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:A:582:HOH:O	3:A:582:HOH:O[2_665]	1.61	0.59

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/238 (84%)	185 (93%)	12 (6%)	2 (1%)	22	18
1	B	199/238 (84%)	175 (88%)	16 (8%)	8 (4%)	5	1
All	All	398/476 (84%)	360 (90%)	28 (7%)	10 (2%)	9	4

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	108	ASP
1	B	115	PRO
1	A	18	GLU
1	B	10	GLY
1	B	103	ARG
1	B	116	LEU
1	B	21	ASN
1	B	110	ASN
1	A	162	LYS
1	B	17	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/209 (86%)	177 (99%)	2 (1%)	84	92
1	B	180/209 (86%)	177 (98%)	3 (2%)	73	84
All	All	359/418 (86%)	354 (99%)	5 (1%)	78	88

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	TYR
1	A	80	ASN
1	B	80	ASN
1	B	107	PRO
1	B	200	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	80	ASN
1	A	81	ASN
1	A	130	ASN
1	B	39	HIS
1	B	80	ASN
1	B	81	ASN
1	B	117	ASN
1	B	120	ASN
1	B	130	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	203/238 (85%)	-0.09	6 (2%) 48 48	16, 32, 64, 93	0
1	B	203/238 (85%)	0.50	22 (10%) 6 6	18, 38, 86, 98	0
All	All	406/476 (85%)	0.21	28 (6%) 17 16	16, 35, 80, 98	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	108	ASP	10.2
1	B	110	ASN	10.1
1	B	116	LEU	8.4
1	B	111	ILE	7.7
1	B	113	ASN	6.7
1	B	107	PRO	5.9
1	A	19	GLN	5.9
1	B	115	PRO	5.8
1	B	109	ALA	5.1
1	B	114	ARG	5.1
1	B	215	SER	3.8
1	B	121	LYS	3.5
1	A	40	TYR	3.5
1	B	102	TYR	3.3
1	B	-1	MET	3.2
1	B	106	ASN	3.1
1	B	19	GLN	3.1
1	B	162	LYS	3.0
1	B	105	PHE	2.9
1	A	37	THR	2.9
1	B	119	TYR	2.8
1	B	112	LEU	2.7
1	B	40	TYR	2.7
1	A	214	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	-2	ALA	2.6
1	B	18	GLU	2.4
1	A	20	GLY	2.4
1	B	125	GLU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	MN	A	501	1/1	0.20	0.42	23,23,23,23	1
2	MN	B	502	1/1	0.06	-2.54	49,49,49,49	0
2	MN	A	503	1/1	0.03	-3.89	47,47,47,47	0

6.5 Other polymers ⓘ

There are no such residues in this entry.