



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 02:57 PM GMT

PDB ID : 1ML6
Title : Crystal Structure of mGSTA2-2 in Complex with the Glutathione Conjugate of Benzo[a]pyrene-7(R),8(S)-Diol-9(S),10(R)-Epoxide
Authors : Gu, Y.; Xiao, B.; Wargo, H.L.; Bucher, M.H.; Singh, S.V.; Ji, X.
Deposited on : 2002-08-30
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	221	
1	B	221	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3986 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutathione S-Transferase GT41A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	0	0
			1783	1149	303	322	9			
1	B	220	Total	C	N	O	S	0	0	0
			1783	1149	303	322	9			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	111	CSO	CYS	MODIFIED RESIDUE	UNP P10648
B	411	CSO	CYS	MODIFIED RESIDUE	UNP P10648

- Molecule 2 is 2-AMINO-4-[1-(CARBOXYMETHYL-CARBAMOYL)-2-(9-HYDROXY-7,8-DIOXO-7,8,9,10-TETRAHYDRO-BENZO[DEF]CHRYSEN-10-YLSULFANYL)-ETHYLCARBAMOYL]-BUTYRICACID (three-letter code: GBX) (formula: $C_{30}H_{27}N_3O_9S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			43	30	3	9	1		
2	B	1	Total	C	N	O	S	0	0
			43	30	3	9	1		

- Molecule 3 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	3	1		
3	B	1	Total	C	O	0	0
			4	3	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	157	Total	O	0	0
			157	157		
4	B	169	Total	O	0	0
			169	169		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glutathione S-Transferase GT41A

Chain A: 



• Molecule 1: Glutathione S-Transferase GT41A

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.07Å 93.90Å 52.60Å 90.00° 92.28° 90.00°	Depositor
Resolution (Å)	26.28 – 1.90 26.28 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (26.28-1.90) 99.7 (26.28-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 1.91Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.200 , 0.249 0.203 , 0.251	Depositor DCC
R_{free} test set	1882 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	22.5	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.4	EDS
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 37774 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3986	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, IPA, GBX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1809	0.62	0/2431
1	B	0.41	0/1809	0.63	0/2431
All	All	0.41	0/3618	0.63	0/4862

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1783	0	1847	16	0
1	B	1783	0	1848	24	0
2	A	43	0	25	0	0
2	B	43	0	25	2	0
3	A	4	0	8	0	0
3	B	4	0	8	0	0
4	A	157	0	0	1	0
4	B	169	0	0	5	0
All	All	3986	0	3761	39	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (39) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:512:ILE:O	1:B:516:ARG:HG3	1.80	0.81
1:B:514:GLU:O	1:B:518:VAL:HG23	1.91	0.70
2:B:530:GBX:HB22	4:B:823:HOH:O	2.02	0.59
1:B:437:LYS:NZ	4:B:804:HOH:O	2.36	0.58
1:A:214:GLU:O	1:A:218:VAL:HG23	2.05	0.56
1:A:27:VAL:HG21	1:A:78:TYR:CZ	2.43	0.54
1:A:160:LEU:HD22	1:A:187:ILE:HG22	1.90	0.54
1:A:208:ASP:O	1:A:212:ILE:HG13	2.09	0.53
1:B:510:LYS:O	1:B:514:GLU:HG3	2.11	0.50
1:B:516:ARG:HA	1:B:521:PHE:HB2	1.96	0.47
1:B:340:LEU:HG	1:B:344:LYS:HE3	1.96	0.47
1:A:216:ARG:HA	1:A:221:PHE:HB2	1.96	0.47
1:A:65:VAL:O	1:A:66:GLN:HB2	2.16	0.46
1:B:475:THR:HB	1:B:476:PRO:HD3	1.96	0.46
1:A:170:ASP:HB3	1:A:173:LEU:CD1	2.46	0.45
1:A:94:TYR:HB2	1:A:155:VAL:HG21	1.99	0.45
1:B:307:HIS:CE1	1:B:332:LYS:HD2	2.51	0.45
1:B:521:PHE:CE1	2:B:530:GBX:SG2	3.10	0.44
1:A:63:LYS:HG3	4:A:772:HOH:O	2.16	0.44
1:B:365:VAL:O	1:B:366:GLN:HB2	2.17	0.44
1:B:349:LEU:HB2	1:B:352:ASP:HA	1.99	0.43
1:B:508:ASP:O	1:B:512:ILE:HG13	2.19	0.43
1:B:461:GLU:HG2	1:B:465:TYR:CE2	2.54	0.43
1:A:7:HIS:CE1	1:A:32:LYS:HD2	2.53	0.43
1:A:85:MET:CE	1:B:377:LYS:HB2	2.49	0.43
1:B:307:HIS:HE1	1:B:332:LYS:HD2	1.84	0.43
1:B:474:LEU:O	1:B:481:LYS:HE2	2.19	0.43
1:B:498:GLN:NE2	4:B:838:HOH:O	2.52	0.43
1:A:77:LYS:N	1:B:385:MET:HE3	2.34	0.42
1:B:348:ASN:ND2	4:B:730:HOH:O	2.52	0.42
1:B:402:THR:HG21	4:B:707:HOH:O	2.20	0.42
1:B:398:ILE:HG13	1:B:435:PHE:CZ	2.55	0.42
1:A:15:MET:HE2	1:A:55:PRO:HB3	2.00	0.42
1:B:375:ALA:CB	1:B:454:ARG:HD2	2.50	0.42
1:B:346:ASP:HB2	1:B:348:ASN:ND2	2.34	0.41
1:B:483:PHE:CZ	1:B:487:ILE:HG13	2.56	0.41
1:A:175:THR:HB	1:A:176:PRO:HD3	2.03	0.41
1:A:40:LEU:HG	1:A:44:LYS:HE3	2.03	0.41
1:A:101:LEU:HD23	1:A:162:LEU:HD21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/221 (98%)	208 (96%)	9 (4%)	0	100	100
1	B	217/221 (98%)	209 (96%)	8 (4%)	0	100	100
All	All	434/442 (98%)	417 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/193 (100%)	187 (97%)	6 (3%)	52	41
1	B	193/193 (100%)	189 (98%)	4 (2%)	66	59
All	All	386/386 (100%)	376 (97%)	10 (3%)	59	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	TYR
1	A	107	GLN
1	A	150	ASN
1	A	169	LEU
1	A	205	PRO
1	A	207	LEU
1	B	381	TYR
1	B	407	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	450	ASN
1	B	469	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	GLN
1	A	150	ASN
1	A	198	GLN
1	B	444	GLN
1	B	450	ASN
1	B	498	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	111	1	6,6,7	7.69	3 (50%)	3,6,8	1.47	1 (33%)
1	CSO	B	411	1	6,6,7	7.39	3 (50%)	3,6,8	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	111	1	-	0/2/5/7	0/0/0/0
1	CSO	B	411	1	-	0/2/5/7	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	111	CSO	O-C	18.03	1.23	1.11
1	B	411	CSO	O-C	17.23	1.23	1.11
1	B	411	CSO	OD-SG	4.90	1.83	1.62
1	A	111	CSO	OD-SG	4.77	1.82	1.62
1	B	411	CSO	CA-C	2.59	1.53	1.48
1	A	111	CSO	CA-C	2.59	1.53	1.48

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	111	CSO	C-CA-N	-2.34	111.49	113.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GBX	A	230	-	47,47,47	3.11	10 (21%)	69,69,69	1.43	5 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	IPA	A	235	-	3,3,3	0.58	0	3,3,3	0.39	0
3	IPA	B	236	-	3,3,3	0.62	0	3,3,3	0.42	0
2	GBX	B	530	-	47,47,47	3.12	10 (21%)	69,69,69	1.43	5 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GBX	A	230	-	-	0/27/47/47	0/0/5/5
3	IPA	A	235	-	-	0/0/0/0	0/0/0/0
3	IPA	B	236	-	-	0/0/0/0	0/0/0/0
2	GBX	B	530	-	-	0/27/47/47	0/0/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	530	GBX	C8-C7	-17.98	1.28	1.54
2	A	230	GBX	C8-C7	-17.82	1.28	1.54
2	A	230	GBX	C9-C8	4.49	1.59	1.52
2	B	530	GBX	C9-C8	4.46	1.59	1.52
2	B	530	GBX	C10-C9	4.38	1.60	1.53
2	A	230	GBX	C10-C9	4.21	1.60	1.53
2	B	530	GBX	C1'-C2'	4.19	1.43	1.37
2	A	230	GBX	C2'-C3'	4.18	1.46	1.40
2	A	230	GBX	C1'-C2'	4.09	1.43	1.37
2	B	530	GBX	C2'-C3'	3.70	1.45	1.40
2	B	530	GBX	C7'-C8'	3.27	1.44	1.36
2	A	230	GBX	C9A-C9'	3.18	1.44	1.34
2	A	230	GBX	C7'-C8'	3.16	1.43	1.36
2	B	530	GBX	C9A-C9'	3.14	1.43	1.34
2	B	530	GBX	C7'-C6'	2.83	1.43	1.36
2	A	230	GBX	C5'-C4'	2.80	1.42	1.34
2	B	530	GBX	C5'-C4'	2.79	1.42	1.34
2	A	230	GBX	C7'-C6'	2.76	1.43	1.36
2	A	230	GBX	C2'-C7	2.24	1.52	1.48
2	B	530	GBX	C2'-C7	2.24	1.52	1.48

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	230	GBX	O8-C8-C9	-6.28	111.47	119.76
2	B	530	GBX	O8-C8-C9	-6.23	111.53	119.76
2	B	530	GBX	O7-C7-C2'	-5.76	111.53	121.58
2	A	230	GBX	O7-C7-C2'	-5.66	111.71	121.58
2	B	530	GBX	O7-C7-C8	-4.49	111.33	117.76
2	A	230	GBX	O7-C7-C8	-4.29	111.62	117.76
2	A	230	GBX	O8-C8-C7	-3.71	111.62	118.79
2	B	530	GBX	O8-C8-C7	-3.68	111.67	118.79
2	A	230	GBX	C2'-C7-C8	-3.12	113.09	117.58
2	B	530	GBX	C2'-C7-C8	-3.07	113.16	117.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/221 (99%)	0.26	11 (5%) 28 28	12, 28, 79, 94	0
1	B	220/221 (99%)	0.19	7 (3%) 45 46	15, 29, 60, 98	0
All	All	440/442 (99%)	0.22	18 (4%) 35 36	12, 29, 72, 98	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	221	PHE	6.7
1	B	521	PHE	5.4
1	B	520	LYS	5.3
1	A	112	PRO	4.7
1	A	210	LYS	4.3
1	A	217	LYS	3.3
1	A	220	LYS	3.3
1	A	111	CSO	3.2
1	A	113	PRO	3.1
1	B	518	VAL	3.0
1	B	517	LYS	2.9
1	A	118	ALA	2.8
1	B	302	GLY	2.7
1	B	514	GLU	2.7
1	A	114	ASP	2.6
1	A	208	ASP	2.2
1	B	318	ILE	2.1
1	A	213	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CSO	A	111	7/8	0.22	-0.16	76,80,84,84	0
1	CSO	B	411	7/8	0.12	-1.00	29,36,38,39	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GBX	B	530	43/43	0.22	1.38	23,69,74,79	0
2	GBX	A	230	43/43	0.21	0.96	22,68,75,78	0
3	IPA	B	236	4/4	0.10	-0.06	48,48,51,52	0
3	IPA	A	235	4/4	0.13	-0.24	35,36,37,42	0

6.5 Other polymers

There are no such residues in this entry.