



Full wwPDB X-ray Structure Validation Report

Oct 7, 2014 – 04:47 PM EDT

PDB ID : 4MMO
Title : The crystal structure of a M20 family metallo-carboxypeptidase Sso-CP2 from *Sulfolobus solfataricus*
Authors : Dupuy, J.; Dutoit, R.; Durisotti, V.; Demarez, M.; Borel, F.; Van Elder, D.; Legrain, C.; Bauvois, C.
Deposited on : 2013-09-09
Resolution : 2.34 Å (reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23828
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23828

1 Overall quality at a glance

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	4049 (2.38-2.30)
Clashscore	79885	1094 (2.36-2.32)
Ramachandran outliers	78287	1080 (2.36-2.32)
Sidechain outliers	78261	1081 (2.36-2.32)
RSRZ outliers	66119	4050 (2.38-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	437	
1	B	437	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	SO4	A	502	-	X
3	SO4	B	502	-	X
4	GOL	B	503	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 12814 atoms, of which 6210 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sso-CP2 metallo-carboxypeptidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	437	Total	C	H	N	O	S	0	0	0
			6845	2194	3431	561	650	9			
1	B	382	Total	C	H	N	O	S	0	0	0
			5611	1838	2771	465	530	7			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	188	Total	O	0	0
			188	188		
5	B	130	Total	O	0	0
			130	130		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sso-CP2 metallo-carboxypeptidase



- Molecule 1: Sso-CP2 metallo-carboxypeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.07Å 89.37Å 161.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.10 – 2.34 49.16 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.1 (46.10-2.34) 99.2 (49.16-2.34)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.206 , 0.253 0.206 , 0.254	Depositor DCC
R_{free} test set	2651 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	46.7	Xtriage
Anisotropy	0.510	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 57.3	EDS
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 53076 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12814	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/3479	0.44	0/4706
1	B	0.23	0/2885	0.42	0/3909
All	All	0.24	0/6364	0.43	0/8615

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3414	3431	11	12	0
1	B	2840	2771	0	7	0
2	A	1	0	0	0	0
3	A	15	0	0	0	0
3	B	10	0	0	0	0
4	B	6	8	0	0	0
5	A	188	0	0	2	0
5	B	130	0	0	3	0
All	All	6604	6210	11	18	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 1.

All (18) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:198:SER:OG	1:A:298:THR:OG1	2.09	0.69
1:A:61:ASN:ND2	5:A:705:HOH:O	2.32	0.63
1:A:198:SER:HG	1:A:298:THR:HG1	1.47	0.62
1:B:291:TYR:OH	1:B:294:LYS:O	2.19	0.61
1:A:364:LYS:NZ	1:A:370:GLU:OE1	2.38	0.56
1:A:249:ASP:OD1	1:A:269:LYS:NZ	2.40	0.55
1:B:303:ARG:NE	5:B:629:HOH:O	2.29	0.55
1:A:359:MET:HG2	1:A:398:ALA:HB3	1.94	0.50
1:A:146:ASP:OD1	5:A:677:HOH:O	2.20	0.46
1:A:198:SER:HA	1:A:300:VAL:HG23	1.99	0.45
1:B:24:ILE:N	5:B:627:HOH:O	2.46	0.44
1:B:247:LYS:NZ	5:B:673:HOH:O	2.45	0.42
1:A:179:VAL:HG11	1:A:381:GLN:HB3	2.02	0.41
1:A:118:HIS:ND1	1:A:428:GLU:OE2	2.48	0.41
1:A:366:VAL:HG21	1:A:426:HIS:HA	2.03	0.41
1:A:288:GLU:HB2	1:B:292:THR:HG21	2.02	0.41
1:B:176:VAL:HG13	1:B:397:SER:HB3	2.02	0.40
1:B:349:THR:HG21	1:B:395:ALA:H	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	435/437 (100%)	428 (98%)	6 (1%)	1 (0%)	56	67
1	B	368/437 (84%)	355 (96%)	13 (4%)	0	100	100
All	All	803/874 (92%)	783 (98%)	19 (2%)	1 (0%)	59	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	399	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/374 (97%)	363 (100%)	0	100	100
1	B	277/374 (74%)	275 (99%)	2 (1%)	91	96
All	All	640/748 (86%)	638 (100%)	2 (0%)	96	98

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	168	ASP
1	B	287	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	70	ASN
1	B	105	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	502	-	4,4,4	0.21	0	6,6,6	0.34	0
3	SO4	A	503	-	4,4,4	0.16	0	6,6,6	0.08	0
3	SO4	A	504	-	4,4,4	0.18	0	6,6,6	0.10	0
3	SO4	B	501	-	4,4,4	0.19	0	6,6,6	0.09	0
3	SO4	B	502	-	4,4,4	0.20	0	6,6,6	0.10	0
4	GOL	B	503	-	5,5,5	0.35	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	502	-	-	0/0/0/0	0/0/0/0
3	SO4	A	503	-	-	0/0/0/0	0/0/0/0
3	SO4	A	504	-	-	0/0/0/0	0/0/0/0
3	SO4	B	501	-	-	0/0/0/0	0/0/0/0
3	SO4	B	502	-	-	0/0/0/0	0/0/0/0
4	GOL	B	503	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	437/437 (100%)	0.45	22 (5%)	28 39	31, 47, 80, 131	0
1	B	382/437 (87%)	1.76	125 (32%)	1 1	32, 90, 172, 241	0
All	All	819/874 (93%)	1.06	147 (17%)	2 3	31, 53, 151, 241	0

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	17	ILE	13.9
1	B	39	LEU	11.2
1	B	418	ASP	10.6
1	B	31	LEU	10.6
1	B	398	ALA	10.5
1	B	120	LEU	9.4
1	B	38	LEU	8.6
1	B	167	LEU	8.3
1	B	16	SER	7.6
1	B	421	TYR	7.2
1	B	364	LYS	7.1
1	B	114	PHE	7.0
1	B	111	ALA	6.8
1	B	417	ILE	6.6
1	B	29	ASN	6.6
1	A	405	TYR	6.5
1	B	366	VAL	6.2
1	B	173	PRO	5.9
1	B	9	ILE	5.9
1	B	437	LEU	5.8
1	B	30	TYR	5.8
1	B	365	LYS	5.7
1	B	48	THR	5.6
1	B	399	ILE	5.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	430	PHE	5.5
1	B	436	ILE	5.4
1	B	124	GLU	5.4
1	B	28	ALA	5.3
1	B	115	ALA	5.3
1	B	119	LEU	5.3
1	B	41	VAL	5.1
1	B	45	LEU	5.1
1	B	32	LYS	5.1
1	B	367	TYR	5.0
1	B	354	THR	5.0
1	B	169	PRO	4.8
1	B	113	LEU	4.8
1	B	26	GLU	4.8
1	B	360	ILE	4.8
1	B	431	LEU	4.7
1	B	53	VAL	4.7
1	B	11	PHE	4.7
1	B	43	ALA	4.6
1	B	24	ILE	4.6
1	B	27	THR	4.5
1	B	103	SER	4.5
1	B	368	GLY	4.5
1	B	52	PRO	4.4
1	B	127	VAL	4.4
1	B	126	ASN	4.3
1	B	20	THR	4.3
1	B	10	GLU	4.3
1	B	44	ASN	4.2
1	B	123	ASN	4.2
1	B	171	GLY	4.2
1	A	404	TYR	4.2
1	B	15	PRO	4.2
1	B	106	LYS	4.2
1	B	166	GLY	4.1
1	B	25	ASP	4.1
1	B	71	HIS	4.1
1	B	51	HIS	4.1
1	B	371	PRO	4.1
1	B	387	VAL	4.1
1	B	34	THR	4.0
1	B	91	THR	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	388	TYR	3.9
1	B	88	PHE	3.9
1	B	426	HIS	3.9
1	B	433	LEU	3.8
1	B	125	LEU	3.8
1	A	388	TYR	3.7
1	B	112	ARG	3.7
1	A	392	ILE	3.5
1	B	33	GLU	3.5
1	B	170	LYS	3.4
1	B	386	PHE	3.4
1	A	391	GLY	3.4
1	B	12	LEU	3.3
1	B	36	GLU	3.3
1	B	392	ILE	3.3
1	B	358	ALA	3.3
1	B	422	LYS	3.3
1	B	390	LEU	3.3
1	B	122	LYS	3.3
1	B	391	GLY	3.2
1	B	168	ASP	3.2
1	B	72	TYR	3.2
1	B	434	TYR	3.2
1	B	393	ARG	3.2
1	A	395	ALA	3.1
1	B	121	ASP	3.1
1	B	61	ASN	3.1
1	A	160	VAL	3.0
1	B	172	ARG	3.0
1	B	35	VAL	3.0
1	B	109	LEU	3.0
1	B	296	SER	3.0
1	B	362	SER	3.0
1	B	394	ASP	3.0
1	B	87	PRO	2.9
1	A	387	VAL	2.9
1	B	164	GLY	2.9
1	A	386	PHE	2.9
1	B	427	THR	2.9
1	B	23	GLY	2.9
1	B	161	ILE	2.8
1	B	13	LYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	296	SER	2.8
1	B	118	HIS	2.7
1	A	349	THR	2.7
1	A	345	TYR	2.6
1	B	8	LEU	2.6
1	A	184	TYR	2.6
1	B	104	ASP	2.6
1	B	90	ALA	2.6
1	B	357	LYS	2.6
1	A	390	LEU	2.6
1	B	221	MET	2.5
1	A	347	VAL	2.5
1	B	424	ILE	2.5
1	B	429	GLU	2.4
1	A	291	TYR	2.4
1	B	355	VAL	2.4
1	B	58	ILE	2.4
1	A	406	SER	2.3
1	B	425	LYS	2.3
1	B	105	ASN	2.3
1	A	183	LEU	2.3
1	B	384	GLY	2.2
1	A	346	PRO	2.2
1	B	133	TYR	2.2
1	B	254	GLU	2.2
1	B	37	LYS	2.2
1	A	384	GLY	2.2
1	B	385	LEU	2.2
1	B	108	THR	2.1
1	B	345	TYR	2.1
1	A	383	MET	2.1
1	B	69	TYR	2.1
1	B	347	VAL	2.1
1	B	131	LEU	2.1
1	B	21	GLY	2.1
1	B	356	VAL	2.1
1	B	47	LYS	2.1
1	B	147	TYR	2.1
1	A	393	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	B	502	5/5	0.23	8.56	75,84,94,94	0
3	SO4	A	502	5/5	0.31	3.27	74,99,111,113	0
4	GOL	B	503	6/6	0.18	2.00	36,62,84,90	0
3	SO4	B	501	5/5	0.12	-0.95	51,63,76,78	0
3	SO4	A	503	5/5	0.11	-0.97	64,78,96,98	0
2	ZN	A	501	1/1	0.12	-1.69	31,31,31,31	1
3	SO4	A	504	5/5	0.10	-2.84	58,77,89,91	0

6.5 Other polymers ⓘ

There are no such residues in this entry.