



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 08:24 PM GMT

PDB ID : 1MVK  
Title : X-ray structure of the tetrameric mutant of the B1 domain of streptococcal protein G  
Authors : Frank, M.K.; Dyda, F.; Dobrodumov, A.; Gronenborn, A.M.  
Deposited on : 2002-09-25  
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	56	
1	B	56	
1	C	56	
1	D	56	
1	E	56	
1	F	56	
1	G	56	
1	H	56	
1	I	56	
1	J	56	
1	K	56	
1	L	56	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4718 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Immunoglobulin G binding protein G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	46	372	238	56	77	1	0	0	0
1	B	46	372	238	56	77	1	0	0	0
1	C	46	372	238	56	77	1	0	0	0
1	D	45	363	233	55	74	1	0	0	0
1	E	45	362	232	55	74	1	0	0	0
1	F	46	372	238	56	77	1	0	0	0
1	G	47	376	240	57	78	1	0	0	0
1	H	50	399	254	61	83	1	0	0	0
1	I	48	383	244	58	80	1	0	0	0
1	J	46	372	238	56	77	1	0	0	0
1	K	46	372	238	56	77	1	0	0	0
1	L	46	370	237	56	76	1	0	0	0

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING MET	UNP P06654
A	2	GLN	THR	ENGINEERED	UNP P06654
A	5	VAL	LEU	ENGINEERED	UNP P06654
A	26	PHE	ALA	ENGINEERED	UNP P06654
A	30	VAL	PHE	ENGINEERED	UNP P06654

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	33	PHE	TYR	ENGINEERED	UNP P06654
A	34	PHE	ALA	ENGINEERED	UNP P06654
B	1	MET	-	INITIATING MET	UNP P06654
B	2	GLN	THR	ENGINEERED	UNP P06654
B	5	VAL	LEU	ENGINEERED	UNP P06654
B	26	PHE	ALA	ENGINEERED	UNP P06654
B	30	VAL	PHE	ENGINEERED	UNP P06654
B	33	PHE	TYR	ENGINEERED	UNP P06654
B	34	PHE	ALA	ENGINEERED	UNP P06654
C	1	MET	-	INITIATING MET	UNP P06654
C	2	GLN	THR	ENGINEERED	UNP P06654
C	5	VAL	LEU	ENGINEERED	UNP P06654
C	26	PHE	ALA	ENGINEERED	UNP P06654
C	30	VAL	PHE	ENGINEERED	UNP P06654
C	33	PHE	TYR	ENGINEERED	UNP P06654
C	34	PHE	ALA	ENGINEERED	UNP P06654
D	1	MET	-	INITIATING MET	UNP P06654
D	2	GLN	THR	ENGINEERED	UNP P06654
D	5	VAL	LEU	ENGINEERED	UNP P06654
D	26	PHE	ALA	ENGINEERED	UNP P06654
D	30	VAL	PHE	ENGINEERED	UNP P06654
D	33	PHE	TYR	ENGINEERED	UNP P06654
D	34	PHE	ALA	ENGINEERED	UNP P06654
E	1	MET	-	INITIATING MET	UNP P06654
E	2	GLN	THR	ENGINEERED	UNP P06654
E	5	VAL	LEU	ENGINEERED	UNP P06654
E	26	PHE	ALA	ENGINEERED	UNP P06654
E	30	VAL	PHE	ENGINEERED	UNP P06654
E	33	PHE	TYR	ENGINEERED	UNP P06654
E	34	PHE	ALA	ENGINEERED	UNP P06654
F	1	MET	-	INITIATING MET	UNP P06654
F	2	GLN	THR	ENGINEERED	UNP P06654
F	5	VAL	LEU	ENGINEERED	UNP P06654
F	26	PHE	ALA	ENGINEERED	UNP P06654
F	30	VAL	PHE	ENGINEERED	UNP P06654
F	33	PHE	TYR	ENGINEERED	UNP P06654
F	34	PHE	ALA	ENGINEERED	UNP P06654
G	1	MET	-	INITIATING MET	UNP P06654
G	2	GLN	THR	ENGINEERED	UNP P06654
G	5	VAL	LEU	ENGINEERED	UNP P06654
G	26	PHE	ALA	ENGINEERED	UNP P06654
G	30	VAL	PHE	ENGINEERED	UNP P06654

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	33	PHE	TYR	ENGINEERED	UNP P06654
G	34	PHE	ALA	ENGINEERED	UNP P06654
H	1	MET	-	INITIATING MET	UNP P06654
H	2	GLN	THR	ENGINEERED	UNP P06654
H	5	VAL	LEU	ENGINEERED	UNP P06654
H	26	PHE	ALA	ENGINEERED	UNP P06654
H	30	VAL	PHE	ENGINEERED	UNP P06654
H	33	PHE	TYR	ENGINEERED	UNP P06654
H	34	PHE	ALA	ENGINEERED	UNP P06654
I	1	MET	-	INITIATING MET	UNP P06654
I	2	GLN	THR	ENGINEERED	UNP P06654
I	5	VAL	LEU	ENGINEERED	UNP P06654
I	26	PHE	ALA	ENGINEERED	UNP P06654
I	30	VAL	PHE	ENGINEERED	UNP P06654
I	33	PHE	TYR	ENGINEERED	UNP P06654
I	34	PHE	ALA	ENGINEERED	UNP P06654
J	1	MET	-	INITIATING MET	UNP P06654
J	2	GLN	THR	ENGINEERED	UNP P06654
J	5	VAL	LEU	ENGINEERED	UNP P06654
J	26	PHE	ALA	ENGINEERED	UNP P06654
J	30	VAL	PHE	ENGINEERED	UNP P06654
J	33	PHE	TYR	ENGINEERED	UNP P06654
J	34	PHE	ALA	ENGINEERED	UNP P06654
K	1	MET	-	INITIATING MET	UNP P06654
K	2	GLN	THR	ENGINEERED	UNP P06654
K	5	VAL	LEU	ENGINEERED	UNP P06654
K	26	PHE	ALA	ENGINEERED	UNP P06654
K	30	VAL	PHE	ENGINEERED	UNP P06654
K	33	PHE	TYR	ENGINEERED	UNP P06654
K	34	PHE	ALA	ENGINEERED	UNP P06654
L	1	MET	-	INITIATING MET	UNP P06654
L	2	GLN	THR	ENGINEERED	UNP P06654
L	5	VAL	LEU	ENGINEERED	UNP P06654
L	26	PHE	ALA	ENGINEERED	UNP P06654
L	30	VAL	PHE	ENGINEERED	UNP P06654
L	33	PHE	TYR	ENGINEERED	UNP P06654
L	34	PHE	ALA	ENGINEERED	UNP P06654

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total O S 5 4 1	0	0
2	K	1	Total O S 5 4 1	0	0
2	H	1	Total O S 5 4 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	22	Total O 22 22	0	0
3	B	25	Total O 25 25	0	0
3	C	14	Total O 14 14	0	0
3	D	22	Total O 22 22	0	0
3	E	10	Total O 10 10	0	0
3	F	23	Total O 23 23	0	0
3	G	17	Total O 17 17	0	0
3	H	20	Total O 20 20	0	0
3	I	16	Total O 16 16	0	0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
3	J	13	Total O 13 13	0	0
3	K	19	Total O 19 19	0	0
3	L	17	Total O 17 17	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Immunoglobulin G binding protein G

Chain A: 



- Molecule 1: Immunoglobulin G binding protein G

Chain B: 




- Molecule 1: Immunoglobulin G binding protein G

Chain C: 



- Molecule 1: Immunoglobulin G binding protein G

Chain D: 



- Molecule 1: Immunoglobulin G binding protein G

Chain E: 




- Molecule 1: Immunoglobulin G binding protein G

Chain F: 



- Molecule 1: Immunoglobulin G binding protein G

Chain G: 



- Molecule 1: Immunoglobulin G binding protein G

Chain H: 



- Molecule 1: Immunoglobulin G binding protein G

Chain I: 



- Molecule 1: Immunoglobulin G binding protein G

Chain J: 




- Molecule 1: Immunoglobulin G binding protein G

Chain K: 



- Molecule 1: Immunoglobulin G binding protein G

Chain L: 



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.10Å 210.40Å 55.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50	Depositor
% Data completeness (in resolution range)	(Not available) (30.00-2.50)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.237 , 0.283	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.44	0/378	0.58	0/511
1	B	0.49	0/378	0.60	0/511
1	C	0.53	0/378	0.57	0/511
1	D	0.43	0/369	0.65	0/499
1	E	0.40	0/368	0.56	0/497
1	F	0.39	0/378	0.52	0/511
1	G	0.42	0/382	0.55	0/516
1	H	0.64	0/405	0.76	1/547 (0.2%)
1	I	0.45	0/389	0.61	0/526
1	J	0.43	0/378	0.54	0/511
1	K	0.45	0/378	0.57	0/511
1	L	0.43	0/376	0.57	0/509
All	All	0.46	0/4557	0.59	1/6160 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	20	ALA	N-CA-C	5.42	125.64	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	372	0	348	6	0
1	B	372	0	348	10	0
1	C	372	0	348	13	0
1	D	363	0	342	13	0
1	E	362	0	340	8	0
1	F	372	0	348	4	0
1	G	376	0	351	1	0
1	H	399	0	378	8	0
1	I	383	0	358	16	0
1	J	372	0	348	15	0
1	K	372	0	348	6	0
1	L	370	0	349	9	0
2	D	5	0	0	0	0
2	H	5	0	0	0	0
2	K	5	0	0	0	0
3	A	22	0	0	2	0
3	B	25	0	0	3	0
3	C	14	0	0	1	0
3	D	22	0	0	1	0
3	E	10	0	0	0	0
3	F	23	0	0	0	0
3	G	17	0	0	0	0
3	H	20	0	0	1	0
3	I	16	0	0	2	0
3	J	13	0	0	0	0
3	K	19	0	0	1	0
3	L	17	0	0	1	0
All	All	4718	0	4206	77	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

The worst 5 of 77 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:2:GLN:HB3	1:C:49:THR:HG22	1.75	0.69
1:J:55:THR:O	1:J:55:THR:HG22	1.92	0.69
1:H:20:ALA:HA	3:H:4071:HOH:O	1.93	0.67
1:B:54:VAL:HG12	1:B:56:GLU:H	1.62	0.64
3:I:4189:HOH:O	1:K:54:VAL:HA	1.98	0.63

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	42/56 (75%)	42 (100%)	0	0	100	100
1	B	42/56 (75%)	41 (98%)	1 (2%)	0	100	100
1	C	42/56 (75%)	41 (98%)	0	1 (2%)	9	13
1	D	41/56 (73%)	39 (95%)	2 (5%)	0	100	100
1	E	41/56 (73%)	41 (100%)	0	0	100	100
1	F	42/56 (75%)	41 (98%)	1 (2%)	0	100	100
1	G	43/56 (77%)	41 (95%)	1 (2%)	1 (2%)	10	14
1	H	46/56 (82%)	45 (98%)	1 (2%)	0	100	100
1	I	44/56 (79%)	44 (100%)	0	0	100	100
1	J	42/56 (75%)	39 (93%)	3 (7%)	0	100	100
1	K	42/56 (75%)	40 (95%)	0	2 (5%)	4	3
1	L	42/56 (75%)	42 (100%)	0	0	100	100
All	All	509/672 (76%)	496 (97%)	9 (2%)	4 (1%)	27	46

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	8	ASN
1	K	54	VAL
1	K	55	THR
1	C	20	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	40/48 (83%)	40 (100%)	0	100	100
1	B	40/48 (83%)	40 (100%)	0	100	100
1	C	40/48 (83%)	40 (100%)	0	100	100
1	D	39/48 (81%)	39 (100%)	0	100	100
1	E	39/48 (81%)	39 (100%)	0	100	100
1	F	40/48 (83%)	40 (100%)	0	100	100
1	G	40/48 (83%)	40 (100%)	0	100	100
1	H	43/48 (90%)	42 (98%)	1 (2%)	63	87
1	I	41/48 (85%)	41 (100%)	0	100	100
1	J	40/48 (83%)	40 (100%)	0	100	100
1	K	40/48 (83%)	39 (98%)	1 (2%)	60	85
1	L	40/48 (83%)	40 (100%)	0	100	100
All	All	482/576 (84%)	480 (100%)	2 (0%)	95	99

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	18	THR
1	K	1	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	2	GLN
1	F	35	ASN
1	I	35	ASN
1	D	8	ASN
1	H	8	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	D	105	-	4,4,4	1.26	0	6,6,6	0.06	0
2	SO4	H	107	-	4,4,4	1.19	0	6,6,6	0.10	0
2	SO4	K	106	-	4,4,4	1.27	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	D	105	-	-	0/0/0/0	0/0/0/0
2	SO4	H	107	-	-	0/0/0/0	0/0/0/0
2	SO4	K	106	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers

EDS was not executed - this section will therefore be empty.