



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 12:36 PM GMT

PDB ID : 1MWC
Title : WILD TYPE MYOGLOBIN WITH CO
Authors : Murshudov, G.N.; Krzywda, S.; Brzozowski, A.M.; Jaskolski, M.; Scott, E.E.;
Klizas, S.A.; Gibson, Q.H.; Olson, J.S.; Wilkinson, A.J.
Deposited on : 1998-08-12
Resolution : 1.70 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2456 (1.70-1.70)
Clashscore	79885	2929 (1.70-1.70)
Ramachandran outliers	78287	2878 (1.70-1.70)
Sidechain outliers	78261	2878 (1.70-1.70)
RSRZ outliers	66119	2456 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	153	
1	B	153	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3097 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (MYOGLOBIN).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	2	15	0
			1256	800	219	234	3			
1	B	153	Total	C	N	O	S	7	8	0
			1228	781	213	231	3			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			2	1	1		
3	B	1	Total	C	O	0	0
			2	1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	292	Total	O	0	0
			292	292		
4	B	231	Total	O	0	0
			231	231		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (MYOGLOBIN)

Chain A: 



- Molecule 1: PROTEIN (MYOGLOBIN)

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	122.18Å 42.17Å 91.67Å 90.00° 92.61° 90.00°	Depositor
Resolution (Å)	24.00 – 1.70 24.97 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (24.00-1.70) 92.9 (24.97-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.71Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.189 , 0.237 0.197 , 0.247	Depositor DCC
R_{free} test set	2426 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.397	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.6	EDS
Estimated twinning fraction	0.124 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 24046 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3097	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 100.00 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 0.0000e+00. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.93	0/1347	1.59	19/1799 (1.1%)
1	B	1.04	2/1283 (0.2%)	1.77	29/1718 (1.7%)
All	All	0.99	2/2630 (0.1%)	1.68	48/3517 (1.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	74	GLY	N-CA	5.08	1.53	1.46
1	B	109	GLU	CD-OE2	5.07	1.31	1.25

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	16	LYS	CG-CD-CE	13.62	152.76	111.90
1	B	146	TYR	CB-CG-CD2	-11.88	113.87	121.00
1	B	4	ASP	CB-CG-OD1	11.14	128.33	118.30
1	A	4	ASP	CB-CG-OD1	10.91	128.12	118.30
1	A	4	ASP	CB-CG-OD2	-10.84	108.54	118.30
1	B	103	TYR	CB-CG-CD2	-10.35	114.79	121.00
1	B	46	PHE	CB-CG-CD1	9.43	127.40	120.80
1	B	4	ASP	CB-CG-OD2	-8.47	110.68	118.30
1	B	16	LYS	CD-CE-NZ	8.38	130.97	111.70
1	B	146	TYR	CB-CG-CD1	7.41	125.45	121.00
1	B	31	ARG	NE-CZ-NH2	-7.35	116.62	120.30
1	A	20	ASP	CB-CG-OD1	7.29	124.86	118.30
1	B	123	PHE	CB-CG-CD1	7.08	125.75	120.80
1	A	109[A]	GLU	OE1-CD-OE2	-7.04	114.84	123.30
1	A	109[B]	GLU	OE1-CD-OE2	-7.04	114.84	123.30
1	B	139	ARG	NE-CZ-NH1	6.94	123.77	120.30
1	B	123	PHE	CB-CG-CD2	-6.90	115.97	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	138	PHE	CB-CG-CD2	-6.62	116.16	120.80
1	A	31	ARG	NE-CZ-NH2	-6.44	117.08	120.30
1	A	139	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	B	7	TRP	CG-CD2-CE3	-6.13	128.38	133.90
1	A	88	PRO	N-CA-CB	5.99	110.48	103.30
1	B	2	LEU	O-C-N	5.98	132.27	122.70
1	B	13[A]	VAL	CA-CB-CG2	-5.88	102.09	110.90
1	B	13[B]	VAL	CA-CB-CG2	-5.88	102.09	110.90
1	B	5	GLY	O-C-N	5.85	132.06	122.70
1	B	20	ASP	CB-CG-OD1	5.84	123.56	118.30
1	B	126	ASP	CB-CG-OD2	-5.80	113.08	118.30
1	A	76	LEU	CB-CG-CD1	-5.74	101.25	111.00
1	B	7	TRP	CE3-CZ3-CH2	5.46	127.21	121.20
1	B	70	THR	CA-CB-CG2	-5.41	104.83	112.40
1	A	7	TRP	CE3-CZ3-CH2	5.37	127.11	121.20
1	A	146	TYR	CB-CG-CD2	-5.36	117.78	121.00
1	A	7	TRP	CG-CD2-CE3	-5.35	129.09	133.90
1	B	148	GLU	CA-CB-CG	5.26	124.97	113.40
1	B	7	TRP	CD1-CG-CD2	-5.25	102.10	106.30
1	A	60	ASP	CB-CG-OD2	-5.22	113.60	118.30
1	B	14	TRP	CG-CD1-NE1	-5.21	104.89	110.10
1	B	141	ASP	CB-CG-OD2	-5.17	113.65	118.30
1	B	103	TYR	CB-CG-CD1	5.14	124.09	121.00
1	A	103	TYR	O-C-N	5.14	130.92	122.70
1	A	8[A]	GLN	CA-CB-CG	-5.11	102.15	113.40
1	A	8[B]	GLN	CA-CB-CG	-5.11	102.15	113.40
1	A	7	TRP	CD2-CE3-CZ3	-5.08	112.19	118.80
1	B	55	MET	O-C-N	-5.07	114.59	122.70
1	A	114	VAL	CA-C-O	5.06	130.72	120.10
1	A	31	ARG	NE-CZ-NH1	5.03	122.81	120.30
1	B	46	PHE	CG-CD1-CE1	5.02	126.32	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the

chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1256	0	1282	22	13
1	B	1228	0	1236	13	5
2	A	43	0	30	4	0
2	B	43	0	30	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	292	0	0	12	11
4	B	231	0	0	9	3
All	All	3097	0	2578	38	17

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (38) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:128[B]:GLN:OE1	4:B:225:HOH:O	1.71	1.06
1:B:128[B]:GLN:OE1	4:B:214:HOH:O	1.76	1.02
1:B:128[B]:GLN:CD	4:B:225:HOH:O	2.07	0.92
1:B:128[B]:GLN:CG	4:B:225:HOH:O	2.17	0.91
1:B:128[B]:GLN:HG2	4:B:225:HOH:O	1.73	0.88
1:B:8:GLN:NE2	4:B:238:HOH:O	2.16	0.78
1:A:27:GLU:HG3	4:A:320:HOH:O	1.91	0.70
1:A:45:LYS:O	1:A:48[B]:HIS:NE2	2.25	0.68
2:A:154:HEM:HBC2	2:A:154:HEM:HMC1	1.75	0.67
1:A:63:LYS:HG3	4:A:434:HOH:O	1.96	0.65
1:A:120:PRO:HB3	4:A:285:HOH:O	1.97	0.64
1:B:27[A]:GLU:HG3	4:B:179:HOH:O	2.00	0.61
1:A:26:GLN:NE2	4:A:281:HOH:O	2.34	0.60
1:B:52:GLU:HG2	1:B:56:LYS:HE2	1.87	0.56
1:B:59:GLU:HG2	1:B:63:LYS:HE3	1.87	0.55
1:B:44:ASP:HA	1:B:47:LYS:HD2	1.89	0.53
1:A:96[B]:LYS:HE2	4:A:289:HOH:O	2.09	0.52
1:A:43:PHE:O	1:A:47[A]:LYS:HD2	2.11	0.51
1:A:101:VAL:HG13	4:A:269:HOH:O	2.09	0.51
1:A:44:ASP:HA	1:A:47[A]:LYS:HD3	1.93	0.50
1:A:89:LEU:HD11	2:A:154:HEM:CHB	2.42	0.50
1:B:31:ARG:NH1	4:B:205:HOH:O	2.40	0.49
1:B:128[B]:GLN:CD	4:B:214:HOH:O	2.36	0.49
1:A:26:GLN:NE2	4:A:320:HOH:O	2.20	0.48
2:A:154:HEM:HBC2	2:A:154:HEM:CMC	2.41	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:48[A]:HIS:NE2	1:A:49:LEU:HD21	2.31	0.46
1:A:145[B]:LYS:HG2	4:A:221:HOH:O	2.16	0.45
1:A:96[A]:LYS:NZ	4:A:194:HOH:O	2.49	0.44
1:B:99:ILE:HD12	2:B:154:HEM:CAC	2.49	0.43
1:A:31:ARG:HA	1:A:31:ARG:HD2	1.85	0.43
1:A:47[B]:LYS:HB2	4:A:351:HOH:O	2.19	0.41
1:A:99:ILE:HD12	2:A:154:HEM:CAC	2.51	0.41
1:A:47[A]:LYS:HB2	4:A:351:HOH:O	2.20	0.40

All (17) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:128[B]:GLN:CD	4:A:241:HOH:O[2.656]	1.22	0.98
1:A:128[B]:GLN:OE1	4:A:241:HOH:O[2.656]	1.22	0.98
1:A:128[B]:GLN:NE2	4:A:189:HOH:O[2.656]	1.36	0.84
1:A:128[A]:GLN:OE1	4:A:202:HOH:O[2.656]	1.41	0.79
1:A:128[B]:GLN:CD	4:A:202:HOH:O[2.656]	1.47	0.73
1:A:128[B]:GLN:CG	4:A:202:HOH:O[2.656]	1.49	0.71
1:A:48[A]:HIS:CD2	1:B:48:HIS:CD2[1.556]	1.61	0.59
1:A:128[B]:GLN:NE2	4:A:202:HOH:O[2.656]	1.70	0.50
1:A:48[A]:HIS:CD2	1:B:48:HIS:NE2[1.556]	1.81	0.39
1:A:128[B]:GLN:CG	4:A:241:HOH:O[2.656]	1.84	0.36
1:A:48[A]:HIS:NE2	1:B:48:HIS:CD2[1.556]	1.90	0.30
1:A:128[A]:GLN:CD	4:A:202:HOH:O[2.656]	1.96	0.24
1:B:109:GLU:OE1	4:B:324:HOH:O[2.645]	2.01	0.19
1:B:109:GLU:OE2	4:B:324:HOH:O[2.645]	2.04	0.16
4:B:355:HOH:O	4:B:382:HOH:O[2.655]	2.14	0.06
1:A:8[A]:GLN:OE1	4:A:396:HOH:O[2.646]	2.16	0.04
4:A:204:HOH:O	4:A:367:HOH:O[2.656]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	166/153 (108%)	156 (94%)	8 (5%)	2 (1%)	19	3
1	B	159/153 (104%)	151 (95%)	8 (5%)	0	100	100
All	All	325/306 (106%)	307 (94%)	16 (5%)	2 (1%)	50	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48[A]	HIS
1	A	48[B]	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	138/124 (111%)	134 (97%)	4 (3%)	55	30
1	B	131/124 (106%)	129 (98%)	2 (2%)	76	60
All	All	269/248 (108%)	263 (98%)	6 (2%)	68	43

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47[A]	LYS
1	A	47[B]	LYS
1	A	102	LYS
1	A	152	GLN
1	B	145	LYS
1	B	152	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	140	ASN
1	B	8	GLN
1	B	26	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	91	GLN
1	B	116	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	154	1,3	49,50,50	2.90	15 (30%)	46,82,82	2.55	20 (43%)
3	CMO	A	155	2	0,1,1	0.00	-	0,0,0	0.00	-
2	HEM	B	154	1,3	49,50,50	4.58	19 (38%)	46,82,82	3.49	17 (36%)
3	CMO	B	155	2	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	154	1,3	-	0/14/114/114	0/0/8/8
3	CMO	A	155	2	-	0/0/0/0	0/0/0/0
2	HEM	B	154	1,3	-	0/14/114/114	0/0/8/8
3	CMO	B	155	2	-	0/0/0/0	0/0/0/0

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	154	HEM	C3D-C4D	19.88	1.49	1.44
2	B	154	HEM	C2D-C1D	13.49	1.47	1.44
2	B	154	HEM	C2B-C1B	13.40	1.47	1.44
2	A	154	HEM	C3D-C4D	10.66	1.47	1.44
2	A	154	HEM	C2B-C1B	6.71	1.46	1.44
2	B	154	HEM	C3B-C2B	-6.63	1.32	1.43
2	A	154	HEM	C3D-C2D	-6.37	1.32	1.43
2	B	154	HEM	C3D-C2D	-5.91	1.33	1.43
2	A	154	HEM	C3C-C2C	-5.86	1.33	1.43
2	B	154	HEM	C3C-C2C	-5.76	1.33	1.43
2	A	154	HEM	C3B-C2B	-5.56	1.34	1.43
2	B	154	HEM	C3B-CAB	4.82	1.55	1.40
2	A	154	HEM	C3C-CAC	4.81	1.55	1.40
2	A	154	HEM	C3B-CAB	4.63	1.55	1.40
2	A	154	HEM	C4A-C3A	4.09	1.45	1.40
2	B	154	HEM	C3C-CAC	4.09	1.53	1.40
2	B	154	HEM	C4A-C3A	3.47	1.44	1.40
2	B	154	HEM	CMB-C2B	3.38	1.57	1.47
2	B	154	HEM	CHC-C1C	3.16	1.42	1.36
2	A	154	HEM	CHD-C4C	3.06	1.41	1.36
2	A	154	HEM	CMD-C2D	2.93	1.56	1.47
2	B	154	HEM	C1B-NB	-2.88	1.33	1.39
2	A	154	HEM	CMB-C2B	2.65	1.55	1.47
2	B	154	HEM	CMD-C2D	2.57	1.55	1.47
2	B	154	HEM	CMC-C2C	2.57	1.55	1.47
2	B	154	HEM	C3B-C4B	2.49	1.47	1.44
2	A	154	HEM	C2D-C1D	-2.45	1.43	1.44
2	A	154	HEM	CMC-C2C	2.37	1.54	1.47
2	A	154	HEM	C3B-C4B	2.32	1.47	1.44
2	B	154	HEM	FE-NB	2.26	2.06	1.97
2	A	154	HEM	C1A-NA	2.23	1.40	1.36
2	B	154	HEM	C1A-NA	2.12	1.40	1.36
2	B	154	HEM	C1C-NC	2.11	1.41	1.38
2	B	154	HEM	C4D-ND	-2.10	1.35	1.39

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	154	HEM	C3B-C4B-NB	-11.63	105.68	114.00
2	A	154	HEM	C3B-C4B-NB	-8.25	108.10	114.00
2	B	154	HEM	CHD-C1D-ND	8.18	131.38	124.58
2	B	154	HEM	CHC-C4B-NB	7.88	131.13	124.58
2	B	154	HEM	C1B-NB-C4B	6.73	112.05	105.16
2	B	154	HEM	C4D-ND-C1D	6.25	111.56	105.16
2	B	154	HEM	C2D-C1D-ND	-5.84	106.03	112.93
2	B	154	HEM	C1A-CHA-C4D	-5.15	120.70	127.47
2	A	154	HEM	CHC-C4B-NB	5.07	128.79	124.58
2	A	154	HEM	C2D-C1D-ND	-4.99	107.03	112.93
2	A	154	HEM	C4D-ND-C1D	4.46	109.72	105.16
2	A	154	HEM	C1B-NB-C4B	4.10	109.36	105.16
2	B	154	HEM	C4A-CHB-C1B	-4.00	122.21	127.47
2	B	154	HEM	C3A-C4A-NA	3.98	112.42	109.41
2	B	154	HEM	C4C-NC-C1C	-3.81	101.58	105.53
2	B	154	HEM	CHB-C1B-NB	3.76	129.47	124.31
2	A	154	HEM	CMA-C3A-C4A	-3.67	122.97	128.62
2	A	154	HEM	C1A-CHA-C4D	-3.61	122.73	127.47
2	B	154	HEM	C4A-NA-C1A	-3.41	102.27	106.76
2	B	154	HEM	CHA-C4D-ND	3.26	128.78	124.31
2	A	154	HEM	CHA-C4D-ND	3.22	128.73	124.31
2	A	154	HEM	CHB-C1B-NB	3.19	128.69	124.31
2	A	154	HEM	CBA-CAA-C2A	-3.19	107.07	112.69
2	A	154	HEM	C3A-C4A-NA	3.05	111.71	109.41
2	B	154	HEM	CMA-C3A-C4A	-3.03	123.97	128.62
2	A	154	HEM	O2A-CGA-CBA	2.87	124.34	114.22
2	A	154	HEM	C4A-NA-C1A	-2.86	102.99	106.76
2	B	154	HEM	CMB-C2B-C3B	2.84	132.86	126.16
2	A	154	HEM	CMA-C3A-C2A	2.66	129.96	124.94
2	B	154	HEM	CHD-C4C-NC	-2.56	122.50	124.73
2	A	154	HEM	CHD-C4C-NC	-2.55	122.52	124.73
2	B	154	HEM	CMA-C3A-C2A	2.45	129.56	124.94
2	A	154	HEM	C4A-CHB-C1B	-2.29	124.45	127.47
2	A	154	HEM	CBD-CAD-C3D	-2.23	109.51	114.37
2	A	154	HEM	CAD-CBD-CGD	2.13	120.13	113.48
2	A	154	HEM	O1A-CGA-CBA	-2.09	115.83	123.03
2	A	154	HEM	O1D-CGD-CBD	-2.02	116.08	123.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	153/153 (100%)	0.12	5 (3%)	44 49	13, 25, 42, 53	1 (0%)
1	B	153/153 (100%)	0.09	4 (2%)	53 58	13, 23, 38, 44	2 (1%)
All	All	306/306 (100%)	0.11	9 (2%)	49 54	13, 24, 41, 53	3 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	153	GLY	7.3
1	A	152	GLN	3.1
1	B	107	ILE	3.0
1	B	152	GLN	2.9
1	B	1	GLY	2.5
1	A	153	GLY	2.2
1	A	48[A]	HIS	2.1
1	A	72	LEU	2.1
1	A	111	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CMO	A	155	2/2	0.11	-0.09	17,17,17,22	0
2	HEM	B	154	43/43	0.10	-0.10	13,19,30,39	0
2	HEM	A	154	43/43	0.11	-0.28	14,23,29,37	0
3	CMO	B	155	2/2	0.07	-1.00	19,19,19,22	0

6.5 Other polymers ⓘ

There are no such residues in this entry.